

APTA Citros Sylvio Moreira, Instituto Agronômico de Campinas, Cordeirópolis, SP, Brazil; ⁹Centro de Protección Vegetal y Biotecnología, IVIA, Apartado Oficial, 46113-Moncada, Valencia, Spain. fgmitter@ufl.edu

One of the main goals of the International Citrus Genome Consortium (ICGC) is to have a freely available, high quality reference genome sequence for *Citrus*. It was concluded that the best assembled sequence would be derived from the most homozygous genome available, and such materials were sought from the international citrus research community. Tissue and DNA samples from three homozygous plants of *Citrus clementina* Hort. ex Tan., cv. Nules, obtained by gynogenesis or by pollen embryogenesis and freely provided by researchers in Spain, France and Italy, were analyzed and characterized by laboratories in several institutions worldwide employing various methods. Specifically, chromosome numbers were determined, and the evaluations of homozygosity and of integrity of the genome through genomic and EST-derived SSR markers, and using microarray technology, were carried out. Plant materials from Spain and France were determined to be haploid, and the accession from Italy was found to be trihaploid, by chromosome counts. Nearly 180 SSR markers were selected, in many cases from previous mapping exercises, to represent as broad and unbiased coverage of the citrus genome as possible, and plant materials were genotyped. Only one SSR locus revealed anomalous results in the French haploid, while the remainder revealed only a single allele product at all other loci surveyed. The microarray results using the Spanish cDNA array revealed anomalous results from only 7 genes of the more than 20, 000 represented on the array. The results obtained by the different screening teams regarding the first step of the *Citrus* genome sequencing project will be reported in detail.

[101]

EST-SSRs in Citrus: Features, Polymorphism and Mappability

Luro F¹, Costantino G¹, Traore S¹, Demonjuz C¹, Albertini MV¹, Terol J², Talon M², Argout X³, Froelicher Y⁴, Ollitrault P⁴, and Morillon R⁴

¹INRA, Unité de Recherche GEQA, INRA San Giuliano, 20230 San Nicolao, France; ²Centro de Genómica, Instituto Valenciano de Investigaciones Agrarias, Valencia, Spain; ³CIRAD AMIS, Montpellier, France; ⁴UPR 'Amélioration génétique d'espèces à multiplication végétative', CIRAD -Montpellier France. luro@corse.inra.fr

From a clementine EST sequences dataset we investigated the presence, occurrence and repartition of different type of SSR in EST and compared the results obtained with other plant species. Among 11000 unigenes

from 37000 ESTs, we have found at least one microsatellite sequence (repeated units size ranged from 2 to 6 nucleotides) in 1500 unigenes (13.6%). This frequency of ESR-SSR (one in 5.2 kb) seems higher than in other species such as rice, barley, wheat etc. Nevertheless, the observed differences can be partially explained by the number of repeat units considered as limit for SSR selection in computational screening. As for many plant species, di and trinucleotides are dominant in citrus ESTs (88% of total SSRs). Trinucleotides are the most abundant (52%) and hexanucleotides, only 2% being under represented when compared to other plants. Dinucleotide microsatellites were preferentially (50%) concentrated in the 5' 100th nucleotides and a large majority (75%) is positioned in untranslated regions (UTR), since trinucleotide microsatellites were equitably distributed inside and outside the translated region (TR) of the ESTs (48 % and 52% respectively). To optimize the use of EST-SSR we assessed their polymorphism among a citrus species set (50 genotypes) representing the genetic diversity. More than 90% of EST-SSR markers were polymorphic. Furthermore, dinucleotide microsatellite markers were more polymorphic than trinucleotide ones, probably related to their distribution that was more often located in the 5'-UTR. We obtained a good agreement of diversity relationships between the citrus species and relatives assessed with EST-SSR markers with the established taxonomy and phylogeny. To end, the heterozygosity of each genotype and all dual combinations were studied to evaluate the percentage of mappable markers. Higher values (> 45%) were observed for putative *Citrus* inter-specific hybrids (lime lemon, or sour orange) than for *Citrus* basic true species (mandarin, pummelo and citron) (<30%). Most favorable combinations for genome mapping were observed in those involving interspecific hybrid genotypes. Those gave higher levels of mappable markers (>70%) with a significant proportion suitable for synteny analysis. Whatever the position of the SSR in the ESTs, the EST-SSR markers are powerful to investigate genetic diversity and genome mapping in citrus.

[102]

Transcriptome Analysis of a Spontaneous Red-flesh Mutant in Sweet Orange (*Citrus sinensis* [L.] Osbeck) During Fruit Development

Liu Q, Zhu AD, Chai LJ, Zhou WJ, Yu KQ, Ding J, Xu J, and Deng XX

National Key Laboratory of Crop Genetic Improvement, Huazhong Agricultural University, Wuhan 430070, China, xxdeng@mail.hzau.edu.cn

Bud mutations arise often in citrus. The selection of mutants is one of the most important breeding channels in citrus. However, the molecular basis of bud mutation has rarely been studied. To identify potential important or