



BOOK OF ABSTRACTS TALKS

Island BIOLOGY

La Réunion
8-13 JULY

2019

📍 **Université de la Réunion**
Campus du Moufia



Island Biology

BOOK OF ABSTRACTS

TALKS

**Third International Conference
on Island Ecology, Evolution and Conservation**

**8-13 July 2019
University of La Réunion
Saint Denis, France**

Editors: Olivier Flores, Claudine Ah-Peng, Nicholas Wilding

Towards a comprehensive understanding of *Oxera*'s island life

Gildas Gâteblé * ¹, Laure Barrabé ^{1,2}, Philippe Birnbaum ^{1,3}, Bryan Drew ⁴, Yuji Isagi ⁵, Ayako Izuno ⁵, Giliane Karnadi-Abdelkader ¹, Ryota Kawai ⁵, Sébastien Lavergne ⁶, Yoshihisa Suyama ⁷

¹ Institut Agronomique néo-Calédonien (IAC) – Equipe ARBOREAL, BP 711, 98810 Mont-Dore, New Caledonia

² Endemia – 3 rue Jules Harbulot, 98800 Nouméa, New Caledonia

³ UMR AMAP, Université de Montpellier, CIRAD, CNRS, INRA, IRD (AMPA) – Centre de coopération internationale en recherche agronomique pour le développement [CIRAD] – UMR AMAP, Université de Montpellier, CIRAD, CNRS, INRA, IRD, 34398 Montpellier, France

⁴ University of Nebraska-Kearney – Department of Biology, University of Nebraska-Kearney, Kearney, NE 68849, United States

⁵ Kyoto University – Laboratory of Forest Biology Division of Forest and Biomaterials Science, Graduate School of Agriculture, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

⁶ Laboratoire d'Ecologie Alpine (LECA) – Centre National de la Recherche Scientifique - CNRS – Laboratoire d'Ecologie Alpine, CNRS - Université Grenoble Alpes, UMR 5553, 38000 Grenoble, France, France

⁷ Tohoku University – Field Science Center, Graduate School of Agricultural Science, Tohoku University, 232-3 Yomogida, Naruko-onsen, Osaki, Miyagi 989-6711, Japan

Since the introduction of the island life concept by Wallace, isolated oceanic and relatively young islands have been the place to be to study and unravel evolutionary patterns among original groups within the tree of life. In these regards, the quite old New Caledonian archipelago has been relatively neglected despite hosting a world-class biome, perhaps because no definitive cases of adaptive radiation have been detected there thus far. Here we present a detailed analysis of the radiation of the genus *Oxera* (Lamiaceae) based on a multidisciplinary study including domestication, interspecific hybridization, karyology, anatomy, systematics, phylogenetics, pollination biology, ecology, conservation and population genetics. New Caledonia is the main center of *Oxera* diversification with about 33 endemic species plus a few more taxa in Papua New Guinea, Australia and Pacific islands. We have shown that *Oxera* originated from a single and quite recent dispersal event in New Caledonia about 4.5 million years ago, and has successfully established and diversified at an outstanding net rate of up to one species per million years. Establishment and rapid initial diversification of *Oxera* in New Caledonia was probably facilitated by arid conditions that may have created new biota at the beginning of the Pliocene. Concurrently, *Oxera* also underwent major shifts in pollination syndromes, dispersal modes and life forms that are now exhibited by its extant major clades. More recently, the drivers of speciation within the main clades are most likely due to allopatry and habitat shifts in relation to climatic oscillations and soil morphogenesis. Ongoing population genetic studies on several species from different clades (e.g. within the monocaulous baladica clade and especially on the *Oxera baladica* species complex) are highlighting more recent stories, while the complete genome sequencing of a taxon (*Oxera pulchella* subsp. *grandiflora*) is likely to deepen our understanding of the past demographic history of several species and/or clades. *In fine*, we hope the results will be of interest for the conservation of the most critically endangered *Oxera* taxa.

Keywords: *Oxera*, New Caledonia, Lamiaceae, radiation, niche shifts, demographic history

*Speaker