MRS. UGO CHIJIOKE (Orcid ID : 0000-0001-6983-5215)

Article type : Original Manuscript

# QUALITY ATTRIBUTES OF *FUFU* IN SOUTH-EAST NIGERIA: GUIDE FOR CASSAVA BREEDERS

### Running title: Quality Attributes of *fufu*

Ugo Chijioke<sup>1\*</sup>,Tessy Madu<sup>1</sup>, Benjamin Okoye<sup>1</sup>, Amaka Promise Ogunka<sup>1</sup>, Mercy Ejechi<sup>1</sup>, Miriam Ofoeze<sup>1</sup>, Chukwudi Ogbete<sup>1</sup>, Damian Njoku<sup>1</sup>, Justin Ewuziem<sup>1</sup>, Kalu Confidence<sup>1</sup>, Nnaemeka Onyemauwa<sup>1</sup>, Blessing Ukeje<sup>1</sup>, Oluchi Achonwa<sup>1</sup>Lora Forsythe<sup>2</sup>, Geneviève Fliedel<sup>3,4</sup> and Chiedozie Egesi<sup>1,5,6</sup>

<sup>1</sup>National Root Crops Research Institute (NRCRI), Umudike, Umuahia, Nigeria <sup>2</sup>Natural Resources Institute, University of Greenwich, Central Avenue, Chatham Maritime, Kent ME4 4TB, United Kingdom

<sup>3</sup> CIRAD, UMR QUALISUD, F-34398 Montpellier, France

<sup>4</sup> Qualisud, Univ Montpellier, CIRAD, Montpellier SupAgro, Univd'Avignon, Univ de La Réunion, Montpellier, France

<sup>5</sup>International Institute of Tropical Agriculture (IITA), Oyo Road, Ibadan, Nigeria <sup>6</sup>Cornell University, Department of Global Development, Ithaca, NY 14853, USA

The peer review history for this article is available at https://publons.com/publon/10.1111/ijfs.14875

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the <u>Version of Record</u>. Please cite this article as <u>doi:</u> 10.1111/ijfs.14875

\* Corresponding Authors' email: ugochijioke4@gmail.com

### Summary

*Fufu* is a popular traditional fermented wet paste food product from cassava. We examined consumer preferences and quality attributes of *fufu* in Abia and Imo States of South-East Nigeria, with special attention to gender differences, for the purpose of providing guidance to breeders. Data were analysed by the use of descriptive and inferential statistics. Participants for the interview were randomly selected from a list of farmers in the study area. Individual (II) interviews were conducted among 80 participants comprising 26 males (32.5%) and 54 females (67.5%). Preferences along the food chain from raw roots to final product were also obtained. Major traits influencing gender-specific consumer preferences are related to appearance, texture and smell. Smoothness, not sticky, easy to swallow and drawability of *fufu* appear to be major traits that drive acceptance by both men and women. Big roots and smooth skin are prioritized for raw material. Some quality characteristics are conditioned largely by variety traits, while others can be modified by adjusting the processing methods. The complexity of producing high quality *fufu* makes it imperative to introduce a multidisciplinary approach into breeding programmes.

**Keywords:** *fufu*, quality characteristics, varieties, gender, preferred traits, consumers, breeding, processing

#### Introduction

Nigeria is the world's largest producer of cassava, hosting a diverse array of cassava farmers and processors, with the large majority being small-scale operators (Forsythe *et al.* 2016). Major cassavabased food products consumed in Nigeria include: *gari, fufu* and *lafun*. Most of these products are made and consumed locally by farming households themselves (IITA 2012). Fufu is a traditional Nigerian fermented food product in southern, western and eastern Nigeria and some other parts of West Africa (Rosales-soto *et al.* 2016). It is usually described as a 'wet paste food product' and ranks second after *gari* as a food product from cassava (http://www.cassavabiz.org/postharvest/fufu). In some parts of Nigeria, it is also called *utaraakpu* (Owolarafe *et al.* 2018,). Uche (2016) reported that

*fufu* has higher profit, gross margin, mark-up and a better monetary prospect. According to Sanni (1998), late in the 20<sup>th</sup>Century, 60% of all the cassava harvested across Nigeria was used in processing *fufu*, and only 5% for *gari*. However, the preference and consumption pattern has been reversed between *fufu* and *gari* in recent times, because of *fufu*'s poor shelf life and tedious processing methods. *Fufu* is ranked next to *gari* as an indigenous food in Southern Nigeria (Egwim *et al.* 2013), and it is popular in many parts of West Africa (Uyoh *et al.* 2009). *Fufu* flour as a convenience food and staple is increasingly becoming very popular in West Africa (Johnson *et al.* 2006).

*Fufu* is usually processed by households and rural processors whose practices may differ by culture, and region. Fermentation is a key component of *fufu* production, an important step to detoxify the cassava pulp (i.e. degrade cyanogenic glucosides), develop the characteristic aroma and flavour of the *fufu*, and also help in preserving it (Flibert *et al.* 2016). *Fufu* is produced by first, peeling and washing the cassava roots, and cutting them into smaller chunks. The method of soaking/ steeping of roots differ among States and processors in South-East Nigeria. Soaking, steeping or fermentation of the cut roots may be carried out either by continuous soaking of chunked roots for a period of 3-5days of fermentation (Mokemiabeka et al. 2011), or by washing and grating of the soaked root after 48hrs of fermentation, followed by re-steeping of grated roots (Omodamiro et al. 2011). The fermented roots or mash are finally sieved, and dewatered to obtain the wet paste. The sour taste, flavour, appearance and texture are mainly recognized as determinants of *fufu* acceptance and quality (Bamidele et al. 2015). The variations in processing methods and differences in physico-chemical properties of cassava varieties alter the texture and organoleptic properties of the cooked *fufu* (Akingbala *et al.*) 1991). Furthermore, Beechof et al. (2018) and Asrat et al. (2010) noted that gender specific crop trait preferences are rarely considered, or prioritized in most breeding programs. These complexities involved in the processing of the product (*fufu*) makes it imperative for the need to introduce a multidisciplinary approach for breeding varieties that meet end-user needs for *fufu*. The study therefore described the quality characteristics that drive purchase and utilization of fresh cassava roots for *fufu* processing, preferred quality characteristics of cassava root for processing of fermented wet *fufu* 

mash, cooking properties of the intermediate *fufu* product, and quality characteristics of *fufu* during consumption

# Materials and Methods

# Study area

The study was conducted in Imo and Abia States in South East region of Nigeria, selected based on intensity of cassava production (Fig. 1). Imo and Abia States are made up of 3 Agricultural Zones each. Imo State is divided into 27 administrative units called Local Government Areas (LGAs), which are grouped into 3 Agricultural Zones of Owerri, Okigwe and Orlu, and 17 LGAs for Abia State (grouped into 3 Agricultural Zones of Aba, Ohafia and Umuahia). The climate can generally be described as tropical-in the humid rainforest agro-ecological zone-with clearly defined wet and dry seasons. Small-holder crop and livestock farming is the predominant occupation of the people (Anderson *et al.* 2017). Generally, a smallholder farmer is involved in cultivating a small piece of land, cultivating food crops, sometimes with small varieties of cash crops (Thorpe & Muriuki, 2001; Herrero *et al.* 2014). In many localities, smallholder farmers practice mixed crop-livestock farming, whereby the number of large ruminants kept is around 3-5 (Thorpe & Muriuki, 2001). The major crops produced include: yam, cassava, rice, maize, cocoyam, cowpea and tomatoes. *Fufu* is a principal food staple in Imo and Abia States.

# Sampling procedure

A multi-stage sampling procedure was used to select sample respondents to identify traits for a high quality crop (cassava) and product (*fufu*), among the cassava producing and consuming households in the region (Forsythe *et al.* 2020). In the first stage, three (3) Agricultural Zones in Imo State and one (1) in Abia State were selected for the study based on level of fufu processing and consumption. In the second stage, four (4) communities in Imo and two (2) in Abia States were selected, also based on intensity of fufu processing and convenience. In the last stage, participants for the interview were randomly selected from a list of farmers provided by the Agricultural Development Programme (ADP) in the communities. ADPs aims at increasing food production for rural dwellers, and raising the income level of small scale farmers, by making provision for

improved seeds, fertilizer, pesticides, credit facilities and infra-structural facilities (Ajayi and Ajala 1999). Individual interviews (II) were conducted, comprising 10 participants in each community of Imo State (giving 40 participants in four communities) and 20 in each community (40 participants) of Abia State, giving a total of 80 participants (54 female and 26 male).

# Data collection and analysis

The data for the study were collected from primary sources by the use of a well-structured questionnaire. The questionnaires were designed to collect a range of information, including household structure, crop production, sales, utilization and consumption of cassava with special emphasis on *fufu*. The Pivot Table function available in Microsoft Excel was used for analysis. The Pivot Table feature is a user-friendly and easy-to-use tool, which is relevant for the kinds of analysis which is required in a trait preference evaluation, following FAO (2016). Characteristics of preferred traits were assigned weights according to how they were prioritised and the importance they were given by respondents following Forsythe *et al.* (2020). To aggregate the characteristics into one table, we gave each of them weights according to how they were prioritized. Comparison of different priorities for men and women, were also depicted to see how the important characteristics differed. To apply weights, frequency (count) for the most important characteristic (1<sup>st</sup> priority) was multiplied by 3, the frequencies for the second priority characteristic by 2, and the frequencies for the third priority characteristic by 3. The results are weighted scores summarised and ranked as:

 $Y = \sum_{k=i}^{n} (N1x3) + (N2x2) + (N3x1)....(1)$ Where:

Y = Summation of weighted scores

N1= total number of respondents indicating highest (3) preferred traits

N2 = total number of respondents indicating second (2) most preferred traits

N3 = total number of respondents indicating third (1) most preferred traits

For each characteristic, total 'points' from men and women were summed up to get a final score and characteristics sort by descending final scores, and ranked. Results were then presented in tables, graphs, and flow diagrams from which inferences were drawn.

## **Results and Discussion**

# Preferred and less-preferred root quality traits that drive selection of cassava varieties for fufu processing

The evaluation of the preferred and less-preferred quality characteristics that drive purchase and utilization of fresh cassava roots for *fufu* processing within South East Nigeria are shown in Figures 2 and 3. The results show divergent views in choice of trait preferences among the male and female participants. Four root quality attributes were identified as the main traits of preference, and they include: root size (moderately-sized roots equivalent to 1 litre capacity bottled water in diameter), heaviness of the root (weight/density of root when held by the hand), appearance/smoothness of root skin (dark coloured peel and roots without rough/wrinkle skin), and colour of root flesh (white coloured root flesh; without dark discoloration). In contrast, bad root colour (dark/ multiple-coloured striped roots), light foamy weight roots (bread-like in texture), small-sized roots, fibrous roots and roots with high moisture content were identified as less-preferred traits that result to cooked *fufu* of low quality.

Size of root appears to be an important raw material (cassava root) trait for both male and female respondents. It was observed that although male and female respondent's assigned high weight (sum of scores of 48 and 69, respectively) to root size, reason(s) for this preference differs by gender (Fig. 2). According to the female respondents, the preferred moderate-size roots were easy to peel, saved operation time and mitigated drudgery (Supplementary Table 1). This is in agreement with the study by Egbeocha *et al.* (2016) and Jimoh *et al.* (2016), who stated that sizes of tubers is one of the factors that is responsible for the demanding nature of peeling as a unit operation during processing. Furthermore Jimoh and Olukunle (2012) reported that peeling efficiency was higher for larger sized cassava roots compared to smaller ones. According to the respondents, small sized roots are associated with inefficient peeling operations, resulting in peel loss and dark coloured intermediate and final product (cooked fufu). This study therefore reveals that selection of root size as the most important root quality attribute for the female cassava root buyers in Abia and Imo States is based on its influence on processing operation such as peeling and colour of derived products (intermediate and cooked *fufu*). On the contrary, the choice of root size by the male respondents seemed to be driven by

direct economic gain obtained by buying the fresh roots, an activity where men are the major key actors (Bentley *et al.* 2017). According to Gonzáles and Johnson (2009), size of cassava is a key determinant of its market price. The author reported that the bigger the root size, the higher the market price. Interestingly, producing cassava with preferred root traits do not imply an increase in production cost (Gonzáles and Johnson 2009). Our findings are in agreement with the study of Teekan *et al.* (2018), who indicated that gender roles are a strong determinant of preference for traits in cassava.

Heaviness is another trait driving the adoption of cassava varieties for *fufu* processing in South-East Nigeria; and may be related to root weight and density. Gender differences were also observed in the sum scores for this trait, females (51) showed higher interest in the trait compared to their male (10) counterparts. The respondents described the term 'heaviness' with morphological features as seen in yam varieties that have stout and sturdy shape (see Supplementary Table 1). The results show that the preferred trait 'heavy root' can be compared to a less preferred trait 'roots that do not have weight', referring to light weight root, which was described by respondents as cassava roots with bread-like texture. These traits [heaviness or light weight of the cassava root (density) was linked to the colour of the intermediate fufu product (fermented *fufu* mash)]. According to the respondents, light weight roots float in the fermenting media during retting and develop dark brown colour, which subsequently affects the colour of the intermediate and final *fufu* product (Supplementary Table 1).

Appearance/smoothness of root skin is another preferred root quality trait driving selection of fresh cassava roots for *fufu* processing. In describing appearance of the root, the respondents stated that roots with dark-coloured peel and regular shape are preferred. According to the respondents, dark coloured peel is an indication that the cassava root is matured. Maturity of the cassava root could be linked to yield of *fufu*. The study by Baafi and Safo-Kantanka (2007) reported an increase in starch yield of cassava varieties of age 12-13 months. Smoothness of the root skin on the other hand is characterized by absence of wrinkles and roughness. According to the respondents, smoothness of the root skin attracts buyers and encourages ease of peeling. Egbeocha *et al.* (2016) and Jimoh *et al.* (2016) in their study reported that irregularity in the appearance of tubers is one of the factors

responsible for the absence of efficient cassava peelers in Nigeria. This study therefore reveals a link between appearance and smoothness of cassava root to yield of end product and ease of peeling.

White colour of the inner cassava flesh is another trait that was highly rated as seen in the result (Fig 2 and 3). According to the respondents, over 90% of consumers prefer white coloured *fufu*. Ayetigbo *et al.* (2018) reported that colour of cassava flesh is retained in the derived products. Hence Bechoff *et al.* (2018) reported that yellow cassava flesh arising from the presence of carotenoids in the root results in yellow coloured intermediate and cooked *fufu*. By implication, the use of non-white coloured cassava flesh for *fufu* processing will result in intermediate and cooked *fufu* with non-white colour. This will invariably reduce the acceptance of such *fufu* within the study area. However Sanni *et al.* (1998) reported that *fufu* of good quality will either have a creamy-white, grey or yellow colour. Tomlins *et al.* (2007) further reported that *fufu* flour should be creamier in appearance to increase their acceptability. Flesh color has therefore become vital in the selection of cassava for food (Vimala *et al.* 2010).

# Preferred characteristics of cassava root for processing of fermented wet fufu mash

The results in Fig. 4 show preferred quality characteristics of cassava root for processing of fermented wet *fufu* mash. The results revealed gender dissimilarities in scoring of the preferred processing traits. The most outstanding preferred traits for the females were: 'easy to peel' (85), 'freshness of roots' (28) (indicating absence of root rot and wound) and 'root foaming or retting ability' (22). The traits of interest for the males were 'white colour' (36) and 'freshness of roots' (12). Generally, results from the survey showed that the female respondents attribute more to issues related to processing ability of cassava roots for *fufu* processing compared to their male counterparts. This findings indicate that women are the major processors of fermented *fufu* mash within the study area (Teeken *et al.*, 2018). The traits which were highly preferred by the females are directly linked to certain *fufu* processing operations such as peeling and fermentation of peeled roots. Indigenous knowledge and experience acquired by female processors over years on the effect of these traits on the wet mash, and end product (cooked *fufu*) guide selection of these traits.

According to the female respondents, use of cassava varieties that are 'easy to peel' maximizes the efficiency of labour and time needed to carry out other unit operations such as washing and grating. Barati *et al.* (2020) reported that increase in peeling efficiency is associated with increased peeled surface area and reduced peel loss. This enables the white colour of the flesh to dominate over the dark colour of the peel. Previous studies indicate that cassava peels contain certain phytochemicals such as phenols and tannins responsible for dis-colouation of intermediate and final cassava products such as cassava flour and *fufu* (Hongbete *et al.* 2009, Bindzi *et al.* 2014). Mokemiabeka *et al.* (2011), also reported that *fufu* from well peeled cassava had brighter white colour compared to *fufu* from unpeeled roots. This was attributed to lower tannin content in the peeled roots. Colour, according to Awoyale *et al.* (2018), is a major trait that drives visual appeal and acceptance of cassava products by consumers. This suggests that selection of 'easy to peel' cassava varieties by end users may be due to increased efficiency during manual peeling of such varieties, which invariably reduces tannin and phenol content, resulting in the production of intermediate and final *fufu* with desirable colour attribute.

Freshness of the root indicated by absence of rots/wounds on the roots is another important cassava root quality trait (Fig. 4). This was associated with the presence of 'milkish' white sap at the proximal end of the root. This processors' preferred trait was also linked to the preferred traits (ease of peeling and high retting ability) by the respondents. The preference for freshly harvested roots compared to stored ones may be linked to the onset of Post-harvest physiological deterioration (PPD) of roots shortly after harvest (Zainuddin and Vanderschuren, 2018). PPD is associated with certain undesirable features such as vascular streaking, dis-colouration of roots, reduction in starch quality and shelf life, increased water loss and sugar content (Opara, 1999; Opara, 2009; Buschmann *et al.* 2000; Sánchez *et al.* 2006; Zainuddin and Vanderschuren, 2018). Furthermore, Swain (1979) and Rickard (1986) revealed that adverse effect of storing cassava roots could be linked to the synthesis of anti-nutritional compounds such as polyphenols and tannins which eventually results in dis-colouration of the intermediate and cooked *fufu.* Additionally, Ampe *et al.* (1994) reported that storage of roots prior to retting slightly increased the production of lactate and ethanol during retting, resulting in decreased acceptability of *fufu.* From these previous studies, it could be deduced that the preferred trait 'fresh

root' correlates positively with yield of intermediate and final product and quality in terms of colour and acceptability of *fufu*. Hence, Ampe *et al.* (1994) recommended that retting should be performed with freshly harvested and peeled roots. The study by Omosuli *et al.* (2017) also stated that storage of cassava roots leads to increase in peel loss, decrease in yield of *fufu* flour and cyanogenic potential. According to the respondents, freshness of root is also associated with appreciable moisture content. which is an important factor that facilitates peeling and encourages growth of micro-organisms essential for rapid retting of the roots. This implies that the preferred trait 'freshness of root' is intertwined with other preferred processing traits 'ease of peeling' and 'high retting ability' hence, the importance of this root quality trait to processors and other *fufu* end users.

Foaming or retting ability (easy to ferment) of soaked roots is another preferred processing trait of cassava root mentioned by the female respondents (22) (Fig 4). It is characterized by presence of multiple bubbles covering the soaked roots in the fermenting vessel, and it is a key step in *fufu* processing (Sanni et al. 1998). Variability in *fufu* quality has been attributed to various local practices during retting stage of processing (Sanni et al. 1998). According to Obilie et al. (2004) and Ampe et al. (1994), retting is used to reduce cyanogenic compounds, and to improve the organoleptic quality of cassava by-products such as *fufu*. Additionally, Umeh and Odibo (2014) reported that complete retting of fresh cassava roots results in high yield of wet *fufu* mash and enhances detoxification. The study by Otoo et al. (2018) revealed that the reduced pH achieved through retting impacts sour taste and characteristic aroma to *fufu*. However, prolonged retting and reduced starch content of the fresh cassava roots have been reported to result in intense *fufu* odour that is usually undesirable to consumers (Bechoof et al. 2018; Achi and Akomas, 2006). Retting has also been linked to textural properties of the cooked *fufu*. Isirima et al. (2018) observed a higher and better index for drawability, mouldability, smoothness and colour in *fufu* processed using the retting method compared to other cassava processing methods. The improved textural property of *fufu* processed using retting method is linked to the breaking up of carbohydrate granules to smaller particles through the disintegration of building molecules in the cassava (Isirima et al. 2018). The author further revealed that incomplete break down or absence of retting resulted in products with coarse granules or particles. Hongbete et al. (2009) and Bindzi et al. (2013) furthermore, reported that leaching out of phenol during soaking

(retting) and dewatering of cassava roots enhances the white colour of cassava products such as flour and cooked *fufu*. This study therefore reveals that the trait 'high retting ability' influences most of the organoleptic properties of *fufu* such as colour, aroma, texture and overall acceptance of the end product, hence its importance to the respondents.

# High and low processing properties of fermented fufu mash for cooking of fufu

With regards to the preferred cooking properties of the intermediate *fufu* product (fermented wet *fufu* mash), Fig. 5 shows that the trait 'easy to form dough' is of key interest (high sum of scores) to both the male (12) and female respondents (36), however 'drawing ability' and 'thickness' of the wet mash during cooking were prioritized by the women (74) and (24) respectively, who also are the key actors at this level of *fufu* processing. According to the respondents in our survey, 'dough formation' of mash is related to health status of the crop indicated by the absence of pest and diseases. This is in agreement with previous study by Numfor (1999), who had also related dough formation ability of *fufu* mash to health of the cassava variety used during processing. This highly prioritized trait 'easy to form dough' may also be related to ease of gelatinization and pasting properties of the fermented *fufu* mash. According to Bechoff et al. (2017), fast gelatinization of the starch paste is the trait that is highly appreciated by *lafun* and *fufu* processors. Gelatinization which is manifested by swelling, disruption of hydrogen bonds, crystallite melting with subsequent disappearance of maltese cross, viscosity development, and starch molecules solubilisation, is accompanied by changes in viscosity and formation of paste. On the other hand, pasting properties of *fufu* flour are important quality indices in predicting the behaviour of *fufu* paste during and after cooking (Etudaiye *et al.* 2018). The study by Bindzi et al. (2014) reported that pasting temperatures are indicative of the minimum energy required to initiate rapid absorption of water and swelling of starch granules resulting in increased viscosity and formation of dough. Pasting temperature gives an indication of the minimum temperature for cooking of a given sample (Bindziet al. 2014). The study by Chisenga et al. (2019) further revealed that pasting property of starch, defined by the pasting temperatures and viscosities is affected by amylose content and proportion of components in the food matrix. This study therefore suggest that the trait 'easy formation of dough' is related to lower pasting temperature and time, and enhanced by the utilization of cassava varieties that are free of pest and diseases. Cassava varieties

with such quality traits will therefore require reduced processing time and labour, hence its importance to the respondents.

The study further reveals that the trait 'thickness' of *fufu* mash during processing is another determinant of preference especially among the female respondents. This may be referred to the gel strength and viscosity of the aqueous fermented starch paste which develops during stirring and cooking of final product. This trait can be characterized using pasting and rheological or textural properties of the thick cassava slury obtained from the fermented *fufu* mash. This female preferred processing trait may be influenced negatively by the other female preferred trait (high retting ability). According to Nkoudou *et al.* (2020), accelerated retting process reduces the thickening power of the derived cassava flour. The author further reported that degradation rate and the softening degree of cassava roots influences viscosity; accelerated retting process results in reduced viscosity. Thickness of *fufu* mash may also facilitate formation of dough of the mash and reduce cooking time.

The preferred processing trait, 'drawing ability' may be likened to the degree of cohesiveness of the *fufu* paste during cooking. According to Bechoff *et al.* (2018), cohesiveness, a major textural property that drives consumer acceptance is linked to starch composition especially amylose content. The earlier findings by Numfor (1999) and Rosales-soto *et al.* (2016), indicated that cohesiveness in cassava paste is associated with intermolecular forces within the food, and failure of starch granules to release sufficient amylose. This leads to reduction in cohesiveness of fermented cassava product. In addition, Dufour *et al.* (2002) reported that presence of fibre can reduce the cohesiveness of cassava paste. These textural and pasting characteristics of starch have been associated with cooking quality and texture of various food products (Otegbayo *et al.* 2006). The preferred trait (drawing ability) may have a bearing on retting ability of the cassava roots. Isirima *et al.* (2018) reported an acceptable degree of stickiness, mouldability and drawability for *fufu* made from flour processed by retting compared to those from blanching and direct method of cassava processing

# Preferred and less preferred quality characteristics of fufu during consumption

Two quality traits, 'fufu smoothness' and 'easy to swallow' were of utmost importance (highest sum of scores) to male and female *fufu* consumers in South-East Nigeria (Fig. 6). Our study however shows that female respondents (57) placed higher emphasis on *fufu* smoothness compared to their male counterparts (39). Similarly, the trait 'easy to swallow' was assigned higher weight by greater number of the females (79) than males (38). Bechoff et al. (2018) described smooth fufu as dough that is homogeneous in appearance and hand-feel, and does not have notable fibres, lumps or particles. The author further reveals that *fufu* smoothness is enhanced by the removal of fibre during the sieving operation of fermented roots or mash. This further corroborates the report by Uyoh et al. (2009) who stated that insoluble fibres are disintegrated by cellulolytic and pectinolytic enzymes activities during retting, hence enhancing efficiency of sieving and smoothness of *fufu*. This finding suggests that smoothness of *fufu* may be correlated to the fibre content of the fresh cassava root. Our study further revealed that smoothness of *fufu* was also associated with the trait 'easy to swallow' by the male respondents, implying that the two terms are related. Dziedzoave et al. (1999) also related smoothness to uniformity of particles. Hence the highly preferred textural attributes of cooked *fufu* 'smoothness' and 'easy to swallow', could be related to biophysical properties of the fresh cassava root such as fibre content and efficiency of the unit operation 'retting' and 'sieving' during *fufu* processing.

In contrast, the respondents associated two outstanding traits (stickiness and offensive odour) to low quality attributes of cooked *fufu* (Fig. 7). Stickiness in cooked *fufu* seems to be a trait of key importance to female consumers (23) compared to the male respondents (15). Stickiness could be likened to the textural attribute 'adhesiveness' which is described as the work required in overcoming the attractive force between a product and the contact surface (Singh *et al.* 2013). Bechoff *et al.* (2018) reported high elastic and sticky texture in *eba* and *fufu* prepared from white flesh cassava varieties compared to yellow flesh varieties. This was attributed to their high dry matter and starch contents. The study also shows that 'intense *fufu* odour' is a less preferred trait mainly for female *fufu* consumers (103). However, fewer male respondents (33) indicated that odour of *fufu* lowers their preference for the product. This is in agreement with the report by Uyoh *et al.* (2009) who stated that one major problem in processed *fufu* is the flavour of the product, which may be unacceptable to many people. Shittu and Adedokun (2010) and Tomlins *et al.* (2007) further indicated that the

acceptability of *fufu* by consumers is related to its characteristic aroma. Increased fermentation time had already been reported to correlate positively with the intensity of *fufu* aroma (Bechoff *et al.*, 2018; Achi and Akomas, 2006). However, Bechoff *et al.* (2018) reported a negative correlation between starch content of different cassava varieties and intensity of typical *fufu* aroma.

This study therefore shows that 'smoothness', 'easy to swallow', 'bad odour', and 'stickiness' are the main traits that positively or negatively influence consumption of cooked *fufu* in South-East Nigeria. The quality attributes; stickiness' and 'easy to swallow' are related to the sensory property and texture, while the traits- 'bad odour' and 'smoothness' are related to aroma and appearance respectively. The study also shows that these sensory attributes of *fufu* may be related to the processing parameters 'retting' and 'sieving'.

# Conclusion

The study analyzed consumer preferences and quality attributes of *fufu* in the South-East region of Nigeria between men and women. Major preferred root quality traits identified in locally preferred varieties [Dabere, Imo best, Torokwem, Gbayuomma, Aguoegbulam, akwatakwa, Akpalam Aka, Codelia, Ogwuru ego, Mmaduabuchi, Sakasaka, Akpu da grace, Nwaocha, Nwaibibi, Nwanviumuahia (lady from Umuahia), Yellow root (Vit. C)], TMEB 419, Agric., and TMS 98/0505, within the study area were; root size, heaviness, appearance/smoothness of outer root skin, and inner root flesh colour. 'Ease of peeling', 'freshness of root' and 'high retting ability' (easy to ferment) were identified as the preferred processing traits for the fermented *fufu* mash, while 'easy to form dough', 'thickness', 'drawing ability', stood out as the traits of preference during cooking of the fermented wet *fufu* mash. At the point of consumption, 'smoothness' and 'easy to swallow' were identified as the major determinants of preference, while 'stickiness' and 'intense *fufu* aroma' were identified as less preferred traits. The study therefore reveals that the preference of these traits by the female respondents was directly linked to the processing parameters, and overall quality of the end product. On the contrary, economic gain was observed to be the key determinant driving the choice of the male respondents. The study further indicated that the preferred root qualities; 'root size, heaviness', 'appearance/smoothness of outer root skin' and 'inner root flesh colour has effect on the

processing trait- 'ease of peeling', and on the colour and yield of the fermented *fufu* mash, and cooked dough. The study further suggest that the preferred processing traits; 'ease of peeling', 'freshness of root' and 'high retting ability' (easy to ferment) are intertwined and are associated with colour, yield, organoleptic properties and overall acceptance of cooked *fufu*. However, the retting ability of cassava roots was observed to be a major factor influencing most of the processing traits and those of the cooked product. Our finding therefore reveals an undeniable link between the preferred root quality traits, preferred processing traits and overall quality of cooked *fufu*. Emphasis on root size, heaviness, appearance and root flesh colour, and the relation of these traits to preferred processing and *fufu* quality attributes provides insight for cassava breeders on areas to redirect breeding activities. These findings therefore, make it imperative for introduction of a multi-disciplinary, multi-stakeholder approach into crop improvement programmes, especially the science behind food processing. This will assist in breeding of varieties that meet end-user needs and enhance adoption of these new varieties. The female-preferred traits should be of utmost importance to the breeder, since they are the key actors in the processing operations. There may be need to conduct scientific investigation using throughput methods such as NIRS to identify the biochemical properties that affect these quality traits.

# **Ethical Guidelines**

This study was assessed and approved by the National Research Ethics Committee. Research teams obtained ethical approval prior to the fieldwork. Participants were informed about the study and explained that their participation was entirely voluntary, that they could stop the interview at any point and that the responses would be anonymous. Written consent (signature) was sought and obtained from respondents participating in this study.

### Acknowledgements

This study was supported by the RTBfoods project https://rtbfoods.cirad.fr through a grant OPP1178942: Breeding RTB products for end-user preferences (RTBfoods), to the French Agricultural Research Centre for International Development (CIRAD), Montpellier, France, by the

Bill & Melinda Gates Foundation (BMGF). Special thanks to Clair Hershey for his comments and editing the article.

## **Conflict of Interest**

There is no conflict of interest among Authors of this paper

### References

- Achi, O.K. & Akomas, N.S. (2006).Comparative Assessment of Fermentation Techniques in the Processing of Fufu, a Traditional Fermented Cassava Product. *Pakistan Journal of Nutrition*, 5, 224-229.
- Adebowale, A.A. (2005). Rheology and Storage Properties of Tapioca. An unpublished Ph.D. Thesis Federal University of Agriculture, Abeokuta, Nigeria. 231pp.
- Ajayi, A.R. & Ajala, A. A. (1999). Rural Farmers' Participation in Agricultural Development Project (ADP) and The Effect on The Adoption of Innovations: a Case Study of Ekiti-Akoko ADP in Ondo State of Nigeria. *Ghana Journal of Science*, 39, 83-90
- Akingbala, J.O., Oguntimein, G.B. & Abass, A.B. (1991).Effect of processing methods on the quality and acceptability of fufu from low cyanide cassava. *Journal of the Science of Food Agriculture*, 57, 151–154.
- Aluko, Y. A. (2015). Patriarchy and property rights among Yoruba women in Nigeria., *Feminist Economics*, 21, 56-81. doiDOI: 10.1080/13545701.2015.1015591.
- Ampe, F., Brauman A., Trèche, S. & Agossou, A. (1994). Cassava retting: optimization of a traditional fermentation by an experimental research methodology. *J Sci Food Agric.*, 65, 355-361.
- Anderson, J., Marita, C., Musiime, D. & Thiam, (2017).National Survey and Segmentation of Small Holder Households in Nigeria.Understanding their demand for financial, agricultural and digital solutions.CGAP Working Paper. 93pp. Available at file:///C:/Users/USER/Downloads/Working%20Paper\_CGAP%20Smallholder%20Household% 20Survey\_NGA\_Oct%202017.pdf

- Awoyale, W., Asiedu, R., Kawalawu, W.K.C., Maziya-Dixon, B., Abass, A., Edet, M. & Adetunji,
  M.O. (2018). Assessment of the heavy metals and microbial contamination of garri from
  Liberia. *Food Science and Nutrition*, 6, 62-66.
- Baafi, E. & Safo-Kantanka, O. (2007). Effect of genotype, age and location on cassava starch yield and quality. *Journal of Agronomy*, 6(4), 581-585.
- Bamidele, O.P., Fasogbon, M. B., Oladiran, D.A. & Akande, E.O., (2015). Nutritional composition of fufu analog flour produced from cassava root (Manihot esculenta) and cocoyam (Colocasia esculenta) tuber. *Food science and Nutrition*, 3(6), 597-603.
- Barati, Z., Latif, S., Romuli, S. & Müller J., (2020). Enzyme-Assisted mechanical peeling of cassava tubers. http://www.mdpi.com/2073-4344/10/1/66/s1.
- Bechoff, A., Chijioke, U., Westby, A. & Tomlins, K.I. (2018) 'Yellow is good for you': Consumer perception and acceptability of fortified and biofortified cassava products. *PLoS ONE*, 13(9),1-22 e0203421. https://doi.org/10.1371/journal.pone.0203421
  - The authors (Bechoff, A., Chijioke, U., Westby, A. & Tomlins, K.I. (2018)) stated that varietal difference in the retting ability of cassava roots had significant effect on characteristic aroma of fufu a major factor driving consumer acceptance cooked fufu, this is in agreement with our findings.
- Bechoff, A., Tomlins, K., Fliedel, G., Luis Lopez-lavalle, A.B., Westby, A., Hershey, C. & Dufour,
  D. (2017). Cassava traits and end-user preference: Relating traits to consumer liking, sensory
  perception, and genetics. *Critical Reviews in Food Science and Nutrition*, 58, 547-567. DOIdoi: 10.1080/10408398.2016.1202888
- Bentley, J., Olanrewaju, A., Madu, T., Olaosebikan, O., Abdoulaye, T., Wossen, T., Manyong, V., Kulakow, P., Ayedun, B., Ojide, M., Girma, G., Rabbi, I., Asumugha, G. & Tokula, M. (2017). Cassava farmers' preferences for varieties and seed dissemination system in Nigeria: Gender and regional perspectives. International Institute of Tropical Agriculture, Ibadan.IITA Monograph. 108pp.
- Berry, S.A. (1993). Socio-economic aspect of cassava cultivation and use in Africa: Implication for development of appropriate technology. COSCA Working Paper No.8, 1993, IITA, Ibadan. 24pp.

- Bindzi, J.M., Guiama, V.D. & Ndjouenkeu, R.(2014). Combined smoking and drying kinetics of retted cassava pulp and effects on the physicochemical and pasting properties of the extracted starch. *Food and bioprocess technology*, 7, 2749-2758.
- Blanshard, A. F.J., Dahniya, M.T., Poulter, N.H. & Taylor, A.J. (1994). Quality of processed cassava foods in Sierra Leon. *Journal of the Science of Food and Agriculture*, 64, 425-432.
- M. C. & S. (2003). evaluation-texture. Bourne. Szczesniak, A. Sensory In: B. Caballero, L. Trugo, P. Finglus (Eds.), Encyclopedia of Food Science and Nutrition, Academic press, Amsterdam, The Netherlands. Pp. 5167-5174.
- Buschmann, H., Reilly, K., Rodriguez, M.X., Tohme, J. & Beeching, J.R. (2000). Hydrogen peroxide and flavan-3-ols in storage roots of cassava (Manihot esculentaCrantz) during postharvest deterioration. *J. Agr. Food Chem.*, 48, 5522-5529.
- Chisenga, S. M., Workneh, T.S., Bultosa, G. and Alim, B.A. (2019). Progress in research and applications of cassava flour and starch: A review. *Journal of Food Science Technology*, 56 (6), 2799–2813. https://doi.org/10.1007/s13197-019-03814-6
- Dufour, D., O' Brein, G.M. & Best, R.(2002). Cassava flour and starch: progress in research and development. International Center for Tropical Agriculture (CIAT), Apartado Aereo 6713, Cali, Colombia.
- Dziedzoave, N. T., Ellis, W. O., Oldham, J. H. & Osei-Yaw, A. (1999). Subjective and objective assessment of agbelima (cassava dough) quality. *Food Control*, 10, 63-67.
- Egbeocha, C.C., Asoegwu, S. N. & Okereke, N.A. (2016). A review on performance of cassava peeling machines in Nigeria. *Futo J. Ser. (FUTOJNLS)*. 2, 140-168.
- Egwim, E., Amanabo, M., Yahaya, A. & Bello, M., (2013). Nigerian Indigenous fermented foods: processes and prospects. Intech open science/open mind. http://x.doi.org/10.5772/52877.
- Enete, A., Nweke, F. & Tollens, E. (2002). Contributions of men and women to food crop production labor in Africa: information from COSCA. *Outlook on Agriculture*, 3, 259-265.
- Etudaiye, H.A., Nwabueze, T.U. & Sanni, L.O. (2008). Quality of fufu processed from cassava mosaic disease (CMD) resistant varieties. *African Journal of Food Science*, 3 (3), 061-067.
- FAO (2016). Food and Agriculture Organization. Seed Security Assessment: A Practitioner's Guide.Food texture and viscosity. Sandiego Academic Press.

- Flibert, G., Abel, T. & Aly, S. (2016). African cassava traditional fermented food: the microorganism's contribution to their nutritional and safety values- A review. *International Journal of Current Microbiology and Applied Sciences*, 5(10), 664-687.
- Forsythe, L., Posthumus, H. & Adrienne, M. (2016). A crop of one's own? Women's experiences of cassava commercialization in Nigeria and Malawi. *Journal of Gender, Agriculture and Food Security*, 1, 110-128.
- Forsythe, L., Tufan, H., Bouniol, A., Kleih, U. & Fliedel, G. (2020). An interdisciplinary and participatory methodology to improve user acceptability of root, tuber and banana varieties. *International Journal of Food Science and Technology*, https://ifst.onlinelibrary.wiley.com/doi/10.1111/ijfs.14680.
- González, C. & Johnson, N. (2009).Consumer preferences for table cassava characteristics in Pernambuco, *Brazil.Revista de Economia E Agronegócio*, 7(3), 363-384
- Herrero, M., Thornton, P.K. & Bernués, A. (2014). Exploring future changes in smallholder farming systems by linking socio-economic scenarios with regional and household models, *Global Environmental Change*, 24(1), 165–182,
- Hongbete, F., Mestres, C., Akissoe N. & Nago, M.C., (2009). Effect of processing conditions on cyanide content and colour of cassava flours from West Africa. *African Jjournal of Food Science*, 3(1), 001-006.
- IITA (2012). IITA Annual report on cassava production. http://newint.iita.org/wpcontent/uploads/2026/04/Annual-Report-2012.pdf.
- Isirima, C. B., Iyagba, A. G. & Oliseyenum, O.S. (2018). Organoleptic characteristics of cassava flour and fufu obtained from different cassava varieties (*Manihot spp.*) and tuber processing techniques. *Direct Research Journal of Agriculture and Food Science*, 6(6), 121-127
- Jimoh, M.O., Olukunle, O.J. & Manuwa, S.I. (2016). Modeling of cassava peeling performance using dimensional analysis. *Agric. Eng. Int. CIGR J.* 18, 360-367
- Johnson P.N.T., Gallat, S., Oduro-Yeboah, C., Osei-Yaw, A. & Westby, A. (2006). Sensory properties on instant fufu flour from four high-yielding Ghanaian varieties of cassava. *Tropical Science*, 46, 134-138.

- Karam,L. B., Grossmann,M. V. E., Silva, S. F. R., Ferrero, C. & Zaritzky, N. E. (2005). Gel Textural Characteristics of Corn, Cassava and Yam Starch Blends: A Mixture Surface Response Methodology Approach. *Starch/Stärke*, 57, 62–70
- Mokemiabeka, S., Dhellot, J., Kobawila, S. C. & Louembe, D. (2011). Traditional retting of cassava roots in the ponds or the rivers. *International Journal of Biological and Chemical. Sciences*, 5, 134-141.

The study by Mokemiabeka, S., Dhellot, J., Kobawila, S. C. & Louembe, D. (2011) showed variability in characteristics of the products obtained from retted peeled and unpeeled cassava roots. Fufu from the peeled roots contained less tannin which enhanced whiteness of fufu. The quality trait 'colour' and 'ease of peeling' were highly priotized by the respondents. This buttressed the fact that there is arelationship between ease of peeling and colour of the final product.

- Nkoudou N. Z., Ndjouenkeu, R. and Ngang, J.J.E. (2020). Effect of accelerated retting process on physiochemical and pasting properties of cassava (*Manihot esculenta Crantz*) flours. *Asian Food Science Journal*, 15(4), 45-52
- Numfor, F.A. (1999). Physicochemical changes in cassava starch and flour associated with fermentation, effect in textural properties. *Journal of Food Technology in Africa*, 4, 31-35.
- Nweke, F., Haggblade, S. & Zulu, B. (1994). Building on successes in African agriculture recent growth in African cassava; farm level practice relevant to cassava plant protection. *African Crop Science Journal*, 2, 563-582.
- Nweke, F.I. & Bokanga, M. (1994). Importance of cassava processing for production in sub-Saharan Africa. *Acta Horticulture*, 375, 401-412.
- Obadina, A., Olusola, O.,Sanni, L., Keith, T & Westby, A. (2008). Identification of hazards and critical control points (CCP) for cassava fufu processing in south-west Nigeria. *Food Control*, 19, 22-26. 10.1016/j.foodcont.2007.01.002.
- Obilie, E.M., Tano-Debrah, K. & Amoa-Awua, W.K. (2004). Souring and breakdown of cyanogenic glucosides during the processing of cassava into akyeke. *Int. J. Food Microbiol.*, 93, 115-121.
- Omodamiro, R. M., Oti, E., Etudaiye, H. A., Egesi, C., Olasanmi, B. & Ukpabi, U. J. (2012). Production of fufu from yellow cassava roots using the odourless flour technique and the

traditional method: Evaluation of carotenoids retention in the fufu. *Advances in Applied. Science Research*, 3, 2566-2572.

Omosuli, S.V., Ikujenlola, V. & Abisuwa, A.T. (2017). Quality assessment of stored fresh cassava roots and fufu flour produced from stored roots. *Journal of Food Science and Nutrition Therapy*, 3(1), 009-013.

The study by Mokemiabeka, S., Dhellot, J., Kobawila, S. C. & Louembe, D. (2011) showed variability in characteristics of the products obtained from retted peeled and unpeeled cassava roots. Fufu from the peeled roots contained less tannin which enhanced whiteness of fufu. The quality trait 'colour' and 'ease of peeling' were highly priotized by the respondents. This buttressed the fact that there is arelationship between ease of peeling and colour of the final product.

- Opara, U. L. (1999). Cassava storage.CIGR Handbook of Agricultural Engineering Agro.St.Joseph, MI, American Society of Agricultural Engineers.IV
- Opara, U. L. (2009). Postharvest technology of root and tuber crops. In: R. Dris, R. Niskanen & S. M. Jain (Eds.), *Crop management and post-harvest handling of horticultural product*. Science publishers Inc. Pp. 382-406.
- Otegbayo, B., Aina, J., Asiedu, .R. & Bokanga, .M. (2006).Pasting characteristics of fresh yams (Dioscorea spp.) as indicators of textural quality in a major food product- pounded yam. *Food Chemistry*, 99, 663–669.
- Otoo, G.S., Essumanc, E.K., Gyimahd, V. & Bigson, K. (2018). Quality attributes of fufu: Instrumental and sensory measurement. *Scientific African*, 1, 1-7.
- Owolarafe, O.K., Adetifa, B.O., Oyekanmi, A.C., Lemikan, T.G. & Samuel, T.M (2018). Assessment of fufu production technologies in Ogun State, Nigeria. *Arid Zone Journal of Engineering, Technology and Environment*, 14(4), 547-558.
- Oyewole, O. B. & Odunfa, S.A. (1990). Characterization and distribution of lactic acid bacteria in cassava fermentation during fufu production. *Journal of Applied Bacteriology*, 68, 45-52.
- Oyewole, O. B. & Ogundele, S. L. (2004). Effect of length of fermentation on the functional characteristics of fermented cassava 'fufu'. *Journal of Food Technology in Africa*, 6, 38-40.

- Oyewole, O. B. & Sanni. L.O. (1995). Constraints in traditional cassava processing: the case of 'fufu' production.In: Cassava Food Processing, Agbor Egbe, T., Brauman, A., Griffon, T & Treche, S. (eds.). ORSTOM, Paris. Pp. 523–529.
- Rickard, J.E. (1986). Tannins levels in cassava: comparison of methods of analysis. *J Sci Food Agric.*, 37, 37-42.
- Rosales-soto, M.U., Ross, C.F., Younce, F., Fellman, J.K., Mattinson S. D., Huber K. & Powers J.R.
  (2016). Physico-chemical and sensory evaluation of cooked fermented protein fortified cassava (manihot esculenta crantz) flour. *Advances in Food Ttechnology and Nnutritional Sciences Oopen Journal*, 2(1), 9-18.
- Sánchez, T., Chavez, A.L., Ceballos, H., Rodriguez-Amaya, D.B., Nestel, P. and Ishitani, M. (2006).
  Reduction or delay of post-harvest physiological deterioration in cassava roots with higher carotenoid content. *Journal of the Science of Food and Agriculture*.86, 634-639.
- Sanni, L. O., Kosoko, S. B., Adebowale, A. A. & Adeoye, R. J. (2004). The influence of palm oil and chemical modification on the pasting and sensory properties of fufu flour. *International Journal* of Food Properties, 7, 229-237.
- Sanni, L.O., Kolawole, A.G., Akingbala, J.O. & Kuye, A. (2007). Effect of drying methods on moisture sorption isotherms of fufu at three temperatures. *Drying Technology*, 17, 286-295. DOI: 10.1080/07373939908917530
- Sanni,L.O., Akingbala, J.O., Oguntunde, A.O., Bainbridge, Z.A., Graffham, A.J. & Westby, A. (1998). Processing of fufu from cassava in Nigeria: Problems and prospects for Development. *Science Technology and Development*, 16, 58–71.
- Shields, S.A. (2008). Gender: An Intersectionality perspective. Sex Roles, 59, 301-311.
- Shittu, T.A. & Adedokun, I.I. (2010). Comparative evaluation of the functional and sensory characteristics of three traditional fermented cassava products. *J. Nat. Sci. Engr. Tech.*, 9 (2), 106-116.
- Singh, V., Guizani, N., Al-Alawi, A., Claereboudt, M. & Rahman, M. (2013).Instrumental texture profile analysis (TPA) off date fruits as a function of its physico-chemical properties. *Industrial Crops and Products*, 50, 866-873.

Swain, T. (1979). Tannins and lignins. In: Herbivores: their interaction with secondary plant metabolites (Eds.) Rosenthal, G.A. & Janzen, D. Academic Press, New York. Pp. 657-682.

Teeken, B., Olaosebikan, O., Haleegoah, J. Oladejo, E., Madu, T. Bello, A., Parkes, E., Egesi, C., Kulakow, P., Kirscht, H. & Tufan, H.A. (2018). Cassava trait preferences of men and women farmers in Nigeria: implications for breeding. *Economic Botany*, 72, 263–277. https://doi.org/10.1007/s12231-018-9421-7.

This study strengthened the fact that cassava trait preferences are gender specific. The researchers investigated preferences around cassava traits and varieties, while examining gender based constraints in accessing and adopting cassava planting material

- Thorpe, H.G. & Muriuki, W. (2001). Smallholder dairy production and marketing in Eastern and Southern Africa: regional synthesis. In *Smallholder Dairy Production and Marketing— Opportunities and Constraints*, W. Rangnekar and D. Thorpe, Eds. Pp. 185–198.
- Tomlins, K., Sanni, L., Oyewole, O., Dipeolu, A., Ayinde, I., Adebayo, K. & Westby, A. (2007).
  Consumer acceptability and sensory evaluation of a fermented cassava product (Nigerian fufu).
  *Journal of the Science of Food and Agriculture*, 87, 1949-1956.
- Trinh, K.T. & Glasgow, S. (2012). On the texture profile analysis test. Conference: Chemical 2012:Wellington, New Zealand. 13pp.
- Uche, C. (2016). Cost and returns structure in garri and fufu processing in Rivers State, Nigeria. *Nigerian Agricultural Policy Research Journal*. 1(1), 131-138. http://aprnetworkng.org.
- Umeh, S.O. & Odibo, F.J.C. (2014). Use of different additives in retting cassava tubers for fufu production. *International Journal of Pharmaceutical Science Invention*, 3 (4), 11-17.
- Uyoh, E.A., Ntui, V.O. & Udoma, N.N. (2009). Effect of local cassava fermentation methods on some physicochemical and sensory properties of fufu. *Pakistan Journal of Nutrition*, 8(8), 1123-1125.
- Vimala, B., Thushara, R. & Nambisan, B. (2010). Carotenoid Retention in Yellow-fleshed Cassava during Processing. In Proceedings of the 15th Triennial International Society for Tropical Root Crops (ISTRC) Symposium, Lima, Peru. Pp. 2–6.
- Zainuddin, I. M., Fathoni, A., Sudarmonowati, E., Beeching, J.R., Gruissem, W. & Vanderschuren, H. (2018). Cassava post-harvest physiological deterioration: from triggers to symptoms. *Postharvest Biology and Technology*. 142, 115-123.

Zhao, Y. & Saldan<sup>°</sup>a, M.D. (2019). Hydrolysis of cassava starch, chitosan and their mixtures in pressurized hot water media. *J Supercrit Fluids*, 147(3), 293–301



Figure 1: Survey locations in Imo and Abia States of Nigeria

ijfs\_14875\_f2.doc



Figure 2: Preferred characteristics of good cassava roots for fufu when buying in the market

ijfs\_14875\_f3.doc



Figure 3: Characteristics of cassava roots in the market expected to give a less preferred fufu product

ijfs\_14875\_f4.doc



ijfs\_14875\_f5.doc





ijfs\_14875\_f6.doc



Figure 6: Preferred characteristics of fufu during consumption

ijfs\_14875\_f7.doc



