

Québec, 17 -20 July 2022

Poster N°226-fM78-1411

Do not forget that agroforestry can also provide wood, be it fuel wood or timber, for the benefit of populations; examples in West and Central Africa.

MALLET Bernard¹, BALLE Pity², NJOUKAM Raphael³, PELTIER Régis⁴, WENCELIUS François¹

Background and aims



Environmental services of agroforestry, such as carbon sequestration and biodiversity maintenance, have been highly emphasised during last decade,.

Wood consumption and production, be it fuelwood or timber, is still critical for a large part of rural and urban populations in many African countries. Agroforestry systems plays a large role in wood production. Improved agroforestry systems could contribute to sustainable wood production, and decrease pressure on natural forests, complementary to forest plantations.







Case studies concern more specifically lvory Coast, Cameroon and Democratic Republic of Congo, in relation to authors' experience. Bibliographic analysis was completed for some important agroforestry systems such as cocoa production and forest fallows in humid regions, and agroforestry parklands and hedges in more more wet and dry savannahs.



Results

- Integration of fast-growing trees, mainly local or introduced nitrogen-fixing trees (such as Acacia mangium and A. auriculiformis), in fallow systems has many advantages: improvement of nitrogen and soil fertility, and then food supply (cassava, maize, ...); carbon sequestration, ... Projects carried out around Abidjan, Bafoussam and Kinshasa show that they can also make a significant contribution to wood and charcoal supply to rural and urban populations, and generate incomes for small agroforestry farmers with limited investments.
- Cocoa production was first developed under natural forest shade. Current strategies





intend to restore old cocoa plantations in line with "zero deforestation" strategies for sustainable cocoa production. Integration of service or timber trees in young cocoa plantations contributes to restoring an adapted microclimate, but also to significant production of wood through appropriate agroforestry management.

Integration of woody hedges and more sustainable management of agroforestry parklands, are agroforestry practices contributing to the restoration of soils and the environment in wet and dry savannahs. These AFS also provide a large quantity of wood for various uses for the populations of the villages, complementary to natural ecosystems.



Conclusion

Discussion

Complementary context based analysis are needed for joint social, economic and environmental integration of the multifunctional role of wood production in AFS, including carbon sequestration.



It is critical that policy makers and field operators do not forget the wood production component in agroforestry strategies. This is important for rural and urban populations, complementary to other services and food production provided by agroforestry systems.

References

- > Peltier, Balle, Mallet et al. 1996. Produire du bois énergie dans les jachères de zone guinéenne. Intérêts et limites à travers l'expérience d'Oumé en Basse Côte d'Ivoire
- > Temgoua, Njoukam, Peltier. 2021. Ingéniosité, diversité, vision et adaptabilité des agriculteurs produisant du bois et de PFNL dans les montagnes de l'Ouest-Cameroun.
- > Dubiez et al. 2019. Long term impact of Acacia auriculiformis woodlots growing in rotation with cassava and maize on the carbon and nutrient contents of savannah sandy soils in the humid tropics (Democratic Republic of Congo).
- > Smektala et al. 2005. Parcs agroforestiers sahéliens: de la conservation à l'aménagement.



1 : <u>bernard.mallet@cirad.fr</u> Association des forestiers tropicaux et d'Afrique du Nord, France.







