

Tropical Starches from South-East Asia (1): Morphological, Thermal and Pasting properties.

Guillaume Da^{1,3*}, Thierry Tran², Hong Luong Nga³, Juan Bautista Sanz Hernández²,
Dominique Dufour^{4,5}, Sunee Chotineeranat², Le Thanh Mai³, Klanarong Sriroth²

¹ *Laboratoire GPMA-ENSBANA (EA4181), 1 esplanade Erasme, 21000 Dijon, France*

² *Cassava and Starch Technology Research Unit (CSTRU/BIOTEC), Kasetsart University, Jatujak, Bangkok 10900, Thailand*

³ *Hanoi University of Technology (HUT), IBFT, Hanoi, Vietnam*

⁴ *Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), 73 rue Jean-Francois Breton, TA B-95/16, 34398 Montpellier Cedex 5, France*

⁵ *Centro Internacional de Agricultura Tropical (CIAT), Apdo Aéreo 6713, Cali, Colombia*

*Corresponding author: guillaumeda@gmail.com Tel/Fax: +33380396699/+333806898

Keywords: Starches, Tropical tubers, Physicochemical properties, Functional Properties.

Abstract:

This study attempts at exploring the physicochemical and functional characteristics of several types of tropical starches separated from starch-producing plants from South-East Asia. In Vietnam and Thailand, despite the fact that starch is the major component of tubers, these crops remain underutilized and represent a potential field for post-harvest applications. In this work, selected root and tuber crops were potato, sweet potato, cassava, taro, yam bean and canna. Sago was the tree crop sample; kudzu and mung bean as legumes samples. Common cereals like rice and maize were also selected as references. Granule size was measured by laser diffraction. Thermal properties were obtained by DSC analysis. Viscosity properties were determined by RVA. The results obtained with cereals, legumes and other types of starches, were contrasted, suggesting a separate analysis of starches from root and tubers. The selected root and tuber starches had lipid, protein and ash contents in the range of 0.16-0.52%, 0.08-0.21% and 0.01-0.44% respectively. Significant differences in the granule size distribution between starches were observed with the largest type of granules for canna (55µm). DSC gelatinization temperatures were lower for cassava than for aroids. Further investigations are needed to better understand the relationships between proximate analysis, granule distribution and thermal properties of starches, thus the quality of the foodstuffs manufactured from root and tuber crops.