



**THE AFRICAN ASSOCIATION OF INSECT SCIENTISTS**

P. O. Box 59862, 00200 City Square  
NAIROBI, KENYA



## 18ème Conférence de l'Association Africaine des Entomologistes

## 18th Conference of the African Association of Insect Scientists

Salle de Conférence du Ministère de l'Agriculture Ouaga 2000/  
Conference room of the Ministry of Agriculture Ouaga 2000  
OUAGADOUGOU, BURKINA FASO

16 - 20 Novembre / 16 – 20 November 2009

**"Gestion des insectes ravageurs des cultures  
et vecteurs de maladies pour un  
environnement viable et une sécurité  
alimentaire en Afrique: Développements  
courants"**

**"Insect pest and vector management for  
sustainable environment and food security in  
Africa: Current developments"**

# Programme

<sup>1</sup>International Centre of Insect Physiology and Ecology (*icipe*)<sup>1</sup>  
<sup>2</sup>Jomo Kenyatta University of Agriculture and Technology; dmwangi@icipe.org;  
carol.aldous@lshtm.ac.ke; gitonga@fsc.jkuat.ac.ke; mndugu@fsc.jkuat.ac.ke;  
ahassanali@icipe.org; pnjagi@icipe.org; nmaniania@icipe.org  
\* dmwangi@icipe.org , \*ahassanali@icipe.org

Foraging insects encounter diverse array of microorganisms with potential benefits and risks in their habitats. For instance, termites encounter symbionts and pathogenic fungi in their hemiedaphic habitats. An important question is how these insects distinguish between beneficial and risky fungi species. Our preliminary study showed that *Macrotermes michaelseni* detected an entomopathogenic isolate of *Metarhizium anisopliae* through olfaction and avoided direct physical contact. In addition, we found a differential response of the termites to the isolates of the fungus with different pathogenicity. We hypothesized that this avoidance behaviour reflects different composition of the volatiles emanating from the fungal strains. We collected volatile blends of different isolates and analyzed them by GC-MS analysis. Our results show that there are both qualitative and quantitative differences in these volatiles. We are currently characterizing the active repellent blend responsible for the avoidance behaviour of termites towards pathogenic fungi.

**Keywords:** Termites, Fungus, Interactions and entomopatho chemicals

#### **A Comparative study on the life cycles of the leaf miners *Liriomyza trifolii* (Burgesss) and *Liriomyza sativae* (Blanchard) (Agromyzidae: Diptera).**

Salah Faiza E. E.<sup>1</sup>, Elamin Elamin M.<sup>2</sup>, Eltoum Elameen M. A.<sup>1</sup> & Bordat Dominique<sup>3</sup>

<sup>1</sup> Dept. of Crop Protection, Faculty of Agricultural Sciences, University of Gezira,  
Wad Medani, P.O. Box 20 Sudan. Email: Faizaruba@yahoo.fr

<sup>2</sup> Agricultural Research Corporation, Wad Medani, Sudan.

<sup>3</sup> CIRAD, Montpellier, France.

The life cycles of the leaf miners, *Liriomyza trifolii* (Burgess) and *Liriomyza sativae* (Blanchard), were studied in a growth chamber (Temp,  $25 \pm 1^\circ\text{C}$ ; R.H.  $75 \pm 5\%$ ; L/D 12 hL/12 hD and 3000 – 4000 LUX) at CIRAD, Montpellier, France. The host plant used was haricot bean, (*Phaseolus vulgaris*). The egg incubation period of both species lasted for two days. Highly significant differences ( $P<0.01$ ) were recorded between the number of feeding punctures caused by *L. trifolii* and *L. sativae*. Differences between the larval periods and number of emerging adults were not significant. Upon hatching, during 1996, the number of larvae of *L. trifolii* was significantly higher than that of *L. sativae*, but the total life span of *L. sativae* was significantly longer than that of *L. trifolii*.

**Key words:** life cycle, leaf miners, *Liriomyza*, Diptera

#### **Six nouvelles espèces de psylles du genre *Pseudoeriopsylla* (Hemiptera, Homotomidae), ravageurs des figuiers au Cameroun.**

Dzokou V. J.<sup>1\*</sup>, Tamesse J. L.<sup>1</sup> & Burckhardt D.<sup>2</sup>

<sup>1\*</sup>. Université de Yaoundé I, Ecole Normale Supérieure, Laboratoire de Zoologie, B.P. 47, Yaoundé, Cameroun.  
dzovijo@yahoo.fr

<sup>2</sup>. Naturhistorisches Museum, Augustinergasse 2, CH-4001 Basel, Switzerland

Les psylles sont des insectes piqueurs et suceurs de la sève des plantes hôtes. La famille des Homotomidae regroupe plusieurs sous familles et genres reconnaissables par les quelques caractères suivants : proctigère mâle bisegmenté, paire de tubercule prononcée sur le metapostnotum, plaque sous-génitale femelle dépourvue d'accessoires latéro-dorsales. La sous famille des Macrohomotominae regroupe les psylles dont les traits communs de caractères sont observables sur les ailes antérieures des adultes, avec notamment l'existence d'un ptérostigma et l'absence de la brisure costale. Parmi les 6 espèces de *Pseudoeriopsylla* africaines, une seule est connue du Cameroun, *P. laingi*. Les investigations entreprises à l'Ouest Cameroun de 2005 à 2008 ont permis de