


# The Chlordecone crisis in the French West Indies : Its fate in soils and water

M. Voltz (1), P. Cattán (2), C. Saison (3), A. E. Berns (4), F. Colin (5), A. Crabit (2,5), D. Crevoisier (1), J. Fernandez- Bayo (1,3), J. Levillain (2), L. T. Pak (1), A. Samouelian (1), and Y.M. Cabidoche (6)

(1) INRA, UMR LISAH, Montpellier, France (voltz@supagro.inra.fr), (2) CIRAD, UPR Systemes Bananes et Ananas, Guadeloupe, Capesterre-Belle-Eau, France, (3) IRD, UMR LISAH, Montpellier, France, (4) Forschungszentrum Juelich GmbH, IBG 3, Juelich, Germany, (5) Supagro, UMR LISAH, Montpellier, France, (6) INRA, UR Agropedoclimatique de la Zone Caraibe, Environment and Agronomy, Domaine Duclos, Petit-Bourg, Guadeloupe, France


## The Chlordecone crisis and the CHLORDEXCO Project



In the French West Indies, chlordecone (CLD), an organochlorine pesticide, which is highly persistent in the environment, was applied in banana plantations from 1972 to 1993 against the banana weevil *Cosmopolites sordidus*. Pollution surveys conducted in 2001 by the French Department of Health revealed the presence of chlordecone in soils, rivers, springs over large areas in Guadeloupe and Martinique islands. Contamination of drinking water, food crops, aquatic species by CLD has been observed as well as its presence in blood of men, pregnant women and newborns. There is therefore a large social concern about the extent and evolution of CLD pollution in the French West Indies and its impact on human health and ecosystems. From 2008 to 2012 a multidisciplinary project CHLORDEXCO took place to study the CLD fate in water, soils and the contamination characteristics of aquatic species and food crops. Here, we summarize results obtained on the processes controlling the spatial and temporal patterns of soil and water contamination at the scale of the banana cropping area in Guadeloupe and of the Perou catchment.

## Contamination of percolation water


Percolation experiments on undisturbed soil columns taken from contaminated areas showed that CLD concentration in percolated water varied little (5 to 10 µg/L) and were of same magnitude for andosols and nitisols although their CLD stocks, sorption properties and hydrodynamic behaviours differed strongly. Intense macropore flow in the andosols and classical Darcian flow in nitisol is observed. Transport modelling is currently under way for elucidating this observation.



## Conclusions


Contamination of soil and water in the banana cropped areas of the French West Indies is general and permanent. If no soil remediation strategies can be elaborated this severe environmental contamination will last over centuries as estimated by Cabidoche et al. (2009). Since CLD propagates mainly by percolation through the soils and by subsurface flow, buffer strips, usually recommended in many regions to limit the pollution of water by pesticides, will not be useful.

## The soils and their contamination level



The main soils in the contaminated areas are andosols, nitisols and ferralsols. They formed from the weathering of volcanic ashes. They have a high organic carbon content that varies in part according to the altitude. They also have high content of secondary minerals, allophane for andosols and halloysite for nitisols and ferralsols.


	Andosol	Nitisol	Ferralsol	Reference
$\theta$	0.44	0.385	0.495	Colmet Daage 1965
$Bd$	0.55	1.1	1.1	Cabidoche et al. 2009
$K_{oc}$	20000	2000	8000	Cabidoche et al. 2009
$C(75m)$	5	2	3	Dorel et al. 2003 and Cabidoche et al. 2009
$C(150m)$	7	2	3	Dorel et al. 2003 and Cabidoche et al. 2009
$C(300m)$	9	2	3	



An analysis of the spatial distribution of CLD in soil over 1045 field plots (Levillain et al., 2012) in the banana cropping area of Guadeloupe showed that the soil type had a strong impact. Andosols, with a high sorption capacity ( $K_{oc}$  20 000 L/kg), had the highest CLD concentrations and stocks, unlike Nitisols, which had 10-fold lower sorption capacities. A significant «farm effect», due to between-farm variations of application times and amounts, was also noticed. The observed stocks of CLD clearly correspond to the accumulation in soil of successive treatments and thereby confirm the high persistence of CLD in soil also observed in incubation studies in soil microcosms.

## Contamination features of rivers

At the outlet of the 12.6 Km<sup>2</sup> Perou catchment in Guadeloupe, contamination of surface waters was permanent with CLD concentrations one order of magnitude less than in percolation water. As can be seen on the figure CLD concentrations decreased when discharge increased. This is due to dilution effects caused by storm water stemming mainly from uncontaminated uphill areas. Thus, CLD concentrations were highest in base flow indicating that the main pathway of applied CLD to surface waters is by percolation and groundwater flow.



## References

Cabidoche, Y.-M., Achard, R., Cattán, P., Clermont-Dauphin, C., Massat, F., Sansoulet, J., 2009. Long term pollution by chlordecone of tropical volcanic soils in the French West Indies: a simple leaching model accounts for current residue contents. *Environmental Pollution*, 137, 1697-1705.  
 Levillain, J., Cattán, P., Colin, F., Voltz, M., Cabidoche, Y.-M. How physical environmental factors and farming practices determine soil contamination by a persistent organic pollutant, chlordecone, in the French West Indies. *Agriculture, Ecosystems and Environment*, 159, 123- 132.  
 Lesueur-Jannoyer, M., Cattán, P., Monti, D., Saison, C., Voltz, M., Woignier, T., Cabidoche, 2012. Chlordecone aux Antilles : évolution des systèmes de culture et leur incidence sur la dispersion de la pollution. *Agronomie, Environnement & Sociétés*, 2(1), 45- 48.

## Acknowledgements

This study was supported by the Chlordecone Action Plan, and by the French National Research Agency under reference ANR-08-CES-004-01, CHLORDEXCO project.