

TOXOPLASMOSIS AND TRICHINELLOSIS: AN EPIDEMIOLOGICAL SURVEY OF PIG POPULATION IN MADAGASCAR

ENQUÊTE ÉPIDÉMIOLOGIQUE SUR LA TOXOPLAMOSE ET LA TRICHINELLOSE EN POPULATION PORCINE À MADAGASCAR

ENCUESTA EPIDEMIOLÓGICA SOBRE TOXOPLASMÓISIS Y TRIQUINÓISIS EN CERDOS EN MADAGASCAR

M. Rakotoharinome¹ H. Andriamanivo² R. Blaga^{3,4} C. Perret⁴ S.A. Lacour⁴
A. Grasset-Chevillot⁴ P. Mace⁴ M. Thomas⁴ I. Villena⁵ D. Aubert⁵ P. Boireau⁴ V. Porphyre^{6,*}

Keywords: Swine – Parasite – Zoonosis – Immunodiagnosis – *Trichinella* – *Toxoplasma* – Madagascar.

Mots-clés : Porcin – Parasite – Zoonose – Immunodiagnostic – *Trichinella* – *Toxoplasma* – Madagascar.

Palabras clave: Cerdo – Parásito – Zoonosis – Inmunodiagnóstico – *Trichinella* – *Toxoplasma* – Madagascar.

Besides cysticercosis, scarce information is available on the other meat-borne zoonotic parasitoses in the domestic animal population of Madagascar. *Trichinella* is an intracellular parasitic nematode of mammalian skeletal muscle, causing a serious zoonotic disease. *Trichinella* is transmitted to humans by consumption of raw or undercooked meat from pig, wild boar and other sensitive species (1). *Toxoplasma gondii* is a protozoan parasite that can probably infect all warm-blooded animals (mammals and birds) and humans. Humans become infected postnatal by ingesting tissue cysts from undercooked meat (various species), consuming food or drink contaminated with oocysts, or by accidentally ingesting oocysts from the environment (2). The present study aimed at investigating the extent of two major parasitic diseases, namely toxoplasmosis and trichinellosis, within the Malagasy pig population.

Two hundred and fifty pig serum samples were collected during 2010 in the four major slaughterhouses of Antananarivo, the Malagasy capital. Sampled pigs were raised in 11 different regions (of a total of 22 regions) and transported by traders before

slaughtering. Samples were stored at -80°C and sent for analysis to the Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail (ANSES, Maisons-Alfort, France). Serological investigations were conducted for both pathogens using ELISA technique with commercial test kits: ID Screen® Toxoplasmosis Indirect ELISA kit (IdVet, France) and PrioCHECK® *Trichinella* Ab (Prionics, Switzerland), according to the manufacturer's instructions.

Results on seroprevalence of *T. gondii* in pigs confirm that the zoonotic parasite is present in Madagascar. Preliminary results




Figure 1: *Toxoplasma* prevalence in pigs in Madagascar regions in 2010.

1. Services vétérinaires de Madagascar, ministère de l'Élevage, rue Farafaty Ampandrianomby, Antananarivo, Madagascar.

2. Fofifa-DRZV, ministère de l'Enseignement supérieur et de la Recherche, Ampandrianomby, 101 Antananarivo, Madagascar.

3. Ecole nationale vétérinaire, Université Paris-Est, 7 avenue du Général de Gaulle, 94704 Maisons-Alfort, France.

4. Anses, Laboratoire de santé animale, 14 Rue Pierre et Marie Curie, 94701 Maisons-Alfort, France.

5. USC ANSES Epi-Toxo, Centre national de référence de la toxoplasmosse, CHU Reims, pôle épidémiologie, 45 rue Cognacq-Jay, 51092 Reims Cedex, France.

6. Cirad, UMR Selmet, station de Ligne-Paradis, 7 chemin de l'IRAT, 97410 Saint-Pierre, Réunion, France

* Corresponding author

E-mail: vincent.porphyre@cirad.fr

show a seroprevalence of 22.8% (57/250) for toxoplasmosis. All regions are endemic for toxoplasmosis with a frequency of positive pigs ranging from 12% (n = 8) in the southern province of Toliara, to 21% in Antananarivo (n = 69) or Fianarantsoa (n = 79) in central uplands, and up to 33% (n = 69) in Mahajanga in the coastal northwestern region; no region of the island can be considered free of toxoplasmosis. No correlations were found with sex, breed or age of pigs. From a public health perspective these results emphasize the need to inform the public on the importance of proper meat handling and preparation.

Regarding *Trichinella*, two pigs presented positive values by ELISA based on excretory/secretory antigens but those results were not confirmed in western blot. Even if the serological test for the detection of *Trichinella* provides a high degree of sensitivity and specificity, false negative results occur during the early stages of infection, especially for light to moderate infection. Further investigations correlating serology and muscle analysis need to be conducted to conclude on *Trichinella* status in Madagascar.

National field investigations in pig and human populations will have to be carried out to confirm the present results as well as to correlate the observed prevalence with on-farm risk factors, and with the transmission of trichinellosis and toxoplasmosis to Malagasy consumers, especially to pregnant women.

REFERENCES

1. RICHOMME C., LACOUR S.A., DUCROT C., GILOT-FROMONT E., CASABIANCA F., MAESTRINI O., VALEE I., GRASSET A., VAN DER GIESSEN J., BOIREAU P., 2010. Epidemiological survey of trichinellosis in wild boar (*Sus scrofa*) and fox (*Vulpes vulpes*) in a French insular region, Corsica. *Vet. Parasit.*, **172**: 150-154. DOI: 10.1016/j.vetpar.2010.04.026
2. VILLENA I., DURAND B., AUBERT D., BLAGA R., GEERS R., THOMAS M., PERRET C., ALLIOT A., ESCOTTE-BINET S., THEBAULT A., BOIREAU P., HALOS L., 2012. New strategy for the survey of *Toxoplasma gondii* in meat for human consumption. *Vet. Parasitol.*, **183**: 203-208. DOI: 10.1016/j.vetpar.2011.08.001

Accepted 30 April 2015; Online publication June 2015