

Cassava roots

Description Nutritional aspects Nutritional tables References

Click on the "Nutritional aspects" tab for recommendations for ruminants, pigs, poultry, rabbits, horses, fish and crustaceans



Common names

Cassava, Brazilian arrowroot, tapioca [English]; manioc, tapioca [French]; yuca, mandioca, tapioca, guacamota, casabe, casava [Spanish]; maniok [German]; cassave, maniok [Dutch]; rogo [Hausa]; ketela pohon, ubi kayu, atau singkong [Indonesian]; mandioca [Portuguese]; kamoteng-kahoy, kasaba [Tagalog]; manyok [Turkish]; sắn, khoai mì [Vietnamese]; Ègè [Yoruba]; لكسانا [Arabic]; (Kāsābhā) [Bengali]; 木薯 [Chinese]; מניהוט מצוי [Hebrew]; [Hindi]; キャッサバ [Japanese]; 카사바, 마니옥 [Korean]; [Malayalam]; Маниок съедобный, кассава [Russian]; [Tamil]; มันสำปะหลัง [Thai]

- **Product names:** Cassava roots, cassava meal, cassava root meal, cassava chips, cassava pellets, cassava hard pellets

Species

Manihot esculenta Crantz [Euphorbiaceae]

Synonyms

Jatropha dulcis J. F. Gmel., *Jatropha manihot* L., *Manihot aipi* Pohl, *Manihot dulcis* (J. F. Gmel.) Pax, *Manihot flabellifolia* Pohl, *Manihot leptopoda* (Müll. Arg.) D. J. Rogers & Appan, *Manihot manihot* (L.) Cockerell, nom. inval., *Manihot melanobasis* Müll. Arg., *Manihot palmata* Müll. Arg., *Manihot palmata* var. *leptopoda* Müll. Arg., *Manihot peruviana* Müll. Arg., *Manihot saxicola* Lanj., *Manihot tristis* Müll. Arg., *Manihot tristis* subsp. *saxicola* (Lanj.) D. J. Rogers & Appan, *Manihot utilissima* Pohl (USDA, 2009)

Feed categories

- Roots, tubers and by-products
- Plant products and by-products

Related feed(s)

- Cassava leaves and foliage
- Cassava peels, cassava pomace and other cassava by-products

Description

Cassava (*Manihot esculenta* Crantz) is a shrub grown in the tropics and subtropics for its underground starchy tuberous roots. Cassava roots, also called cassava tubers, are a major staple food for more than 800 million people in the world (Eccocrop, 2011; Lebot, 2009).

Morphology

The cassava plant is a woody shrub, reaching 2 to 4 m in height. The cassava tuber consists of the bark (outermost layer, 0.5-2% of the organ; easily removed by simple scratching), the peel (1-2 mm thick; 8-15% of the tuber; it contains most of the toxic cyanogenic glucosides) and the fleshy starchy parenchyma (83-92% of the tuber) which is the edible part of agricultural importance (Lebot, 2009; Tewe, 1992). Each plant has 5 to 20 starchy elongated tubers. Each tuber may be 20-80 cm long and 5-10 cm in diameter. Average tuber weight is between 4 and 7 kg but specimens up to 40 kg have been recorded (Eccocrop, 2011). The number and size of tubers is highly variable between cultivars and growing conditions (Eccocrop, 2011; Lebot, 2009). There are more than 7000 cassava varieties

Utilisation

Cassava tubers can be eaten boiled, mashed, deep-fried, etc. and there are many food products based on cassava, such as tapioca (cassava starch), a worldwide food ingredient, *fufu* (cassava flour boiled in water) and *garri* (fermented cassava mash), the two last popular foods in West and Central Africa. Cassava tubers also provide starch for ethanol production (Kuijper et al., 2007). Other cassava products include the finger-like leaves, which are consumed as vegetables or used as feed (see the [Cassava foliage](#) datasheet) and numerous by-products (notably pomace and peels) of the cassava processing industries (from starch, ethanol and cassava food production, etc.), which are also potential feeds (see the [Cassava by-products](#) datasheet). Cassava flour unsuitable for human consumption is also recycled as animal feed (Boscolo et al., 2002a).

More than a third of cassava production is used for animal feeding (FAO, 2011):

- **Fresh roots**, whole, broken or sliced
- **Dried cassava chips**: cassava chips sun-dried or artificially dried
- **Cassava root meal**: ground cassava chips
- **Cassava pellets**: ground and pelletized cassava chips. **Hard pellets** are particularly compact industrial pellets.

Automatic translation

Sélectionner une langue

Feed categories

All feeds

Forage plants

- ▶ Cereal and grass forages
- ▶ Legume forages
- ▶ Forage trees
- ▶ Aquatic plants
- ▶ Other forage plants

Plant products/by-products

- ▶ Cereal grains and by-products
- ▶ Legume seeds and by-products
- ▶ Oil plants and by-products
- ▶ Fruits and by-products
- ▶ Roots, tubers and by-products
- ▶ Sugar processing by-products
- ▶ Plant oils and fats
- ▶ Other plant by-products

Feeds of animal origin

- ▶ Animal by-products
- ▶ Dairy products/by-products
- ▶ Animal fats and oils
- ▶ Insects

Other feeds

- ▶ Minerals
- ▶ Other products

Latin names

Plant and animal families

Plant and animal species

Resources

Broadening horizons

Literature search

Image search

Glossary

External resources

- ▶ Literature databases
- ▶ Feeds and plants databases
- ▶ Organisations & networks
- ▶ Books
- ▶ Journals

Distribution

A native to South America, cassava is now widespread throughout the tropics and subtropics, including Sub-Saharan Africa and South-East Asia. The main production areas are within 30°N and 30°S and from sea level to an altitude of 2000 m, depending on the latitude ([Ecoport, 2009](#)).

Optimal growth conditions are an annual average day-temperature over 18-20°C, annual rainfall ranging from 500 mm to 3500 mm, high solar radiation and light, well-drained and acid soils. Cassava may withstand light frosts at higher altitudes and cloudy conditions in the hot humid lowland equatorial belt. A hardy plant, cassava is highly tolerant to poor soil conditions, drought and pests ([Vongsamphanh et al., 2004](#)), but it does not grow well in heavy, rocky and gravelly soils. It is susceptible to waterlogged, saline and alkaline soils. Zinc deficiency should be avoided while very low P levels are well accepted.

Cassava root production has been increasing steadily since the 1960s and has surged since 2000 (increased 40% between 1997 and 2007, from 161 to 224 million tons). Its use in animal feeding has also increased from 25% of the crop in 1997 to 34% in 2007 (76 million tons). In 2010, 52% of cassava was produced in Africa, 33% in Asia and 15% in Latin America ([FAO, 2011](#)).

The production of cassava chips and pellets for animal feeding started in Thailand in the 1960s, fuelled by European demand for energy sources cheaper than cereal grains, which were then highly subsidized in the EU. Shipping expenses and European concerns about dust pollution motivated a shift from chips to pellets in the late 1960s and to hard pellets in the early 1980s. Cassava exports to Europe climbed until the mid-1980s (the Netherlands imported 45-50% of worldwide dried cassava), when the EU set importation quotas ([FAO, 2001a](#)). The European market gradually evaporated and was replaced in the mid-2000s by China, which now imports 85% of the dried cassava produced worldwide. Today, Thailand remains the major exporter of dried cassava (80% in 2009), far ahead of Vietnam (14%) ([FAO, 2011](#)).

Processes

Fresh cassava tubers, particularly high-quality ones, are very perishable. They deteriorate within two or three days of harvest and therefore must be processed quickly ([Müller et al., 1975](#); [Tewe, 1992](#)).

Tubers intended for industrial animal feeding are sliced and dried, and then usually ground or pelletized. The technologies used at different scales of chip and pellet production are similar and cassava chips can be produced by simple techniques in the household or village as well as on a large mechanized scale. The selection of a technology depends on the amount of cassava to be processed, the availability of capital and labor cost, as well as the availability of relatively cheap energy ([Hahn et al., 1992](#)).

The first step is usually washing, followed by peeling. The roots are then sliced, either by hand or mechanically. Cassava chips may have different sizes and shapes, rectangular, cubic, thick sliced, depending on the slicing and drying methods. Drying may be natural or artificial. Sun-drying is done on concrete floors or on trays. Sun-drying is a very labor intensive operation, requiring about 35-40 laborers per hectare of drying floor. Chips dried on trays look better and are more uniformly dried than those dried on concrete floors. Artificial drying is done using static or moving bed dryers, or rotary dryers. Cassava chips can be sold directly, ground into cassava meal, or pelletized. During pelletizing, chips are heated and moistened and then forced into continuous die presses. Pelletizing results in a product that is 25-40% denser, more uniform, more durable, less dusty and easier to handle ([Hahn et al., 1992](#)).

Because peeling operations require time, alternative methods to produce chips and pellets without peeling have been developed. One such method consists in grating and chopping unpeeled tubers, mixing them with cassava foliage in a 4:1 ratio and passing the mixture through a pelletizer ([Tewe, 2004](#)).

In humid places where sun-drying is not easy, cassava roots can be ensiled alone (clean cassava roots + 0.5% salt) or mixed with rice straw or cassava leaves ([Le Duc Ngoan et al., 2002](#); [Premkumar et al., 2001](#); [Kavana et al., 2005](#)).

Forage management

Cassava is generally propagated by stem cuttings. However, under natural conditions as well as in plant breeding, propagation by seed is common and farmers in Africa are known to occasionally use spontaneous seedlings for subsequent planting ([Lokko et al., 2007](#)). Starch accumulation within the tubers occurs some 180-200 days after planting when they begin to thicken and store large quantities of starch. Because older tubers have the highest starch content, the best harvest period ranges from 9 to 24 months after planting. Cassava roots for animal feeding are commonly harvested from the 9th to the 12th month after planting ([Kuiper et al., 2007](#); [Régnier, 2011](#); [Gomez, 1991](#)). Harvesting is the most expensive part of cassava production. In order to improve tuber preservation, stems and leaves are cut two weeks before harvesting, leaving only a few centimetres of stems above ground. Uprooting must be done carefully because damaged tubers spoil readily ([Kuiper et al., 2007](#)). In 2009, the average worldwide tuber yield was 13 t/ha ([FAO, 2011](#)).

Environmental impact

Most cassava is produced by smallholder farmers living in marginal and fragile environments, and particularly on erosion-prone, acid and infertile soils. This ability to produce on poor soils, where most other crops would fail, has given cassava a reputation as a safeguard against food scarcity. However, there are serious environmental concerns about cassava production ([FAO, 2001b](#)). For the environmental impact of cassava processing, see the [Cassava by-products](#) datasheet.

Soil nutrient depletion

Cassava production can be detrimental to soil fertility through crop removal of nutrients. Due to the low value of cassava products, the application of manure and chemical fertilizers, which could easily correct nutrient depletion, may not be economically justified or affordable for smallholders. However, it should be noted that, at current yield levels, soil nutrient depletion by cassava is lower than depletion caused by other crops ([FAO, 2001b](#)).

Erosion

Cassava production can result in serious erosion when the crop is grown on slopes or on light soils. Good agronomic practices (adequate fertilizer, closer plant spacing, planting on contour ridges, intercropping, reduced tillage), used alone or in combination, can reduce erosion by 50-90%. Properly managed cassava production on slopes does not necessarily cause erosion ([FAO, 2001b](#)).

Water pollution

It is considered unlikely that cassava production results in water pollution, as it is grown mainly by poor farmers who apply no or very low rates of fertilizers, pesticides and herbicides. However, this may change in the future ([FAO, 2001b](#)).

Biodiversity

Cassava production does not seem to have had widespread effects on biodiversity, though some local situations may merit attention, such as deforestation in the north-east of Thailand or the competition with native cassava species in Latin America ([FAO, 2001b](#)).

Datasheet citation

Heuzé V., Tran G., Archimède H., Régnier C., Bastianelli D., Lebas F., 2016. *Cassava roots*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/527>. Last updated on April 12, 2016, 10:16

English correction by Tim Smith (Animal Science consultant) and H  l  ne Thiollet (AFZ)

Image credits

● David Monniaux ● Denis Bastianelli, CIRAD ● Vincent Porphyre, CIRAD ● Vincent Porphyre, CIRAD ● Fran  ois Lebas



Cassava roots

[Description](#) [Nutritional aspects](#) [Nutritional tables](#) [References](#)

Nutritional attributes

Cassava roots contain a large amount of starch, ranging from 70 to 85% DM, which increases with the stage of harvesting (Régnier, 2011; Ly, 1998). Cassava roots are therefore considered as an energy feed. However, their protein content (typically < 3%) is lower than that of cereal grains. Cassava can be substituted for cereals at high level in rations for all classes of livestock and poultry, provided that it is supplemented with a nitrogen source. The fibre content is also extremely low (NDF < 10% DM), which makes cassava roots highly digestible in all livestock species. HCN content may or may not be a problem, depending on the variety, process and livestock species.

Potential constraints

The cassava plant contains 2 glycosides, linamarin (80% of total glycoside) and lotaustralin (20%), which are acted upon by a cell-wall enzyme to liberate hydrogen cyanide (HCN), which is lethal to animals. HCN concentrations depend on cultivar, environmental conditions, plant age, number of harvest (for the foliage) and on the organ considered. While there is a continuous gradient of HCN content between varieties (Peroni et al., 2007), cassava varieties are usually divided into two groups:

- **Bitter varieties** have roots containing 0.02-0.03% HCN (DM basis) and fresh leaves containing up to 0.2% HCN (DM basis) (Murugeswari et al., 2006). Values up to 0.22% DM have been reported in fresh roots (Smith, 1988). These varieties have to be processed before being fed to animals.
- **Sweet varieties** have roots containing less than 0.01% HCN and fresh leaves containing about 0.1% HCN (DM basis) (Murugeswari et al., 2006). These varieties can be fed raw. Most commercial varieties belong to this group.

Bitter varieties often have longer and thicker roots than the sweet varieties, but there is no simple and safe method to assess HCN content.

Intensive use of cassava in animal feeding is possible after removal of the cyanogenic glucosides. It is generally assumed that roots containing less than 0.01% (100 mg/kg) of HCN in the DM are safe for use in animal diets (Buitrago et al., 2002b). Hydrogen cyanide is easily destroyed by simple treatments and many detoxification processes have been tested. Drying or ensiling cassava roots are the main processes for detoxification and storage (Gomez et al., 1988b; Tewe, 1992). In South America, ground cassava roots are put in nets, and then washed and squeezed until the toxic substance is eliminated. The toxic elements can also be removed by cooking, or by drying slices of the roots for about two weeks. Sun-drying appeared to be more efficient than oven-drying (60°C) (Panigrahi et al., 1992; Tewe, 1992).

The presence of HCN makes cassava products insect-resistant and easy to store. Adding 15% cassava root meal to a concentrate feed also improves its resistance to pests (Göhl, 1982).

Ruminants

Both fresh and dried cassava roots are consumed by ruminants in different forms (chips, ground, pelleted). Cassava starch has a large amylopectin content (70%) making it a suitable energy source for ruminants when combined with non-protein nitrogen in feeds (Müller, 1977).

Fresh cassava

The use of fresh cassava roots of bitter varieties is limited by their HCN content. When properly processed, they may serve as a basic energy source for intensive cattle feeding (Müller et al., 1975).

Dried cassava

Dried cassava roots have given satisfactory results as the principal energy source in ruminant production systems (Göhl, 1982). Studies indicate that the inclusion of cassava, to partly replace cereal grains (maize, barley, sorghum) up to 30-40%, gave satisfactory animal performance with no negative effects on animal health in finishing beef and dairy cattle, growing goats and lambs (Chanjula et al., 2007; Wachirapakorn et al., 2001; Sommart et al., 2000; Holzer et al., 1997; Zinn et al., 1991; Göhl, 1982). When cassava tubers are supplemented with non-protein nitrogen, minerals, vitamins, and roughage, high performances have been registered with dairy and beef cattle, sheep and goats (Smith, 1988). Palatability can be enhanced by the addition of molasses if pelleting is not possible (Göhl, 1982).

The energy value of cassava roots is about 85-93% that of maize grain, depending on the quality and starch content of the roots (Sauvant et al., 2004). In beef cattle, dried cassava was found to be as digestible as steam-flaked maize, but more so than sorghum grain (Zinn et al., 1991; Holzer et al., 1997). Because of the rapid degradation of cassava starch in the rumen, split-feeding several times a day can help ensure efficient utilization of nitrogen-deficient basal feed (Smith, 1988).


Dairy cattle

In some experiments, replacing maize with cassava resulted in lower milk yields but also in lower production costs. However, milk yield increased in other feed trials. Using cassava as an energy supplement in grazing cows had a positive effect on milk yield (+20%). Supplementing a fresh or ensiled sugarcane-based diet with cassava did not change milk production (Smith, 1988).

Beef cattle

In beef cattle, including cassava pellets up to 65% of the DM does not seem to affect health, carcass quality or overall performance when the diets are carefully balanced (Göhl, 1982). There were no significant differences in the performance of

Automatic translation

 Sélectionner une langue ▼

Feed categories

All feeds

Forage plants

- ▶ Cereal and grass forages
- ▶ Legume forages
- ▶ Forage trees
- ▶ Aquatic plants
- ▶ Other forage plants

Plant products/by-products

- ▶ Cereal grains and by-products
- ▶ Legume seeds and by-products
- ▶ Oil plants and by-products
- ▶ Fruits and by-products
- ▶ Roots, tubers and by-products
- ▶ Sugar processing by-products
- ▶ Plant oils and fats
- ▶ Other plant by-products

Feeds of animal origin

- ▶ Animal by-products
- ▶ Dairy products/by-products
- ▶ Animal fats and oils
- ▶ Insects

Other feeds

- ▶ Minerals
- ▶ Other products

Latin names

Plant and animal families

Plant and animal species

Resources

Broadening horizons

Literature search

Image search

Glossary

External resources

- ▶ Literature databases
- ▶ Feeds and plants databases
- ▶ Organisations & networks
- ▶ Books
- ▶ Journals

Holstein-Friesian male calves (180 kg live weight) fed a mixed diet containing 80% concentrate when 40% of the grain was replaced by cassava, except for a small increase in DM intake. The average daily gain was 1200 g and the energy conversion efficiency into live weight was reduced by 8%, depending on the nature of the protein source ([Holzer et al., 1997](#)).

Sheep

Substitution of maize cobs with cassava (20%) in sheep fed pangola grass (*Digitaria eriantha*) hay improved digestibility, body weight gain, and rumen function ([Smith, 1988](#)). Supplementation by 20 to 80% increased the digestibility of a rice straw-based diet but reduced the digestibility of a molasses-urea-based diet ([Devendra, 1977](#)).

Goats

In goats, replacing maize grain by cassava roots reduced performance at substitution levels of 40% and 60% ([Smith, 1988](#)). Supplementation of *Gliricidia sepium* with cassava roots at 30 g DM/kg $W^{0.75}$ reduced growth rate ([Smith, 1988](#)). Improved digestibility and similar growth rate to a control diet were reported with a 75:25 *Gliricidia/Leucaena* mix supplemented with cassava roots at 15 or 30 g DM/kg $W^{0.75}$. This discrepancy between the two last studies was attributed to energy-nitrogen synchronization in the rumen ([Smith, 1988](#)).

Pigs

Due to their high starch content, cassava roots are an excellent source of energy for pigs and can be used in fresh, ensiled or dried forms ([Göhl, 1982](#)). The digestible energy value of dried cassava roots for pigs varies between 14.5 and 16.5 MJ/kg DM ([Sauvant et al., 2004](#); [Rostagno et al., 2005](#); [Régnier, 2011](#)). These variations can be attributed to differences in chemical composition, especially in the starch and fibre fractions ([Régnier, 2011](#)). Because cassava peels are more than twice as fibrous as the root pulp, peeling improves energy digestibility and energy content.

For growing-finishing pigs, it is possible to include up to 60% of dried cassava root in the diet. The inclusion rate depends on the growth stage of the pig, and also on the form of distribution. The maximum intake of cassava is about 100 g DM/kg $W^{0.60}$ for dried ground cassava ([Gomez, 1991](#); [Régnier, 2011](#)). Cassava root meal is a palatable ingredient in the diets of young pigs ([Göhl, 1982](#)).

The major concern in feeding cassava roots to pigs is the presence of HCN, especially in bitter cultivars. There is a negative relationship between HCN content and cassava root intake ([Régnier, 2011](#)).

Poultry

Dried cassava tubers can be efficiently used in poultry feeding. The problems related with cyanogenic compounds are overcome by the use of sweet varieties and/or proper post-harvest treatments as simple as sun-drying on a concrete floor ([Gomez et al., 1983b](#); [Chauynarong et al., 2009](#)). Other processes such as boiling, autoclaving and fermentation have been found to be efficient but are unnecessary when sun-drying is sufficient. However, grinding cassava results in high levels of fine particles which can reduce feed intake and possibly irritate respiratory organs ([Garcia et al., 1999](#)). Pelletting reduces dust and increases the bulk density, favoring an increase in feed intake especially in young animals.

The protein content of cassava is low, which requires correcting when formulating a diet. Particularly, sulphur-containing amino acids (methionine and cystine) have to be supplied in large amounts because they can be altered during the metabolization of HCN. The metabolizable energy value of good cassava meal (72% starch) is equivalent to that of maize ([Sauvant et al., 2004](#)). Lower quality cassava (less starch, more fibre) has lower ME values, and the ME of unpeeled cassava meal is reduced to about 85% that of maize ([Agwunobi et al., 2000](#)).

Cassava tubers have also been used together with cassava foliage as a whole plant feed ([Akinfala et al., 2002](#)) (see the [Cassava foliage](#) datasheet).

Broilers

In well-formulated diets, good quality cassava can be used at high levels in broilers without reducing performance ([Chauynarong et al., 2009](#); [Daghir, 2008](#)). For example the inclusion of 50% pelleted cassava results in performance comparable to that obtained with maize diets ([Stevenson et al., 1983](#)). With more than 30% unpeeled cassava meal in the diet, some authors report a reduction of feed intake, resulting in a non-significant decrease in growth while maintaining feed efficiency ([Mafouo Ngandjou et al., 2011](#)). Fine grinding (<1 mm) can decrease performance compared to coarse grinding ([Mafouo Ngandjou et al., 2010](#)). Feed consumption can be affected in young animals at high inclusion rates (50%) while older animals maintain their performance ([Brum et al., 1990](#)). Early reports of growth depression with cassava were probably due to high HCN levels or protein/amino acid deficiencies ([Chauynarong et al., 2009](#)).

While there is no hard limit on the inclusion level of high-grade cassava in pelleted diets for grower-finisher broilers, the low protein content of cassava limits its inclusion to 30-40% to meet dietary requirements. When diets are fed in meal form, inclusion of cassava should not be higher than 20-30%, particularly in young animals ([Buitrago et al., 2002a](#)). Lesser grade cassava root products, such as dusty, unpeeled, or high-HCN roots or roots processed with little or no quality control, should be used more carefully, and the inclusion rate should not exceed 20% of the diet.

Layers

High levels of cassava meal can be used in layer diets when the HCN level is low and the diet is balanced for protein and amino acids ([Buitrago, 1990](#)). Up to 50% cassava did not significantly decrease production, feed efficiency and body weight. Water consumption increased when cassava was fed in meal form, while this effect was not observed with pelletting. Egg mass produced was also improved by pelletting ([Stevenson, 1984](#)).

Unpeeled cassava meal could be included at 30% in layers, completely replacing maize grain in the diet without adverse effects, including on egg quality (weight, shell, Haugh units, etc.) ([Eruvbetine et al., 1997](#)). However, the low carotenoid content of cassava requires supplementation with natural or synthetic sources of pigments if the egg yolk colour is to be maintained ([Garcia et al., 1999](#)).

Good quality cassava meal can be used in layer diets without limit provided that the diet is properly balanced, especially with amino acids. As in broilers, lower quality cassava should not exceed 20-30% of the diet.

Geese

When cassava meal was included at high levels (up to 45%) in geese diets, feed intake was maintained but performance and feed efficiency decreased ([Sahle et al., 1992](#)).

Turkeys

While early research reported problems when including cassava in turkey diets ([Göhl, 1982](#)), no evidence of negative effects has since been found in scientific literature when correctly formulated diets are fed.

Rabbits

Sun-dried cassava chips are used by traditional farmers in tropical countries such as Ghana ([Mamattah, 1979](#)), Tanzania ([Mgheni, 1979](#)), Uganda ([Lukfahr, 1998](#)) and Nigeria ([Mailafia et al., 2010](#)). Cassava roots slices are also a common ingredient for complete rabbit feeds in many tropical countries such as Cameroon ([Fomunyan et al., 1984](#)) or Vietnam ([Doan Thi Gang et al., 2006](#)). The inclusion level is generally 25 to 30% of the diet.

Several studies have investigated the ability of sun-dried cassava root meal to replace cereal grains such as maize and barley, or other concentrate ingredients, in rabbit diets ([Ikurior et al., 1998](#); [Radwan et al., 1989](#)). When experimental diets are correctly balanced, no differences are observed in growth or reproduction performance with inclusion levels up to 20-30%. Including cassava roots in the diet does not affect the quality of rabbit meat (physico-chemical composition and meat acceptability) ([Omole et al., 1983](#); [Onifade et al., 1993](#); [Soliman, 1994](#); [Oso et al., 2010](#)). In experiments levels up to 45-50% have been tested without any adverse effects in growing rabbits ([Radwan et al., 1989](#)) or breeding does ([Eshiett et al., 1980](#)).

The main issue when formulating balanced rabbit diets with cassava roots is their very low protein content, a problem that is usually addressed by increasing the proportion of soybean meal in the diet ([Oke, 1978](#)). However, cassava protein contains low levels of sulphur-containing amino acids, particularly methionine, which is required to eliminate the HCN released by bacterial activity in the digestive tract. Supplementary protein must supply sufficient methionine to achieve this. For instance, supplementing a diet containing 20% cassava meal and cassava leaf meal, also a low-methionine ingredient, for growing rabbits, resulted in a growth rate 17% lower than that obtained with the control diet ([Abd-El-Baki et al., 1993](#)). Thus, the use of cassava roots in rabbit feeding requires at least one source of sulphur-containing amino acids (more than 3.8% of the protein) such as maize bran, or pure DL-methionine. The total sulphur-containing amino acid content of the complete diet should never be lower than 3.7-3.8% of the dietary protein.

The addition of palm oil (5%) seems to be an alternative way to reduce the negative influence of the cyanogenic glucosides ([Omole et al., 1983](#)). Supplemental oil delays the bacterial decomposition and therefore decreases HCN absorption ([Tewe, 1992](#)).

Cassava roots, even when sliced and dried, have a mild goitrogenic effect, as evidenced by the low levels of serum thyroxin and cholesterol and by enlarged thyroid glands ([Ratnakumar et al., 1992](#)). However, rabbits fed a 25% cassava root diet had an increased growth rate compared to that of the control group, which indicates that the goitrogenic effect could be negligible in growing rabbits ([Ratnakumar et al., 1992](#)).

Fish

Cassava roots have been tested in many fish species as an energy source.

Tilapia

Digestibility and energy values of cassava chips reported in the literature for cassava are highly variable. Authors agree on the apparent protein digestibility (88-90%) but give different DM digestibility values (70% vs. 78%) and digestible energy values (6.7 vs. 13.2 MJ/kg DM) ([Campeche et al., 2011](#); [Pezzato et al., 2004](#)). However, cassava flour, unsuitable for human consumption, was found to be highly digestible (91% for DM and 97% for protein) with a much higher digestible energy value than cassava chips (15.4 MJ/kg DM) ([Boscolo et al., 2002a](#)).

Cassava root meal was found to be suitable for replacing 50% of white or yellow maize grain in the diets of young Nile tilapias ([El-Baki et al., 1999](#)). Another experiment concluded that discarded cassava flour could be used in the feeding of Nile tilapia fingerlings up to an inclusion level of 24%, wholly replacing maize grain without a decrease in performance ([Boscolo et al., 2002b](#)). It was possible to feed tilapias with mixtures of 76-80% fresh or sun-dried cassava leaves, 12-16% rice bran and 5% cassava roots ([Chhay Ty et al., 2010](#)).

Cassava meal was found to be less palatable to tilapias than sunflower meal, maize grain, maize gluten meal and animal by-products, but more palatable than wheat, soybean meal and cottonseed meal ([Pereira-da-Silva et al., 2000](#)).

African catfish

In *Clarias gariepinus* fingerlings, replacement of 33 to 100% of maize grain by cassava flour resulted in a reduced performance ([Akegbejo-Samsons, 1999](#)). However, an economic analysis showed that cassava root meal could replace maize in the diet of hybrid African catfish *Clarias gariepinus* x *Heterobranchus longifilis* profitably up to 100% inclusion, with the optimal economic performance at 66% level of inclusion. HCN content increased with the level of cassava in the diet but was always within the tolerable range for the normal metabolism of the fish ([Abu et al., 2010a](#); [Abu et al., 2010b](#)).

Carp

In grass carp (*Ctenopharyngodon idella*) fingerlings, cassava meal could replace up to 100% maize grain (30% of the diet) with no detrimental effect on the final weight, final length, feed conversion ratio, condition index and survival rate ([Lacerda et al., 2005](#)).

In common carp (*Cyprinus carpio* L.) a diet containing 47% cassava root meal has a slightly lower energy digestibility (87 vs. 90%) than diets containing maize or wheat starch. Growth and feed conversion efficiency were not influenced by starch source. The DM, fat and energy content in carp given cassava meal was significantly lower than that of carp given maize or wheat starch ([Schwarz et al., 1993](#)).

Characids

In South American characids, black pacu (tambaqui), *Colossoma macropomum* and red pacu (*Piaractus brachypomus*), cassava root, plantain fruit and peach-palm fruit (*Bactris gasipaes*) gave a better growth performance than wheat bran and wheat middlings in diets containing 30% of the test ingredient ([Lochmann et al., 2009](#)). Sun-dried, ground cassava chips could be fed to *Colossoma macropomum* at 5% of body weight per day, together with commercial chicken feed given at 1% of body weight ([Souza et al., 1998](#)).

Crustaceans

Prawns

Cassava meal can be totally substituted for wheat flour in extruded white shrimp (*Litopenaeus vannamei*) diets without any adverse effects on performance. It also improved the immunity development of the animals (Songluk et al., 2010).

Feeding giant tiger prawns (*Penaeus monodon*) with the cooked meat of golden snails and cooked cassava chips (60:40 on fresh weight) yielded the highest net income, compared with maize alone, and helped address the problem of snail infestation in rice fields (Bombeo-Tuburan et al., 1995).

Cassava meal could replace 100% of maize grain (51% of the total diet) in the diet of Malaysian prawns (*Macrobrachium rosenbergii*) without detrimental effects (Correia et al., 1996; Gomes et al., 1996). Dry matter, protein and energy digestibilities in that species are about 47-54%, 74-77% and 44-45% respectively, with heated cassava being slightly more digestible than dried cassava (Gomes et al., 1997).

Crabs

In the mud crab species *Scylla paramamosain*, cassava meal included at 30 or 45% in the diet reduced digestibility of DM, protein and energy compared to maize flour, rice bran and soybean meal and was therefore a less valuable ingredient (Phuong Ha Truong et al., 2009).

Other species

African giant land snail (*Archachatina marginata*)

A mixture of cassava flour and groundnut cake was used successfully to feed African giant land snails (Amubode et al., 1995).

Datasheet citation

Heuzé V., Tran G., Archimède H., Régnier C., Bastianelli D., Lebas F., 2016. Cassava roots. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/527> Last updated on April 12, 2016, 10:16

English correction by Tim Smith (Animal Science consultant) and Hélène Thiollet (AFZ)

Image credits

● David Monniaux ● Denis Bastianelli, CIRAD ● Vincent Porphyre, CIRAD ● Vincent Porphyre, CIRAD ● François Lebas

[+](#) Share / Save [f](#) [t](#) [r](#)



Cassava roots

[Description](#) [Nutritional aspects](#) [Nutritional tables](#) [References](#)

Tables of chemical composition and nutritional value

- Cassava tubers, dehydrated
- Cassava tubers, fresh
- Cassava tubers, peeled, fresh

Avg: average or predicted value; SD: standard deviation; Min: minimum value; Max: maximum value; Nb: number of values (samples) used

Cassava tubers, dehydrated




Main analysis	Unit	Avg	SD	Min	Max	Nb
Dry matter	% as fed	87.6	1.2	84.3	92.8	3354
Crude protein	% DM	2.9	0.6	1.4	4.5	1040
Crude fibre	% DM	3.9	1.2	1.7	8.2	3083
NDF	% DM	8.0	2.4	4.1	12.0	63 *
ADF	% DM	5.4	1.7	2.6	8.4	60 *
Lignin	% DM	1.7	1.0	0.1	3.8	50 *
Ether extract	% DM	0.7	0.3	0.2	1.4	221
Ash	% DM	3.9	1.9	1.4	8.6	2081
Starch (polarimetry)	% DM	80.4	4.1	69.1	88.6	3118
Total sugars	% DM	2.4	1.0	0.9	4.6	112
Gross energy	MJ/kg DM	16.8	0.2	16.2	17.2	51 *

Minerals	Unit	Avg	SD	Min	Max	Nb
Calcium	g/kg DM	1.7	0.7	0.6	3.3	125
Phosphorus	g/kg DM	1.1	0.3	0.6	1.8	125
Potassium	g/kg DM	9.9	2.8	5.8	17.5	64
Sodium	g/kg DM	0.3	0.2	0.1	0.8	18
Magnesium	g/kg DM	0.9	0.2	0.5	1.4	55
Manganese	mg/kg DM	23	11	7	43	7
Zinc	mg/kg DM	33	38	7	116	7
Copper	mg/kg DM	5	2	2	7	7
Iron	mg/kg DM	24	29	6	57	3

Amino acids	Unit	Avg	SD	Min	Max	Nb
Alanine	% protein	5.3	1.1	3.8	6.8	7
Arginine	% protein	5.0	1.9	2.8	7.6	6
Aspartic acid	% protein	6.6	0.9	5.7	7.9	7
Cystine	% protein	1.6	0.6	0.4	2.7	8
Glutamic acid	% protein	12.5	3.7	6.5	17.5	7
Glycine	% protein	3.4	0.4	3.0	4.0	6
Histidine	% protein	3.6	2.0	1.3	6.6	6
Isoleucine	% protein	2.7	0.7	1.3	3.3	7
Leucine	% protein	5.1	0.8	4.2	6.1	6
Lysine	% protein	3.9	0.5	3.2	4.8	17
Methionine	% protein	1.6	0.3	0.9	1.9	10
Phenylalanine	% protein	2.9	0.9	1.3	3.9	7
Proline	% protein	3.3	0.5	2.8	3.8	3
Serine	% protein	3.2	0.8	1.7	4.2	7
Threonine	% protein	2.9	0.8	1.3	3.8	8
Tryptophan	% protein	0.8		0.5	1.0	2
Tyrosine	% protein	1.7	0.9	0.4	2.7	4
Valine	% protein	4.5	1.8	1.7	7.7	7

Ruminant nutritive values	Unit	Avg	SD	Min	Max	Nb
OM digestibility, Ruminant	%	88.8	2.8	87.9	95.1	5 *

Automatic translation

 Sélectionner une langue

Feed categories

All feeds

Forage plants

- ▶ Cereal and grass forages
- ▶ Legume forages
- ▶ Forage trees
- ▶ Aquatic plants
- ▶ Other forage plants

Plant products/by-products

- ▶ Cereal grains and by-products
- ▶ Legume seeds and by-products
- ▶ Oil plants and by-products
- ▶ Fruits and by-products
- ▶ Roots, tubers and by-products
- ▶ Sugar processing by-products
- ▶ Plant oils and fats
- ▶ Other plant by-products

Feeds of animal origin

- ▶ Animal by-products
- ▶ Dairy products/by-products
- ▶ Animal fats and oils
- ▶ Insects

Other feeds

- ▶ Minerals
- ▶ Other products

Latin names

Plant and animal families

Plant and animal species

Resources

Broadening horizons

Literature search

Image search

Glossary

External resources

- ▶ Literature databases
- ▶ Feeds and plants databases
- ▶ Organisations & networks
- ▶ Books
- ▶ Journals

Energy digestibility, ruminants	%	84.5	3.3	84.4	92.8	5 *
DE ruminants	MJ/kg DM	14.2				*
ME ruminants	MJ/kg DM	12.2	0.7	11.5	12.9	5 *
Nitrogen digestibility, ruminants	%	35.3				*
a (N)	%	65.0				1
b (N)	%	34.4				1
c (N)	h-1	0.260				1
Nitrogen degradability (effective, k=4%)	%	95				*
Nitrogen degradability (effective, k=6%)	%	93		64	93	2 *

Pig nutritive values	Unit	Avg	SD	Min	Max	Nb
Energy digestibility, growing pig	%	90.8	4.1	84.6	99.4	18 *
DE growing pig	MJ/kg DM	15.3	0.5	14.3	15.8	15 *
ME growing pig	MJ/kg DM	15.0	0.7	14.2	15.7	8 *
NE growing pig	MJ/kg DM	12.2				*
Nitrogen digestibility, growing pig	%	52.3	21.8	16.7	83.8	13

Poultry nutritive values	Unit	Avg	SD	Min	Max	Nb
AMEn cockerel	MJ/kg DM	15.1	0.4	13.8	15.1	4 *
AMEn broiler	MJ/kg DM	15.1	1.3	13.4	15.8	3 *

The asterisk * indicates that the average value was obtained by an equation.

References

AFZ, 2011; Agunbiade et al., 2004; Anderson et al., 1991; Arieli, 1992; Bach Knudsen, 1997; Bui Huy Nhu Phuc, 2003; Carré et al., 1986; Carvalho Junior et al., 2009; Chanjula et al., 2003; Chhay Ty et al., 2006; Chhay Ty et al., 2007; Chhay Ty et al., 2007; Chiv Phiny et al., 2008; Chou et al., 1973; CIRAD, 1991; Cirad, 2008; Dao Lan Nhi et al., 2001; De Boever et al., 1984; De Boever et al., 1988; De Boever et al., 1994; DePeters et al., 1992; Du Thanh Hang et al., 2009; Fetuga et al., 1976; Gowda et al., 2004; Guillaume, 1978; Holm, 1971; Hutagalung, 1977; Jondreville et al., 1991; Jongbloed et al., 1990; Jongbloed et al., 1993; Kendall et al., 1982; Kuan et al., 1982; Le Duc Ngoan et al., 2001; Ledin et al., 2002; Lekule et al., 1990; Malibougou et al., 1998; Morgan et al., 1980; Muindi et al., 1981; Nguyen Van Hao et al., 2001; Nguyen Van Thu et al., 2009; Noblet et al., 1989; Perez et al., 1981; Perez et al., 1984; Pham Ho Hai et al., 2009; Pozy et al., 1996; Premaratne et al., 1998; Rajaguru et al., 1985; Ranaweera et al., 1981; Ravindran et al., 1994; Sahle et al., 1992; Sonaiya et al., 1983; Stevenson et al., 1983; Stevenson, 1984; Szylit et al., 1977; Thim Sokha et al., 2008; URZ, 2009; Van Cauwenberghé et al., 1996; Vanthong Phengvichith et al., 2007; Vervaeke et al., 1989; Wainman et al., 1984

Last updated on 24/10/2012 00:45:26

Cassava tubers, fresh



Main analysis	Unit	Avg	SD	Min	Max	Nb
Dry matter	% as fed	37.6	5.7	31.9	46.2	11
Crude protein	% DM	2.6	0.9	1.4	4.6	12
Crude fibre	% DM	3.7	1.0	2.0	5.7	11
NDF	% DM	7.8				*
ADF	% DM	5.3				*
Lignin	% DM	1.6				*
Ether extract	% DM	0.8	0.3	0.4	1.5	12
Ash	% DM	2.8	0.8	2.1	4.8	12
Starch (polarimetry)	% DM	80.8				*
Gross energy	MJ/kg DM	17.1				*

Minerals	Unit	Avg	SD	Min	Max	Nb
Calcium	g/kg DM	1.6	0.3	1.0	2.0	8
Phosphorus	g/kg DM	1.2	0.5	0.2	1.9	8
Potassium	g/kg DM	7.7	2.4	5.2	11.7	7
Magnesium	g/kg DM	1.1	0.2	0.8	1.5	7

Amino acids	Unit	Avg	SD	Min	Max	Nb
Arginine	% protein	7.7				1
Histidine	% protein	1.5				1
Isoleucine	% protein	5.3				1
Leucine	% protein	5.6				1
Lysine	% protein	6.2				1
Methionine	% protein	0.6				1
Phenylalanine	% protein	3.5				1
Threonine	% protein	3.8				1
Tryptophan	% protein	0.5				1

Valine	% protein	4.5				1
Ruminant nutritive values						
	Unit	Avg	SD	Min	Max	Nb
OM digestibility, Ruminant	%	89.1				*
Energy digestibility, ruminants	%	85.0				*
DE ruminants	MJ/kg DM	14.5				*
ME ruminants	MJ/kg DM	12.4				*
Nitrogen digestibility, ruminants	%	31.4				*
Pig nutritive values						
	Unit	Avg	SD	Min	Max	Nb
Energy digestibility, growing pig	%	92.1				*
DE growing pig	MJ/kg DM	15.7				*
ME _n growing pig	MJ/kg DM	15.4				*
NE growing pig	MJ/kg DM	12.6				*
Poultry nutritive values						
	Unit	Avg	SD	Min	Max	Nb
AMEn cockerel	MJ/kg DM	15.2				*
AMEn broiler	MJ/kg DM	15.2				*

The asterisk * indicates that the average value was obtained by an equation.

References

CIRAD, 1991; Dongmeza et al., 2009; French, 1937; Maner et al., 1967; Oyenuga, 1968; Ramachandran et al., 1956

Last updated on 24/10/2012 00:43:56

Cassava tubers, peeled, fresh



Main analysis						
	Unit	Avg	SD	Min	Max	Nb
Dry matter	% as fed	28.5		28.5	28.5	2
Crude protein	% DM	2.2		1.7	2.6	2
Crude fibre	% DM	1.0		0.4	1.6	2
NDF	% DM	3.7				*
ADF	% DM	1.6				*
Lignin	% DM	0.0				*
Ether extract	% DM	0.6		0.5	0.7	2
Ash	% DM	3.8		2.4	5.2	2
Gross energy	MJ/kg DM	16.7				*
Minerals						
	Unit	Avg	SD	Min	Max	Nb
Calcium	g/kg DM	1.0				1
Phosphorus	g/kg DM	0.4				1
Ruminant nutritive values						
	Unit	Avg	SD	Min	Max	Nb
OM digestibility, Ruminant	%	93.7				*
Energy digestibility, ruminants	%	89.2				*
DE ruminants	MJ/kg DM	14.9				*
ME ruminants	MJ/kg DM	12.8				*
Nitrogen digestibility, ruminants	%	33.8				*
Pig nutritive values						
	Unit	Avg	SD	Min	Max	Nb
Energy digestibility, growing pig	%	95.6				*
DE growing pig	MJ/kg DM	16.0				*
ME _n growing pig	MJ/kg DM	15.7				*
NE growing pig	MJ/kg DM	12.8				*
Nitrogen digestibility, growing pig	%	78.2				*

The asterisk * indicates that the average value was obtained by an equation.

References

Oyenuga, 1968

Last updated on 24/10/2012 00:43:56

Datasheet citation

Heuzé V., Tran G., Archimède H., Régnier C., Bastianelli D., Lebas F., 2016. Cassava roots. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/527>. Last updated on April 12, 2016, 10:16

English correction by Tim Smith (Animal Science consultant) and H el ene Thiollet (AFZ)

Image credits

- David Monniaux
- Denis Bastianelli, CIRAD
- Vincent Porphyre, CIRAD
- Vincent Porphyre, CIRAD
- Fran ois Lebas


 Share / Save   





Cassava roots

[Description](#)
[Nutritional aspects](#)
[Nutritional tables](#)
[References](#)


References


Abd-El-Baki, S. M. ; Nowar, M. S. ; Bassuny, S. M. ; Hassona, E. M. ; Soliman, E. S., 1993. Cassava as new animal feed in Egypt. 3. Pelleted complete cassava feed for growing rabbits. *World Rabbit Science*, 1 (4): 139-145 

Abu, O. M. G. ; Sanni, L. O. ; Erondy, E. S. ; Akinrotimi, O. A., 2010. Economic viability of replacing maize with whole cassava root meal in the diet of hybrid cat-fish. Abu, O. M. G. ; Sanni, L. O. ; Erondy, E. S. ; Akinrotimi, O. A. 

Abu, O. M. G. ; Sanni, L. O. ; Erondy, E. S. ; Akinrotimi, O. A., 2010. Chemical composition and cyanide levels of hybrid catfish fed whole cassava root meal in replacement of maize. *J. Food Technol.*, 8 (2): 52-57 


Agwunobi, L. N. ; Okeke, J. E., 2000. Metabolisable energy of some improved cassava cultivars for broiler chicken. *Afr. J. Root Tuber Crops*, 4 (1), 35-37


Akegbejo-Samsons, Y., 1999. The use of cassava flour as a substitute for yellow maize in diets for *Clarias gariepinus* fingerlings. *J. Aquacult. Trop.*, 14 (3): 247-253 


Akinfala, E. O. ; Aderibigbe, A. O. ; Matanmi, O., 2002. Evaluation of the nutritive value of whole cassava plant as replacement for maize in the starter diets for broiler chicken. *Livest. Res. Rural Dev.*, 14 (6) 

Amubode, F. O. ; Ogogo, A. U., 1995. Performance of snails (*Archachatina marginata*) fed varying levels of calorie-protein supplementary diets. *Nigerian J. Forestry*, 24/25: 36-43


Blakley, R. L. ; Coop, I. E., 1949. The metabolism and toxicity of cyanide and cyanogenic glycosides in sheep. 11. Detoxification of hydrocyanic acid. *New Zeal. J. Sci. Technol.*, 31 (3) A: 1-16


Bombero-Tuburan, I. ; Fukumoto, S. ; Rodriguez, E. M., 1995. Use of the golden apple snail, cassava, and maize as feeds for the tiger shrimp, *Penaeus monodon*, in ponds. *Aquaculture*, 131 (1-2): 91-100 


Boscolo, W. R. ; Hayashi, C. ; Meurer, F., 2002. Apparent digestibility of the energy and nutrients of conventional and alternative foods for Nile tilapia (*Oreochromis niloticus*). *Rev. Bras. Zootec.*, 31 (2): 539-545 


Boscolo, W. R. ; Hayashi, C. ; Meurer, F., 2002. Cassava by-product meal (*Manihot esculenta*) on feeding of Nile tilapia (*Oreochromis niloticus* L.) fingerlings. *Rev. Bras. Zootec.*, 31 (2): 546-551 


Brum, P. A. R. de ; Guidoni, A. L. ; Albino, L. F. T. ; Cesar, J. S., 1990. Whole cassava meal in diets for broiler chickens. *Pesq. Agropec. Bras.*, 25 (10): 1367-1373

Buitrago, J. A. ; Gil Llanos, J. L. ; Patiño, B. O., 2002. Cassava in poultry nutrition. *Cuadernos Avícolas* 14, FENAVI-FONAV, Cali (Colombia). 44p. 

Buitrago, J. ; Ospina, B. ; Gil, J. L. ; Aparicio, H., 2002. Cassava root and leaf meal as the main ingredient in poultry feeding: some experiences in Colombia. In: Howeler, R.H. Cassava research and development in Asia: Exploring new opportunities for an ancient crop. 7th Regional Cassava Workshop, Oct 28- Nov 1, 2002 in Bangkok, Thailand 


Buitrago, J. A., 1990. The use of cassava in animal feeding. *Centro Internacional de Agricultura Tropical (CIAT), Cali (Colombia)*, 446 p. 


Campeche, D. F. B. ; Moraes, S. A. de ; Lima, V. T. ; Sousa, S. M. de N. ; Oliveira, S. T. L. de ; Souza, M. G. de ; Paulino, R. V., 2011. Chemical composition and apparent digestibility of feed found in the Brazilian semiarid region for tilapia rosa feeding on cultivation. *Ciencia Rural*, 41 (2): 343-348 

Cereda, M. P. ; Mattos, M. C. Y., 1996. Linamarin: the toxic compound of cassava. *J. Venomous Animals and Toxins, Botucatu*, 2 (1) 


Chanjula, P. ; Ngampongsai, W. ; Wanapat, M., 2007. Effects of replacing ground corn with cassava chip in concentrate on feed intake, nutrient utilization, rumen fermentation characteristics and microbial populations in goats. *Asian-Aust. J. Anim. Sci.* 20 (10): 1557-1566


Chapoutot, P., 1998. Étude de la dégradation *in situ* des constituants pariétaux des aliments pour ruminants. Thèse Docteur en Sciences Agronomiques, Institut National Agronomique Paris-Grignon, Paris (FRA), 1998/11/17.

Chara, J. D., 1992. Level of starch and root of cassava (*Manihot esculenta*) as energy source in the diet of Peking ducks (*Anas platyrhynchos*). *Livest. Res. Rural Dev.*, 4 (2): 13-19 


Chauynarong, N. ; Elangovan, A. V. ; Iji, P. A., 2009. The potential of cassava products in diets for poultry. *World Poultry Sci. J.*, 65 (1): 23-36 


Chedly, K. ; Lee, S., 1999. Silage from by-products for smallholders. *FAO Electronic Conference on Tropical Silage* 


Chhay Ty; Ly, J. ; Rodríguez, L., 2001. An approach to ensiling conditions for preservation of cassava foliage in Cambodia. *Livest. Res. Rural Dev.*, 13 (2) 


Chhay Ty; Borin, K. ; Sopharith, N. ; Preston, T. R. ; Aye, T. M., 2010. Effect of sun-dried and fresh cassava leaves on growth of Tilapia (*Oreochromis niloticus*) fish fed basal diets of rice bran or rice bran mixed with cassava root meal. *Livest. Res. Rural Dev.*, 22 (3): 43 

Chou, K. C. ; Nah, K. C. ; Muller, Z., 1973. Replacement of maize by high level of tapioca meal in rations for growing/finishing pigs. *Kajian Veterinaire, Malasya-Singapore*, 5 (1): 3-10

Chumpawadee, S. ; Chantiratikul, A. ; Chantiratikul, P., 2007. Chemical compositions and nutritional evaluation of energy feeds for ruminants using an *in vitro* gas production technique. *Pak. J. Nutr.*, 6 (6): 607-612 

Conceição, W. L. F. ; Figueirêdo, A. V. de ; Nascimento, H. T. S. dos ; Vasconcelos, V. R. ; Alves, A. A. ; Filho, L. A. D., 2009. Nutritional value of diets containing cassava scrapings for feedlot sheep. *Acta Scientiarum - Animal Sciences*, 31 (4): 397-402 

Correia, E. de S. ; Gomes, S. Z., 1996. Effect of substituting maize with cassava meal on growth, feed conversion and survival rate in Malaysian prawns (*Macrobrachium rosenbergii* De Man, 1879). *Rev. Bras. Zootec.*, 25 (4): 583-594 

Daghir, N. J., 2008. Poultry production in hot climates. Second Edition, Cabi Series, CABI 

Automatic translation

Feed categories

All feeds

Forage plants

- ▶ Cereal and grass forages
- ▶ Legume forages
- ▶ Forage trees
- ▶ Aquatic plants
- ▶ Other forage plants

Plant products/by-products

- ▶ Cereal grains and by-products
- ▶ Legume seeds and by-products
- ▶ Oil plants and by-products
- ▶ Fruits and by-products
- ▶ Roots, tubers and by-products
- ▶ Sugar processing by-products
- ▶ Plant oils and fats
- ▶ Other plant by-products

Feeds of animal origin

- ▶ Animal by-products
- ▶ Dairy products/by-products
- ▶ Animal fats and oils
- ▶ Insects

Other feeds

- ▶ Minerals
- ▶ Other products

Latin names

Plant and animal families

Plant and animal species

Resources

Broadening horizons

Literature search

Image search

Glossary

External resources

- ▶ Literature databases
- ▶ Feeds and plants databases
- ▶ Organisations & networks
- ▶ Books
- ▶ Journals

- Devendra, C., 1977. Cassava as a feed source for ruminants. In Cassava as animal feed. Proc. of Cassava as Animal Feed Workshop, Eds B. Nestel and M. Graham, 18-20 April 1977, University of Guelph, Ontario, Canada. IDRC: Ottawa, 107-119 
- Devendra, C., 1983. New dietary protein sources for animal production in South East Asia. Feed information and animal production. Proceedings of the Second Symposium of the International Network of Feed Information Centres. 1983, 479-483
- Dinh Van Binh; Bui Van Chinh; Preston, T. R., 1991. Molasses urea blocks as supplements for rabbits. Livest. Res. Rural Dev., 3 (2): 13-16 
- Doan Thi Gang; Khuc Thi Hue; Dinh Van Binh; Nguyen Thi Mui, 2006. Effect of Guinea grass on feed intake, digestibility and growth performance of rabbits fed a molasses block and either water spinach (*Ipomoea aquatica*) or sweet potato (*Ipomoea batatas* L) vines. Workshop on Forages for Pigs and Rabbits, 21-24 August 2006, MEKARN-CelAgrid 
- Du Thanh Hang ; Nguyen Quang Linh ; Everts, H. ; Beynen, A. C., 2009. Ileal and total tract digestibility in growing pigs fed cassava root meal and rice bran with inclusion of cassava leaves, sweet potato vine, duckweed and stylosanthes foliage. Livest. Res. Rural Dev., 21 (1) 
- Ecocrop, 2011. Ecocrop database. FAO 
- Ecoport, 2009. Ecoport database. Ecoport 
- Ecoport, 2011. Ecoport database. Ecoport 
- El-Baki, S. M. A. ; Ghoneim, S. I. ; El-Husseiny, H. M. ; El-Gendy, K. M. ; Marghany, M., 1999. Cassava as a new animal feed in Egypt. 9. Cassava root meal in Nile tilapia (*Oreochromis niloticus*) diets. Egyptian J. Nutr. Feeds, 2 (Special issue): 753-763
- Eruvbetine, D. ; Oguntona, E. B., 1997. Unpeeled cassava root meal in diets for laying hens. Trop. Agric. (Trinidad), 74 (4): 299 - 302 
- Eshiett, N. O. ; Ademosun, A. A. ; Omole, T. A., 1980. Effect of feeding cassava root meal on reproduction and growth of rabbits. J. Nutr., 110 (4) : 697-702 
- FAO, 2001. A review of cassava in Asia with country case studies on Thailand and Viet Nam. Proceedings of the validation forum on the global cassava development strategy, Volume 3. FAO, Rome, 26-28 april 2000 
- FAO, 2001. Strategic environmental assessment. An assessment of the impact of cassava production and processing on the environment and biodiversity. Proceedings of the validation forum on the global cassava development strategy, Volume 5. FAO, Rome, 26-28 April 2000 
- FAO, 2001. The global cassava development strategy and implementation plan. Proceedings of the validation forum on the global cassava development strategy, Volume1. FAO, Rome, 26-28 april 2000 
- FAO, 2004. A review of cassava in Latin America and the Caribbean with country case studies on Brazil and Colombia. Proceedings of the validation forum on the global cassava development strategy, Volume 4. FAO, Rome, 26-28 april 2000 
- FAO, 2005. A review of cassava in Africa with country case studies on Nigeria, Ghana, the United Republic of Tanzania, Uganda and Benin. Proceedings of the validation forum on the global cassava development strategy, Volume5. FAO, Rome, 26-28 april 2000 
- FAO, 2011. FAOSTAT. Food and Agriculture Organization of the United Nations 
- Fetuga, B. L. ; Oluyemi, J.A., 1976. The metabolizable energy of some tropical tuber meals for chicks. Poult. Sci., 55 (3): 868-873 
- Figuerola, M. ; Paneque, G. ; Marrero, L., 1986. Study of six varieties of cassava (*Manihot esculenta*) at two harvest times for pig feeding. Ciencia y Tecnica en la Agricultura, Viandas Tropicales, 9 (2): 27-38
- Fomunyan, R. T. ; Adegbola, A. A. ; Oke, O. L., 1984. The reproductive, growth and carcass traits of rabbits fed cassava-based diets supplemented with palm oil. Food Chem., 14 (4) : 263-272 
- French, M. H., 1937. The nutritive value of cassava roots. Rep. vet. Dep., Tanganyika, 81-82
- Garcia, M. ; Dale, N., 1999. Cassava root meal for poultry. J. Appl. Poult. Res., 8 (1): 132-137 
- Garcia Gallego, M. ; Bazoco, J. ; Akharbach, H. ; Suarez, M. D. ; Sanz, A., 1994. Utilization of different carbohydrates by the European eel (*Anguilla anguilla*). Aquaculture, 124 (1/4): 99-108 
- Göhl, B., 1982. Les aliments du bétail sous les tropiques. FAO, Division de Production et Santé Animale, Roma, Italy 
- Gomes, S. Z. ; Correia, E. de S., 1996. Effect of substituting maize with cassava meal on voluntary intake of DM by Malaysian prawns (*Macrobrachium rosenbergii* De Man, 1879). Rev. Bras. Zootec., 25 (4): 595-604 
- Gomes, S. Z. ; Pena, M. del C. G., 1997. Apparent digestibility of cassava (*Manihot esculenta*) by freshwater prawn (*Macrobrachium rosenbergii*). Rev. Bras. Zootec., 26 (5): 858-862 
- Gomez, G. ; Valdivieso, M., 1983. Cassava meal for baby pig feeding. Nutr. Rep. Int., 28 (3): 547-558
- Gomez, G. ; Valdivieso, M. ; Santos, J. ; Hoyos, C., 1983. Evaluation of cassava root meal prepared from low or high cyanide containing cultivars in pig and broiler diets. Nutr. Rep. Int., 28 (4): 693-704
- Gomez, G. ; Noma, A. T., 1986. The amino acid composition of cassava leaves, foliage, root tissues and whole root chips. Nutr. Rep. Int., 33 (4): 595-601
- Gomez, G. ; Valdivieso, M. ; Santos, J., 1988. Cassava whole root chips silage for growing finishing pigs. Nutr. Rep. Int., 37 (5): 1081-1092. Centro Internacional de Agricultura Tropical (CIAT), Cali, Colombia.
- Gomez, G. ; Valdivieso, M., 1988. The effects of ensiling cassava whole root chips on cyanide elimination. Nutr. Rep. Int., 37 (6): 1161-1166
- Gomez, G., 1991. Use of cassava products in pig feeding. Pig News and Information 12, 387-390
- Hahn, S. K. ; Reynolds, L. ; Egbunike, G. N., 1992. Cassava as livestock feed in Africa. Proc. IITA/ILCA/Univ. of Ibadan Workshop on the Potential Utilization of Cassava as Livestock Feed in Africa, 14-18 November 1988, Ibadan, Nigeria 
- Hershey, C., 1994. Manihot genetic diversity. In: International Network for Cassava genetic resources. Report on the first meeting of the International meeting for cassava genetic resources, CIAT, Cali, Colombia 18-23 august 1992. International Crop Network series N°10. International Plant genetic Re
- Holzer, Z. ; Aharoni, Y. ; Lubimov, V. ; Brosh, A., 1997. The feasibility of replacement of grain by tapioca in diets for growing-fattening cattle. Anim. Feed Sci. Technol., 64: 133-141 

- Hongthong Phimmasan; Ledin, I., 2005. Effect of supplementing on-farm a diet based on maize, rice bran and cassava chip with Stylo 184 or native grass on feed intake and growth in rabbits. MSc thesis, Swedish University of Agricultural Sciences, Department of Animal Nutrition and Management, Uppsala, Sweden, 2005
- Hongthong Phimmasan; Ledin, I., 2005. Effect of supplementing a diet based on maize, rice bran and cassava chip with three different improved forages on feed intake, digestibility and growth in rabbit. MSc thesis, Swedish University of Agricultural Sciences, Department of Animal Nutrition and Management, Uppsala, Sweden, 2005
- Hongthong Phimmasan, 2005. Evaluation of tropical forages as feeds for growing rabbits. MSc thesis, Swedish University of Agricultural Sciences, Department of Animal Nutrition and Management, Uppsala, Sweden, 2005
- Hutagalung, R. I., 1977. Additives other than methionine in cassava diets. In: Cassava as animal feed, Proceedings of a workshop held at the University of Guelph, 18-20 April 1977, 18-32
- Ikurior, S. A. ; Akem, J. D., 1998. Replacing maize with cassava root meal or its mixture with brewers yeast slurry in rabbit diets. Nigerian J. Anim. Prod., 25 (1-2) : 31-35
- Kanjanapruthipong, J., 1998. The use of cassava in cattle feeding. Clayuca (Latin america and caribbean consortium to support Cassava research and development) - Thai Tapioca Development Institute
- Kantho, U. ; Juttupompong, S., 2002. Clean cassava chips for animal feeding in thailand. In: Howeler, R.H. Cassava research and development in Asia: Exploring new opportunities for an ancient crop. 7th Regional Cassava Workshop, Oct 28- Nov 1, 2002 in Bangkok, Thailand
- Kavana, P. Y. ; Mtunda, K. ; Abass, A. ; Rweyendera, V., 2005. Promotion of cassava leaves silage utilization for smallholder dairy production in Eastern coast of Tanzania. Livest. Res. Rural Dev., 17 (4)
- Kiura, J.N. ; Bimbuzi, S. ; Mwakina D. ; Furaha, G., 2008. Processing cassava root for dairy cattle feeding. Kari Information Brochures II, Kenya Agricultural Research Institute
- Kiura, J. N. ; Ndung'u, J. M. ; Muli, B. M., 2010. Processing cassava into chips in coastal Kenya: commercial potential in the future. 12th KARI Biennial Scientific Conference: Transforming Agriculture for improved livelihoods through Agricultural Product Value Chains. Kenya Agricultural Research Institute
- Kuiper, L. ; Ekmecki, B. ; Hamelink, C. ; Hettinga, W. ; Meyer, S. ; Koop, K., 2007. Bio-ethanol from cassava. Project number: PBIONL062937. Ecofys Netherlands BV, Utrecht
- Lacerda, C. H. F. ; Hayashi, C. ; Soares, C. M. ; Boscolo, W. R. ; Kavata, L. C. B., 2005. Replacement of corn *Zea mays* L. by cassava *Manihot esculenta* crants meal in grass-carp *Ctenopharyngodon idella* fingerlings diets. Acta Scientiarum - Animal Sciences, 27 (2): 241-245
- Le Duc Ngoan; Nguyen Thi Hoa Ly, 2002. The use of cassava roots and leaves for feeding pigs in Vietnam. In: Howeler, R.H. Cassava research and development in Asia: Exploring new opportunities for an ancient crop. 7th Regional Cassava Workshop, Oct 28- Nov 1, 2002 in Bangkok, Thailand
- Lebot, V., 2009. Tropical root and tuber crops: cassava, sweet potato, yams and aroids. Crop production science in horticulture (17), CAB books, CABI, Wallingford, UK
- Lekule, F. P. ; Just, A. ; Mtenga, L. A., 1988. The responses in growth and carcass quality of barrows and gilts to diets of local feeds. E. Afr. Agric. For. J., 53 (3): 105-109
- Liu Jian Ping; Zhuang Zhong Tang, 2001. The use of dry cassava roots and silage from leaves for pig feeding in Yunnan province of China. In: Howeler, R.H.; Tan, S. L. (Eds). Cassava's potential in Asia in the 21th century: present situation and future research and development needs. Proc. 6th Regional Workshop, Ho Chi Minh City, Vietnam, February 21-25, 2000
- Lochmann, R. ; Chen, R. G. ; Chu-Koo, F. W. ; Camargo, W. N. ; Kohler, C. C. ; Kasper, C., 2009. Effects of carbohydrate-rich alternative feedstuffs on growth, survival, body composition, hematology, and nonspecific immune response of black pacu, *Colossoma macropomum*, and red pacu, *Piaractus brachypomus*. J. World Aquacult. Soc., 40 (1): 33-44
- Lokko, Y. ; Okogbenin, E. ; Mba, C. ; Dixon, A. ; Raji, A. ; Fregene, M., 2007. Cassava. In: Chittaranjan Kole, 2007. Pulses, Sugar and Tuber Crops. Genome Mapping and Molecular Breeding in Plants, Volume 3. Springer
- Longe, O. G. ; Tona, G. O., 1988. Metabolizable energy values of some tropical feedstuffs for poultry. Trop. Agric. (Trinidad), 65 (4):358-360
- Lukefahr, S. D., 1998. Rabbit production in Uganda : Potential *versus* opportunity. World Rabbit Science, 6 (3-4): 331-340
- Ly, J., 1998. Cassava roots (*Manihot esculenta* Crantz) for pigs; A short review on its nutrient content. Revista Computadorizada de Produccion Porcina, 5: 1-13
- Machin, D. ; Nyvold, S., 1992. Roots, tubers, plantains and bananas in animal feeding. Proceedings of the FAO Expert Consultation held in CIAT, Cali, Colombia 21–25 January 1991; FAO Animal Production and Health Paper - 95
- Madalla, N., 2008. Novel feed ingredients for Nile tilapia (*Oreochromis niloticus* L.) . In: PhD Thesis, Inst. Aquaculture, Univ. Stirling, Scotland, UK
- Mafouo Ngandjou, H. ; Tegua, A. ; Mube, H. K. ; Diarra, M., 2010. Effect of cassava flour particle size as alternative food energy source on broiler growth parameters. Livest. Res. Rural Dev., 22 (11)
- Mafouo Ngandjou, H. ; Tegua, A. ; Kana, J. R. ; Mube, H. K. ; Diarra, M., 2011. Effect of the level of incorporation of cassava flour in the diet on broiler growth parameters. Livest. Res. Rural Dev., 23 (4)
- Mailafia, S.; Onakpa, M. M.; Owoleke, O. E., 2010. Problems and prospects of rabbit production in Nigeria - A review. Bayero Journal of Pure and Applied Science, 3 (2): 20-25
- Mamattah, N., 1979. Sociological aspects of introducing rabbits into farm practices. Trop. Anim. Prod., 4 (3) : 295
- Maner, J. N. ; Buitrago, J. ; Jimenez, I., 1967. Utilisation of yuca in swine feeding. Proc. Int. Symp. Tropical Root Crops, Trinidad, 2, Section VI, 62-71
- Mgheni, M., 1979. Rabbit husbandry in Tanzania. Trop. Anim. Prod., 4 (3): 292
- Min Wang; Yuan Hu; Zhiliang Tan; Shaoxun Tang, Zhihong Sun; Xuefeng Han, 2008. In situ ruminal phosphorus degradation of selected three classes of feedstuffs in goats. Livest. Sci. 117 (2-3): 233-237
- Moore, C. P. ; Cock, J. H., 1985. Cassava forage silage as a feed source for zebu calves in the tropics. Trop. Agric. (Trinidad), 62 (2): 142-144
- Müller, Z. O. ; Chou, K. C. ; Nah, K.C., 1975. Cassava as a total substitute for cereals in livestock and poultry rations. Proceedings of the 1974 Tropical Products Institute Conference, 1–5 April, 85–95
- Müller, Z. O., 1977. Improving the quality of cassava root and leaf product technology. In: Cassava as animal feed. Proceeding, Cassava as animal feed Workshop, Nestel, B.; Graham, M. 18-20 april 1977 University of Guelph, Ontario, Canada. IDRC: Ottawa, 120-126

- Murugesrawi, R. ; Balakrishnan, V. ; Vijayakumar, R., 2006. Studies to assess the suitable conservation method for tapioca leaves for effective utilization by ruminants. *Livest. Res. Rural Dev.*, 18 (3)
- Nanda, S. K. ; Jyothi, A. N. ; Balagopalan, C., 2002. Cassava waste treatment and residue management in India. In: Howeler, R.H. Cassava research and development in Asia: Exploring new opportunities for an ancient crop. 7th Regional Cassava Workshop, Oct 28- Nov 1, 2002 in Bangkok, Thailand
- Nang'ayo, F. ; Omanyua, G. ; Bokanga, M. ; Odera, M. ; Muchiri, N. ; Ali, Z. ; Werehire, P., 2005. A strategy for industrialisation of cassava in Africa. In: Proceedings of a small group meeting, 14-18 November 2005, Ibadan, Nigeria. Nairobi, Kenya: African Agricultural Technology Foundation
- Nwokolo, E., 1987. Leaf meals of cassava (*Manihot esculenta* Crantz) and siam weed (*Eupatorium odoratum* L.) as nutrient sources in poultry diets. *Nutr. Rep. Int.*, 36 (4): 819-826
- Oke, O. L., 1978. Problems in the use of cassava as animal feed. *Anim. Feed Sci. Technol.*, 3 (4) : 345-380
- Omole, T. A. ; Onwudike, O. C., 1983. Effect of palm oil on the use of cassava peel meal by rabbits. *Trop. Anim. Prod.*, 8 (1): 27-32
- Onifade, A. A. ; Tewe, O. O., 1993. Alternative tropical energy feed resources in rabbit diets: growth performance, diet's digestibility and blood composition. *World Rabbit Science*, 1 (1) : 17-24
- Oso, A. O. ; Oso, O. ; Bamgbose, A. M. ; Eruvbetine, D., 2010. Utilization of unpeeled cassava (*Manihot esculenta*) root meal in diets of weaner rabbits. *Livest. Sci.*, 127 (2) : 192-196
- Ospina, B. ; Wheatley, C., 1992. Processing of cassava tuber meals and chips. In: Machin, D.; Nyvold, S., 1992. Roots, tubers, plantains and bananas in animal feeding. Proceedings of the FAO Expert Consultation held in CIAT, Cali, Colombia 21-25 January 1991; FAO Animal Production and Health Paper - 95
- Oyenuga, V. A., 1968. Nigeria's foods and foodstuffs. Ibadan, University Press
- Panditharatne, S. ; Allen, V. G. ; Fontenot, J. P. ; Jayasuriya, M. C. N., 1986. Ensiling characteristics of tropical grasses as influenced by stage of growth, additives and chopping length. *J. Anim. Sci.*, 63 (1): 197-207
- Panigrahi, S. ; Rickard, J. ; O'Brien, G. M. ; Gay, C., 1992. Effects of different rates of drying cassava root on its toxicity to broiler chicks. *Br. Poult. Sci.*, 33 (5):1025-1041
- Payne, M.; Owen, E.; Capper, B. S.; Wood, J. F. Radwan, M. A.H, 1988. Incorporation of grass silage, whole cereal grains, cassava and cottonseed meal into diets of rabbits kept in a simulated tropical environment. *Trop. Anim. Health Prod.*, 20 (4): 212-218
- Pereira-da-Silva, E. M. ; Pezzato, L. E., 2000. Response of Nile tilapia (*Oreochromis niloticus*) to the attraction and palatability of the used ingredients in the feeding of fishes. *Rev. Bras. Zootec.*, 29 (5): 1273-1280
- Peroni, N. ; Kageyama, P. Y. ; Begossi, A., 2007. Molecular differentiation, diversity, and folk classification of "sweet" and "bitter" cassava (*Manihot esculenta*) in Caicara and Caboclo management systems (Brazil). *Genetic Resources and Crop Evolution*, 54 (6): 1333-1349
- Pezzato, L. E. ; Miranda, E. C. de ; Barros, M. M. ; Furuya, W. M. ; Pinto, L. G. Q., 2004. Apparent digestibility of dry matter and crude protein and digestible energy of some alternative ingredients by Nile tilapia *Oreochromis niloticus*. *Acta Scientiarum - Animal Sciences*, 26 (3): 329-337
- Phuc, B. H. N. ; Lindberg, J. E., 2000. Ileal and total tract digestibility in growing pigs given cassava root meal diets with inclusion of cassava leaves, leucaena leaves and groundnut foliage. *Anim. Sci.*, 71: 301-308
- Phuong Ha Truong ; Anderson, A. J. ; Mather, P. B. ; Paterson, B. D. ; Richardson, N. A., 2009. Apparent digestibility of selected feed ingredients in diets formulated for the sub-adult mud crab, *Scylla paramamosain*, in Vietnam. *Aquacult. Res.*, 40 (3): 322-328
- Premkumar, T. ; Padmaja, G. ; Moorthy, S. N. ; Nanda, S. K.;George, M. ; Balagopalan, C., 2001. New cassava products of future potential in India. In: Howeler, R.H.; Tan, S. L. (Eds). Cassava's potential in Asia in the 21st century: present situation and future research and development needs. Proc. 6th Regional Workshop, Ho Chi Minh City, Vietnam, February 21-25, 2000
- Radwan, M. A. H. ; Partridge, G. G. ; Allan, S. J. ; Fordyce, R. A., 1989. Cassava root meal in diets for growing rabbits. *Trop. Anim. Health Prod.*, 21 (1) : 32-36
- Ramachandran, M. ; Phansalkar, S. V., 1956. Essential amino-acid composition of certain vegetable foodstuffs. *Indian J. med. Res.*, 44: 501-509
- Raposo de Medeiros, S. ; Machado, P. F., 1993. Effect of the replacement of steam treated sugar cane bagasse by milo on ruminal fermentation in bovines and *in vivo* digestibility in sheep. *Livest. Res. Rural Dev.*, 5 (2): 16-24
- Raposo de Medeiros, S. ; Machado, P. F., 1993. Effect of the replacement of steam treated sugarcane bagasse by milo upon performance of finishing cattle. *Livest. Res. Rural Dev.*, 5 (2): 25-30
- Ratnakumar, J. N. ; Rajan, A., 1992. Goitrogenic effect of cassava in broiler rabbits. *Indian J. Anim. Sci.*, 62 (7): 670-676
- Ravindran, V. ; Kornegay, E. T. ; Notter, D. R. ; Rajaguru, A. S. B., 1984. Utilization of cassava leaf meal in swine diets. *Anim. Sci. Res. Report*, Virginia Agricultural Experimental Station. No. 4, 92-96
- Ravindran, V. ; Kornegay, E. T. ; Rajaguru, A. S. B. ; Notter, D. R., 1987. Cassava leaf meal as a replacement for coconut oil meal in pig diet. *J. Sci. Food Agric.*, 41 (1): 45-53
- Régnier, C., 2011. Valorisation des ressources alimentaires tropicales (feuilles et tubercules) chez le porc. Thèse (INRA Antilles-Guyane, Unité de Recherches Zootechniques – URZ)
- Reynolds, L. ; Adediran, S. O., 1987. The effects of browse supplementation on the productivity of West African Dwarf sheep over two reproductive cycles. Goat production in the humid tropics. Proceedings of a workshop at the University of Ife, Ile Ife, Nigeria, 20 24 July 1987
- Rickard, J. E., 1986. Tannin levels in cassava, a comparison of methods of analysis. *J. Sci. Food Agric.*, 37 (1): 37-42
- Rostagno, H. S. ; Teixeira, A. ; Donzele, J. L. ; Gomes, P. C. ; De Oliveira, R. F. M. ; Lopes, D. C. ; Ferreira, A. J. P. ; Toledo Barreto, S. L., 2005. Brazilian Tables for Poultry and Swine: composition of feedstuffs and nutritional requirements. Universidade Federal de Viçosa, Departamento de Zootecnia, MG, Brazil
- Sahle, M. ; Coleou, J. ; Haas, C., 1992. Nutritional value of cassava meal in diets for geese. *Anim. Feed Sci. Technol.*, 36 (1-2): 29-40
- Sauvant, D.; Perez, J. M.; Tran, G., 2004. Tables INRA-AFZ de composition et de valeur nutritive des matières premières destinées aux animaux d'élevage: 2ème édition. ISBN 2738011586, 306 p. INRA Editions Versailles
- Schwarz, F. J.; Kirchgessner, M., 1993. Digestibility, growth and carcass composition of carp (*Cyprinus carpio* L.) fed different starches. *Arch. Tierernähr.*, 43 (3): 275-282
- Senez, J. C. ; Raimbault, M. ; Deschamps, F., 1983. Protein enrichment of starchy substrates by solid state fermentation.

Food and Nutrition Bulletin. 1983, Suppl. 7, 52-61

Smith, O. B., 1988. A review of ruminant responses to cassava-based diets. In: Hahn, S. K.; Reynolds, L., Egbunike, G. N. (Eds). Cassava as livestock in Africa.

Soliman, M. A., 1994. A study of some factors affecting rabbits meat quality. Egyptian J. Rabbit Sci., 4 (1) : 113-122

Sommart, K.; Wanapat, M.; Rowlinson, P.; Parker, D. S.; Climee, P.; Panishying, S., 2000. The use of cassava chips as an energy source for lactating dairy cows fed with rice straw. Asian-Aust. J. Anim. Sci. 13: 1094-1101

Songluk, K.; Kanto, U.; Juttapornpong, S.; Jintasathaporn, O., 2010. Effect of cassava meal on growth performance and immunological system in white shrimp (*Litopenaeus vannamei*). J. Agric. Res. Ext., 27 (3): 39-46

Souza, R. A. L. de; Castro Filho, B. O. de; Rodrigues, M. de J. J.; Peret, A. C.; Teixeira, R. N. G., 1998. Growth of tambaqui fish, *Colossoma macropomum* (Cuvier, 1818) (Pisces-Characidae), in tanks with ground manioc as food. Boletim da Faculdade de Ciências Agrárias do Para, Brazil, 29: 23-31

Stevenson, M. H.; Graham, W. D., 1983. The chemical composition and true metabolisable energy content of cassava root meal imported into Northern Ireland. J. Sci. Food Agric., 34 (10): 1105-1106

Stevenson, M. H., 1984. The nutritional value of cassava root meal in laying hen diets. J. Sci. Food Agric., 35: 36-40

Tewe, O. O., 1992. Detoxification of cassava products and effects of residual toxins on consuming animals. In: Machin, D.; Nyvold, S., 1992. Roots, tubers, plantains and bananas in animal feeding. Proceedings of the FAO Expert Consultation held in CIAT, Cali, Colombia 21–25 January 1991; FAO Animal Production and Health Paper - 95

Tewe, O. O., 2004. The global cassava development strategy: cassava for livestock feed in Sub-Saharan Africa. IFAD and FAO

Thomas, K.; Singh, R. A., 1985. Feeding pigs in tropics. 1. Effect of plane of feeding and feed particle size on growth. Kerala J. Vet. Sci., 15 (2): 51-60

Tudor, G. D.; McGuigan, K. R.; Norton, B. W., 1985. The effects of three protein sources on the growth and feed utilization of cattle fed cassava. J. Agric. Sci., 104: 11 - 18.

Ugwu, L. L. C.; Asogwa, M. O.; Mgbenka, B. O., 2004. Influence of dietary levels of cassava (*Manihot esculenta*) peel meal on feed efficiency and productive protein value of young tilapia (*Oreochromis niloticus*, Trewavas). J. Sustain. Agric. Environ., 6 (2): 148-156

Van Eys, J. E.; Pulungan, H.; Rangkuti, M.; Johnson, W. L., 1987. Cassava meal as supplement to napier grass diets for growing sheep and goats. Anim. Feed Sci. Technol., 18 (3): 197-207

Vongsamphanh, P.; Wanapat, M., 2004. Comparison of cassava hay yield and chemical composition of local and introduced varieties and effects of levels of cassava hay supplementation in native beef cattle fed on rice straw. Livest. Res. Rural Dev., 16 (8)

Wachirapakorn, C.; Wanapat, M.; Sornsungnern, N.; Kowsuwan, S., 2001. Optimum cassava root chip levels in lactating cow diets. International Workshop (July 23-25 2001) in Khon Kaen University, Thailand - Current Research and Development on Use of Cassava as Animal Feed. Mekarn

Wanapat, M.; Prasertsuck, S.; Chantai, S., 1985. Effects of ensiling rice straw with urea and supplementing with dried cassava leaves on digestion by water buffaloes. Trop. Anim. Prod., 10 (1): 44-49

Zinn, R. A.; De Peters, E. J., 1991. Comparative feeding value of tapioca pellets for feedlot cattle. J. Anim. Sci., 69: 4726-4733

148 references found

Datasheet citation

Heuzé V., Tran G., Archimède H., Régnier C., Bastianelli D., Lebas F., 2016. Cassava roots. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/527>. Last updated on April 12, 2016, 10:16

English correction by Tim Smith (Animal Science consultant) and Hélène Thiollot (AFZ)

Image credits

● David Monniaux ● Denis Bastianelli, CIRAD ● Vincent Porphyre, CIRAD ● Vincent Porphyre, CIRAD ● François Lebas

[+](#) Share / Save [f](#) [t](#) [r](#)

