

---

# Are mycorrhizal-bacterial community networks impacted by the sexual type of their host tree?

Hervé Sanguin\*<sup>†1</sup>, Frédéric Mahé\*<sup>1</sup>, Pierre Tisseyre<sup>1</sup>, Estelle Tournier<sup>1</sup>, Lahcen Ouahmane<sup>2</sup>, Christine Le Roux<sup>1</sup>, Robin Duponnois<sup>1</sup>, Frédéric Médail<sup>3</sup>, Magda Bou Dagher-Kharrat<sup>4</sup>, Stefano La Malfa<sup>5</sup>, Alex Baumel<sup>3</sup>, and Yves Prin\*<sup>1</sup>

<sup>1</sup>Laboratoire des symbioses tropicales et méditerranéennes (LSTM) – Centre de coopération internationale en recherche agronomique pour le développement [CIRAD] : UMR82, Institut de recherche pour le développement [IRD] : UMR113 – Campus international de Baillarguet - TA 10 / J - 34398 Montpellier Cedex 5, France

<sup>2</sup>University of Cadi Ayyad (UCA) – UCAM, FSSM, Bd Moulay abdellah, 40000, Marrakesh, Morocco

<sup>3</sup>Institut méditerranéen de biodiversité et d'écologie marine et continentale (IMBE) – INEE, Université d'Avignon et des Pays de Vaucluse, Institut de recherche pour le développement [IRD] : UMR237, Aix Marseille Université, CNRS : UMR7263, INSB, INSU – Aix Marseille Université, Campus Etoile, Faculté St-Jérôme case 421 Av. . escadrille Normandie-Niemen 13397 MARSEILLE CEDEX 20, France

<sup>4</sup>University Saint-Joseph (USJ) – Rectorat de l'USJ Rue de Damas, BP 17-5208 - Mar Mikhaël Beyrouth - 1104 2020 Liban, Lebanon

<sup>5</sup>University of Catania, Department of Agriculture, Food and Environment (UniCt, Di3A) – Università degli Studi di Catania Piazza Università, 2 95131 Catania, Italy

## Abstract

The determination of tree sexual type during the production of seedlings in nursery constitutes a major issue for the set up of efficient orchards. In the case of carob trees (*Ceratonia siliqua*), a multi-use tree of high socio-economical added value in Mediterranean regions, the majority is dioecious but some are also hermaphrodite. Physiological differences between carob male and female have been described but the development of robust indicators of tree sexuality remains of particular interest for the carob sector.

Carob tree is highly dependent on arbuscular mycorrhizal (AM) symbiosis, and bacterial endophytes are also hypothesized to play a role in carob nutrition. Symbiotic communities are known to be key factors of plant nutrition and growth, affecting greatly plant physiology. The intimate connection between physiology and symbiotic community of carob may suggest a potential specificity of symbionts regarding the carob sexual type.

In the framework of the international project DYNAMIC (Deciphering sYmbiotic Networks in cArob-based MedIterranean agro-eCosystems), a metabarcoding approach was developed to characterized the symbiotic microbial diversity and assemblages associated to carob trees in the Mediterranean basin. The results may provide new guidelines for the determination of bio-indicators of tree sexual type and the design of more productive orchards.

**Keywords:** Symbiotic community, diversity, Carob, New Generation Sequencing

---

\*Speaker

<sup>†</sup>Corresponding author: herve.sanguin@cirad.fr