

Tropentag 2018

International Research on Food Security, Natural
Resource Management and Rural Development

Global food security and food safety: The role of universities

Book of abstracts

Editor: Eric Tielkes

Reviewers/scientific committee: Geert Baert, Jan Baetens,
Veerle Fievez, Geert Haesaert, Bernhard Freyer, Carl Lachat, Roel Merckx,
Eva Schlecht, Stijn Speelman, Barbara Sturm, Patrick Van Damme,
Gilbert Van Stappen, Wouter Vanhove, Annelies Verdoolaege,
Vladimir Verner, Kaat Verzelen, Ann Water-Bayers, Anthony Whitbread,
Stephan Winter

Editorial assistance: Mercedes Gotterbarm, Laetitia Samuel

Impressum

Bibliografische Information der Deutschen Nationalbibliothek

Die Deutsche Nationalbibliothek verzeichnet diese Publikation in der Deutschen Nationalbibliografie; detaillierte bibliografische Daten sind im Internet über <http://dnb.ddb.de> abrufbar.

Tropentag 2018: Global food security and food safety: the role of universities
Tielkes, E. (ed.) 1. Aufl. - Weikersheim: Margraf Publishers GmbH, 2018

© Margraf Publishers GmbH, Weikersheim
Kanalstraße 21, 97990 Weikersheim
Telefon: 07934 - 3071
Telefax: 07934 - 8156
www.margraf-publishers.eu

Alle Rechte vorbehalten. Ohne ausdrückliche Genehmigung des Verlages ist es nicht gestattet, das Buch oder Teile daraus auf fotomechanischem Weg (Fotokopie Mikrokopie) zu vervielfältigen.

The authors of the articles are solely responsible for the content of their contribution. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means without prior permission of the copyright owners.

ISBN: 978-3-8236-1760-0

Online-Version: <http://www.tropentag.de/>



Margraf Publishers GmbH

Oil Palm Root System Characteristics in a 12 Year Old Density Trial in Nigeria

REINOUT IMPENS¹, PATRICK VAN DAMME¹, XAVIER BONNEAU²

¹*Ghent University, Dept. of Plant Production - Lab. for Tropical Agronomy, Belgium*

²*Agricultural Research Centre for International Development (CIRAD), Performance of Tropical Production & Processing Systems (Persyst), France*

Oil palm plantations have a typical economic life cycle of 25 years. As a monocotyle-neous perennial crop the root system is radially distributed and shows no secondary growth but continuously expands. Oil palms are planted in an equilateral triangle design and the planting distance, determining the planting density, is fixed at the time of establishment. In time the roots from neighbouring palms will overlap and competition for soil resources can limit the growth and production of individual palms.

This paper presents the results from a root analysis study carried out on an oil palm density trial in Nigeria. The biomass and characteristics of the root system in an oil palm density trial have never been thoroughly studied so this would further the understanding of how planting distance affects underground processes in oil palm plantations for a particular environment.

The trial field was planted in 2005 following a Fisher block design with 6 treatments and 4 replicates. Plots consist of 72 palms of which the inner 20 are considered as useful (double row of border palms). The analysis was performed on 3 densities, D1, D3 and D6 corresponding to a planting distance of 9.5, 8.5 and 7.5 m and planting densities of 128, 160 and 205 palms ha⁻¹ respectively. A simplified voronoi method was used for the root analysis on one palm per plot. The voronoi volume was divided in 4 or 5 zones and 4 horizons giving 16 or 20 sampling volumes. For each of these volumes all the roots were removed and sorted in primary, secondary, fine and dead roots. Biomass for each was determined and extrapolated to a palm and plot level. Scanned images of the roots were also taken and analysed with WinRhizo to determine the root length, surface and volume.

Keywords: *Elaeis guineensis*, oil palm, planting density, root architecture, root system