

[← Previous](#)
[Next >](#)

 Details


 Figures


 Literature
Cited

DISEASE NOTES

 OPEN

First Report of Orange Rust Caused by *Puccinia kuehnii* on Sugarcane on the Island of Reunion

J. Hubert, C. Jeandel, L. Costet, B. Hostachy, A. S Dupuis, A. Coddeville, L. Barau, and R. loos

Affiliations ▾

Authors and Affiliations

J. Hubert¹C. Jeandel¹L. Costet²B. Hostachy³A. S Dupuis⁴A. Coddeville⁴L. Barau⁵R. loos¹ †

¹ANSES Laboratoire de la Santé des Végétaux, Unité de Mycologie, 54220 Malzéville, France

²CIRAD, UMR PVBMT, Saint-Pierre, La Réunion, France

³ANSES Laboratoire de la Santé des Végétaux, Unité des Ravageurs et Agents Pathogènes Tropicaux, Saint-Pierre, La Réunion, France

⁴DAAF 974, Service de l'Alimentation, La Réunion, France

⁵ERCANE, 97490 Saint-Denis, La Réunion, France

Published Online: 28 Aug 2019 |
<https://doi.org/10.1094/PDIS-04-19-0750-PDN>

Puccinia kuehnii (W. Krüger) E. J. Butler, the causal agent of orange rust, is reported as one of the two known widely spread rust species on commercial sugarcane (*Saccharum* spp.). This pathogen affects plant foliage, and the symptoms are characterized by the appearance of small elongated uredinial lesions (pustules), typically light orange to yellowish-brown, oval in shape, producing urediniospores. On highly susceptible cultivars, abundant pustules develop and coalesce on the lower leaf surface, causing tissue death. On such cultivars, orange rust has severe yield and economic


Just Published
Subscribe

ISSN: 0191-2917
 e-ISSN: 1943-7692

Metrics

Downloaded 128 times

Article History

Published: 28 Aug 2019

First Look: 25 Jun 2019

Accepted: 17 Jun 2019

Information

This article is in the public domain and copyrightable. It may be freely reprint customary crediting of the source The American Phytopathological Society, 2

Keywords

FUNGI

TROPICAL PLANTS

impacts (Magarey et al. 2001). This pathogen has expanded its range of distribution over the last 10 years and is now affecting America and Africa. In July and August 2018, in different areas of Reunion Island, typical rust symptoms were observed on leaves of R062006 and R062007 sugarcane varieties in selection. The intensity of the disease was low to moderate. Samples were collected on both varieties in fields from the ERCANE breeding station in Saint-Louis (Le Gol) in the southwest (-21.265027; 55.379830) and in Saint-Philippe (Le Baril) in the southeast (-21.359220; 55.731959) for laboratory analysis. The lesions were examined under a dissecting microscope. Microscopic slides were prepared from material by carefully picking up spores in rust sori with a needle. The presence of the causal fungus of orange rust was confirmed by the observation of typical microscopic features. Urediniospores were mostly obovoid to pyriform, echinulate, variable in size, 35.9 to 57.1 × 16.4 to 29.5 µm. Walls were 1 to 2.5 µm thick, most often with apical thickening up to 8.3 µm. A few inconspicuous hyaline paraphyses with a thin wall (1 to 2 µm) were present. No teliospore could be observed (Dixon et al. 2010; Virtudazo et al. 2001). Identity of the species involved was determined by sequencing the internal transcribed spacer (ITS) region of the rDNA followed by comparison with reference sequences available in GenBank. Leaf material containing uredinial lesions was cut into pieces about 2 × 3 mm and transferred into 2-ml microtubes. Total DNA was then extracted using a commercial plant DNA extraction kit (DNeasy Plant Mini Kit, Qiagen), and the fungal ITS region was amplified by PCR using specific primers PkPm-F/R, located respectively in ITS1 and ITS2 (Glynn et al. 2010). The nucleotide sequence was determined and deposited on GenBank (MK578656 to MK578657). Analysis of the sequence by BLAST showed 100 and 99% identity with *P. kuehnii*, over a 494- and 493-bp length, respectively, which was consistent with the morphological features observed. This is the first report of *P. kuehnii* in Reunion Island. Up to now, the disease has only been found in cultivar trials in Reunion. Orange rust has spread rapidly in the sister Mascarene Island of Mauritius since it was initially detected in March 2018. Out of four cultivars present in Mauritius and planted on Reunion, three tend to be moderately or slightly susceptible, and one is symptomless (S. Saumtally, personal communication). Following the emergence of brown rust caused by *Puccinia melanocephala* Syd. & P. Syd. in Reunion in 1965 (Boyer de la Giroday et al. 1979), resistant

cultivars, such as R570 bearing the *Brul* resistance gene (Costet et al. 2012), have been bred and planted to control this rust species. The same work will have to be conducted by the breeders in the years to come in order to take into account this new threat.

The author(s) declare no conflict of interest.



The American Phytopathological Society (APS)

📍 3340 Pilot Knob Road, St. Paul, MN 55121 USA

☎ +1.651.454.7250

FAX +1.651.454.0766

 APS