

BOOK OF ABSTRACTS  
TALKS





# Island BIOLOGY

La Réunion  
8-13 JULY

2019

📍 **Université de la Réunion**  
**Campus du Moufia**



**Island Biology**  
**BOOK OF ABSTRACTS**

TALKS

**Third International Conference  
on Island Ecology, Evolution and Conservation**

**8-13 July 2019  
University of La Réunion  
Saint Denis, France**

## Diversity and community composition of herbivorous thrips vary along environmental gradients, but plant effects remain an important driver

Niry T. Dianzinga <sup>\*† 1</sup>, Marie-Ludders Moutoussamy <sup>2</sup>, Joëlle Sadeyen <sup>2</sup>, Lala H. R. Ravaomanarivo <sup>3</sup>, Samuel Nibouche <sup>2</sup>, Enric Frago <sup>2,4</sup>

<sup>1</sup> Peuplements végétaux et bioagresseurs en milieu tropical (UMR PVBMT) – Faculté des Sciences et techniques - Université de La Réunion 15 avenue René Cassin CS92003 97744 Saint-Denis Cédex 9 pour le Cirad Station de Ligne-Paradis - 7 chemin de l'IRAT - 97410 Saint-Pierre, Réunion

<sup>2</sup> UMR Peuplement Végétaux et Bioagresseurs en Milieu Tropical (UMR PVBMT - INRA) – 97410 St-Pierre, Réunion

<sup>3</sup> Département d'Entomologie [Antananarivo] – Faculté des Sciences B.P 906 - 101 Antananarivo, Madagascar

<sup>4</sup> Centre de Biologie pour la Gestion des Populations (CBGP) – Centre de Coopération Internationale en Recherche Agronomique pour le Développement : UMR55 – 755 avenue du Campus Agropolis, 34988 Montpellier sur Lez, France

Elevational gradients are a powerful tool to explore how species communities respond to changes in environmental conditions along short geographic distances. It is usually observed that species diversity decrease monotonically with elevation, or follow a mid-elevation peak. These effects, however, may vary among taxa and geographic regions. Changes driven by elevation may also depend on landscape characteristics like habitat heterogeneity and patchiness, but also on seasonal variations such as those occurring between cold and hot seasons. Here, we used as biological model thrips (Insecta: Thysanoptera), which is a diverse group of slender insects, to investigate the drivers of insect diversity and of community composition along elevational gradients in Reunion Island. We sampled thrips on 106 plant species belonging to 44 families along seven replicated elevated gradients from 0 to 1,600 m in both summer and winter conditions. We identified 4,278 specimens representing 41 species and we estimated their diversity both at the local (alpha) and at the compositional (beta) diversity level. These metrics were related with elevation, and with landscape characteristics around each site, which included habitat diversity and patchiness (or fragmentation). We found that different thrips communities are directly related to host plants: invasive plants hosted a large diversity of thrips, whereas some endemic ones hosted unique communities. Elevation, patchiness and season were variables with an evident effect on both alpha and beta diversity. Beta diversity revealed that diversity of thrips was larger at low elevations, and this change was mostly dominated by richness differences (i.e. low elevation sites had more unique species than high elevation ones). Our results show that important diversity patterns can be appreciated through the lens of beta diversity, and may help to understand how regional pools of insects are distributed in relationship with topography, landscape and climatic conditions.

**Keywords:** tropical island, environmental gradients, beta diversity, alpha diversity, thrips

---

\*Speaker

†Corresponding author: [niry.dianzinga@gmail.com](mailto:niry.dianzinga@gmail.com)