

BOOK OF ABSTRACTS

TALKS




Island BIOLOGY

La Réunion
8-13 JULY

2019

📍 **Université de la Réunion**
Campus du Moufia



Island Biology
BOOK OF ABSTRACTS

TALKS

**Third International Conference
on Island Ecology, Evolution and Conservation**

**8-13 July 2019
University of La Réunion
Saint Denis, France**

Editors: Olivier Flores, Claudine Ah-Peng, Nicholas Wilding

Do endemic trees flora make endemic forests? Insights from New Caledonian forests

Philippe Birnbaum ^{*† 2,1}, Thomas Ibanez ³, Robin Pouteau ⁴, Hervé Vandrot ⁵, Vanessa Hequet ⁶, Grégoire Blanchard ⁵, Jeremy Girardi ⁵, Dimitri Justeau-Allaire ¹, Jean-Jérôme Cassan ⁷

² Institut Agronomique Néo-Calédonien (IAC) – Centre IRD de Nouméa, New Caledonia

¹ Botanique et Modélisation de l'Architecture des Plantes et des Végétations (UMR AMAP) – Centre de Coopération Internationale en Recherche Agronomique pour le Développement : UMR51-2015, Institut Agronomique Néo-Calédonien – Bd de la Lironde TA A-51/ PS 2 34398 Montpellier cedex 5, France

³ Department of Biology, University of Hawai – University of Hawai'i at Hilo, United States

⁴ Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and Conservation – Taizhou University, Taizhou, China

⁵ Institut Agronomique Néo-Calédonien (IAC) – Nouméa, Nouvelle-Calédonie, New Caledonia

⁶ Botanique et Modélisation de l'Architecture des Plantes et des Végétations (UMR AMAP-NC) – Centre IRD de Nouméa BP A5 - 98848 - Nouméa, New Caledonia

⁷ Service impact environnement et conservation (SIEC-DDEE - province Nord) – Kohné, Province Nord, New Caledonia

New Caledonia homes a rich and highly original flora with a species endemic rate > 75% and fascinating representation of relict taxa (gymnosperms and basal angiosperms). As a result, previous studies on the island flora have mostly focused on the taxonomy and biogeographical origins of this exceptional flora while few studies have attempted to understand the spatial distribution of species and the structure and diversity of species assemblages. Here, we present new insights into the diversity, structure, and ecology of trees communities derived from the New Caledonian Plant Inventory and Permanent Plot Network (NC-PIPPN). NC-PIPPN consists of standardized forest inventories scattered throughout the New Caledonian main island. This network groups together ca. 450 plots including more than 70 000 occurrences of woody plants (trees, shrubs, lianas, tree ferns, and palms) belonging to more than 950 mostly endemics species. Most species are distributed along wide environmental ranges (ca. 900 m of elevation and 2200 mm of mean annual rainfall) and contrasted substrates (volcano-sedimentary, ultramafics and calcareous). Wide environmental ranges, however, do not significantly correlate with large spatial distribution or high local abundance. As in other places in the tropics, the diversity of the forests is support by a highly uneven species abundance distribution. Less than 20% of tree species account for > 50% of all known occurrences, while half of the tree species contribute to < 16% of occurrences. Local abundance is also independant to spatial distribution: some rare species at the island scale are locally abundant while some frequent species at the island scale are locally rare. The spatial distribution of species results in highly heterogeneous forests (high beta diversity) that contrasts with a relatively homogeneous structure of communities. Despite a highly original flora and a pattern of aggregative species distribution, New Caledonian forests are not so distinguishable from other forests in the South Pacific region. Our results suggest that the New Caledonian rainforests are mostly constrained by geographical features (area and isolation of the archipelago) and climatic features (e.g. cyclonic frequency) while the flora uniqueness poorly contribute to the forest structure.

Keywords: New Caledonia, diversity, communities, trees

*Speaker

†Corresponding author: birnbaum@cirad.fr