

SHORT COMMUNICATION

The Windblown: Possible Explanations for Dinophyte DNA in Forest Soils

Marc Gottschling^a , Lucas Czech^{b,c} , Frédéric Mahé^{d,e} , Sina Adl^f  & Micah Dunthorn^{*g,h} 

a Department Biologie, Systematische Botanik und Mykologie, GeoBio-Center, Ludwig-Maximilians-Universität München, Munich, D-80638, Germany

b Computational Molecular Evolution Group, Heidelberg Institute for Theoretical Studies, Heidelberg, D-69118, Germany

c Department of Plant Biology, Carnegie Institution for Science, Stanford, California, 94305, USA

d CIRAD, UMR BGPI, Montpellier, F-34398, France

e BGPI, Université de Montpellier, CIRAD, IRD, Montpellier SupAgro, , Montpellier, France

f Department of Soil Sciences, College of Agriculture and Bioresources, University of Saskatchewan, Saskatoon, SK, S7N 5A8, Canada

g Eukaryotic Microbiology, Faculty of Biology, Universität Duisburg-Essen, Essen, D-45141, Germany

h Centre for Water and Environmental Research (ZWU), Universität Duisburg-Essen, Essen, D-45141, Germany

Keywords

Biodiversity; biogeography; dinoflagellates; environmental DNA; phylogeny.

*Correspondence

M. Dunthorn, Eukaryotic Microbiology, Faculty of Biology, Universität Duisburg-Essen, Universitätsstrasse 5, D-45141 Essen, Germany

Telephone number: +49-(0)-201-183-2453;
FAX number: +49 (0)201 183 3768
e-mail: micah.dunthorn@uni-due.de

ABSTRACT

Dinophytes are widely distributed in marine- and fresh-waters, but have yet to be conclusively documented in terrestrial environments. Here, we evaluated the presence of these protists from an environmental DNA metabarcoding dataset of Neotropical rainforest soils. Using a phylogenetic placement approach with a reference alignment and tree, we showed that the numerous sequencing reads that were phylogenetically placed as dinophytes did not correlate with taxonomic assignment, environmental preference, nutritional mode, or dormancy. All the dinophytes in the soils are rather windblown dispersal units of aquatic species and are not biologically active residents of terrestrial environments.

Received: 23 July 2020; revised 12 October 2020; accepted October 28, 2020.

doi:10.1111/jeu.12833

ENVIRONMENTAL high-throughput sequencing (HTS) studies of protists have now been performed for over a decade (Santoferrara et al. 2020). During that time, a large diversity of dinophyte DNA sequences has been uncovered. Dinophytes are an ecologically and economically important group of protists that exhibit many types of life styles and nutritional modes, including phototrophic, mixotrophic, and heterotrophic forms as well as some being parasitic (Saldarriaga and Taylor 2017). All known dinophytes are from marine or freshwater environments (Adl et al. 2019), except maybe the unarmored (athecate) *Esoptrodinium* (Fawcett and Parrow 2014). As dinophytes constitute a considerable fraction of the plankton and play an important role in the global aquatic ecosystem, HTS studies have detected their DNA from waters sampled from the polar regions through to the tropics (Annenkova

et al. 2020; Decelle et al. 2018; Elferink et al. 2017; Giner et al. 2020; Gottschling et al. 2020; Le Bescot et al. 2016; Lentendu et al. 2018; de Vargas et al. 2015). HTS studies have also detected DNA of dinophytes in terrestrial environments (Bates et al. 2013; Geisen et al. 2015; Mahé et al. 2017; Venter et al. 2017; Voss et al. 2019), although they are not expected to be there.

Aquatic protists can sometimes be detected in terrestrial environments, notably riparian soils, such as foraminifera (Lejzerowicz et al. 2010; Meisterfeld et al. 2001) and possibly haptophytes (Mahé et al. 2017). However, that does not mean that the normally aquatic protists are biologically active in soils or other drier environments (Geisen et al. 2018). In the absence of observing putative soil dinophytes using direct microscopic observations, here we used Mahé et al.'s (2017) metabarcoding data from three

lowland Neotropical rainforest soils to ask whether the presence of dinophytes in those soils correlates with taxonomic assignment, environmental preference, nutritional mode, or dormancy.

MATERIALS AND METHODS

Environmental sampling and data

Sampling and sequencing of tropical soils originally took place in lowland rainforest in Costa Rica, Panama, and Ecuador (Mahé et al. 2017). While the methodologies and bioinformatics can be found in detail in Mahé et al. (2017), here in brief: the extracted soil DNAs were amplified for the hypervariable V4 region of the SSU-rRNA locus using general eukaryotic primers (Stoeck et al. 2010); this short region has relatively strong phylogenetic signal, although it is not as strong as the full-length SSU-rRNA (Dunthorn et al. 2014; Gottschling et al. 2020). Illumina sequencing reads were clustered into OTUs using swarm v2 (Mahé et al. 2015) and taxonomically assigned to the Protist Ribosomal Reference database v203 (Guillou et al. 2013) using VSEARCH v1.6.0 (Rognes et al. 2016). The 269 OTUs that were assigned to the dinophytes by Mahé et al. (2017) were extracted and used here for phylogenetic placements (File S1).

Reference tree

From GenBank (Sayers et al. 2020), 228 ingroup dinophytes, plus 10 outgroups, were downloaded, then aligned with MAFFT v6.624b (Katoh and Standley 2013) using the —auto option (File S2). Based on previous analyses (Chacón and Gottschling 2020; Gottschling et al. 2012, 2020; Žerdoner Čalasan et al. 2019), the full sequences of each species were used without excluding ambiguously aligned positions sites. Phylogenetic inferences of the reference alignment were carried out by using Maximum Likelihood (ML), as described in detail by Gottschling et al. (2012, 2020), as implemented RAxML v8.2.10 (Stamatakis 2014) with the GTR + G substitution model. To determine the best-fitted ML tree, we executed 10-tree searches from distinct random stepwise addition sequence Maximum Parsimony starting trees and performed 1,000 non-parametric bootstrap replicates. Reference alignment and tree are available as Supporting Information (Files S3-S4).

Phylogenetic placement of environmental OTUs

The OTU representative sequences obtained from Swarm were aligned against the reference alignment using PaPaRa v2.0 (Berger and Stamatakis 2011) and phylogenetically placed onto the ML reference tree using the Evolutionary Placement Algorithm (Berger et al. 2011) as implemented in RAxML. Next, all OTUs that were placed with at least 95% probability (combined likelihood weight ratios) in the dinophyte clade were extracted and visualized, using Gappa v0.1.0 (Czech et al. 2020). For details on the extraction, see Czech et al. (2018); details of the

workflow are published in a GitHub code repository (<https://github.com/lczech/dinoflagellate-paper>).

RESULTS AND DISCUSSION

For the dinophyte reference alignment and tree, we included a broad representative taxon sample covering the known DNA sequence diversity with comprehensive sequence information, similar to previous strategies (Chacón and Gottschling 2020; Gottschling et al. 2012, 2020; Gu et al. 2013). The alignment was 7,270 bp long and had 3,753 parsimony informative sites (52%, 15.7 per terminal taxon). The ML tree had many bipartitions with high if not maximal bootstrap values. The Dinophyceae was inferred here to be monophyletic, and it contained the well-known subclades: Dinophysales, Gonyaulacales, Gymnodiniales, Peridiniales, Prorocentrales, †Suessiales as well as Amphidomataceae, Brachydiniaceae, and Tovelliaceae. Only 207 OTUs from the Mahé et al.'s (2017) that were assigned to the dinophytes, phylogenetically placed here across the reference tree with high likelihood weight scores (Fig. 1, File S5).

There was no exclusive correlation with taxonomic assignment. Some of the dinophyte OTUs formed distinct clades of sequences that were unknown until the present study. However, the proportion of such undescribed diversity is low compared to other microbial lineages such as the Fungi (Bass et al. 2018; Jones et al. 2011). They placed onto early branches, which comprise heterotrophic, mainly parasitic species (Bachvaroff et al. 2012; Gómez et al. 2009; Gu et al. 2013; Saldarriaga et al. 2003), but also within the Peridiniales. Most of the OTUs, though, placed within already known lineages of the Gymnodinaceae, Peridiniales, and †Suessiales, a truly heterogeneous set of dinophytes including unarmored and thecate algae as well. There is no morphological trait that the OTUs would therefore necessarily share and that would unite them with these different taxa. Some thecate groups such as Dinophysales, Gonyaulacales, Prorocentrales, and Protoperidiniae (Peridiniales) did not include any of the OTUs obtained from the environmental samples.

There was no exclusive correlation with habitat preference. Some of the dinophyte OTUs placed onto branches that contain freshwater species such as the Gymnodinaceae (Kretschmann et al. 2015; Romeikat et al. 2020), Tovelliaceae (Lindberg et al. 2005), and peridinialean *Naiadinium* comprising a freshwater lineage within otherwise marine *Scrippsiella* s.l. (Kretschmann et al. 2014; Luo et al. 2016). This placement of OTUs within freshwater clades was similar to what was shown for the haptophyte OTUs from the same rainforest soils (Mahé et al. 2017). However, none of the OTUs placed with the Peridinaceae, which is one of the most prominent freshwater dinophyte lineages (Gottschling et al. 2020; Moestrup and Calado 2018), but some OTUs placed on branches that contain just marine species, such as the Amphidomataceae (Tillmann et al. 2014) and Brachydiniaceae (Berg Holtz et al. 2006; Henrichs et al. 2011). Groups such as




Figure 1 A molecular reference tree recognizing major groups of dinophytes. Maximum likelihood (ML) tree of 228 representative dinophyte sequences (with strain number information), plus 10 outgroups, as inferred from a SSUrRNA nucleotide alignment (3,767 parsimony informative positions). Numbers on branches are ML bootstrap (above) and Bayesian support values (below) for the clusters (*maximal support values; values < 50 not shown).

Dinophysales, Gonyaulacales, Prorocentrales, and Protoperidiiniaceae (Peridiniales) are primarily marine and did not include any of the OTUs obtained from the environmental samples.

There was no exclusive correlation with nutritional mode. Some of the dinophyte OTUs placed onto branches that predominantly contain phototrophic species, such as Gymnodiniaceae, Peridiniales, and †Suessiales. Other OTUs placed with the heterotrophic species such as in Thoracosphaeraceae (i.e. *Pfiesteria* and relatives), but not in the consistently heterotrophic Dinophysales and Protoperidiiniaceae (Peridiniales). Transitions between phototrophic and heterotrophic modes are thought to occur in

the dinophytes (Fawcett and Parrow 2014; Jeong et al. 2012), but there is no phylogenetic signal for this trait at high taxonomic levels. Additionally, some OTUs placed onto early branches that include many parasitic species (Bachvaroff et al. 2012; Gómez et al. 2009; Saldarriaga et al. 2003), and they placed onto later branches that also include parasites, such as the Gymnodiniaceae (Gómez et al. 2009; Kretschmann et al. 2015; Romekat et al. 2020) and Peridiniales (Coats et al. 2010; Gottschling and Söhner 2013).

There was no exclusive correlation with dormancy practice. In addition to flagellated trophic cells, coccoid stages are integral part in the life-history of many dinophytes

from marine and freshwater environments. Coccoid cells are particularly abundant in the Gonyaulacales where no OTUs placed, or in the Peridiniales where many OTUs placed (Dale 1983; Evitt and Davidson 1964; Fensome et al. 1993; Wall 1971). The exact ecological functions of coccoid cells are not worked out rigorously for more than a handful of dinophyte species, but may frequently correspond to resting and/or dormancy stages (Fensome et al. 1993). Deposited in sediments, coccoid cells have the potential to preserve the local biodiversity like diaspores in a seed bank (Dale 1983). They can be considered either as old remnants from formerly aquatic, but now currently terrestrial environments (Boere et al. 2011; Sønstebo et al. 2010) or the result of random dispersal (Foissner 2006, 2011) by birds (Kretschmann et al. 2018; Tesson et al. 2018) or by wind. However, the difference between the taxonomic assignments of the rainforest soil samples (see above) may indicate that the ability to form coccoid cells during life history is not decisive for their terrestrial occurrence. An ecological group that has been receiving more interest in the past years and that may also be considered for the evaluation of the terrestrial samples is benthic dinophytes living in the intertidal (Hoppnerath et al. 2014); phylogenetically, it is a heterogeneous assemblage recruiting members particularly from Gonyaulacales and Peridiniales. But the bigger question remains, as to whether there are any soil dinophytes at all, or we are simply detecting windblown cells or lost dormant cells.

CONCLUSION

The presence of dinophyte DNA sequences in the Neotropical rainforest soils—as an exemplar of a terrestrial environment—did not correlate with taxonomic assignment, environmental preference, dormancy practice during life history, or nutrition mode/organismal interaction. The reason for their presence thus remains to be identified. We have to keep in mind that presence of DNA does not necessarily indicate biological activity. The environmental DNA sequences identified here are scattered unevenly across the classification, but it may represent more easily dispersed surface algae. To completely resolve this paradox, microscopy on soil samples with dinophyte DNA, especially with fluorescence *in situ* hybridization probes designed from the molecular OTUs, needs to be performed to verify their presence; however, dinophytes have never been conclusively observed in soils even after two centuries of observations. As suggested for other protists (Ehrenberg 1849), the most likely explanation for dinophyte DNA in soils is that they were passively dispersed there—they are the windblown.

ACKNOWLEDGMENTS

We thank Matthew Parrow and Gerard Versteegh for helpful edits, and Alexandros Stamatakis for computational support. Funding came from the Deutsche Forschungsgemeinschaft (grant DU1319/5-1) to MD.

LITERATURE CITED

- Adl, S. M., Bass, D., Lane, C. E., Lukeš, J., Schoch, C. L., Smirnov, A., Agatha, S., Berney, C., Brown, M. W., Burki, F., Cárdenas, P., Čepička, I., Chistyakova, L., Campo, J., Dunthorn, M., Edvardsen, B., Eglit, Y., Guillou, L., Hampl, V., Heiss, A. A., Hoppnerath, M., James, T. Y., Karnkowska, A., Karpov, S., Kim, E., Kolisko, M., Kudryavtsev, A., Lahr, D. J. G., Lara, E., Le Gall, L., Lynn, D. H., Mann, D. G., Massana, R., Mitchell, E. A. D., Morrow, C., Park, J. S., Pawłowski, J. W., Powell, M. J., Richter, D. J., Rueckert, S., Shadwick, L., Shimano, S., Spiegel, F. W., Torruella, G., Youssef, N., Zlatogursky, V. & Zhang, Q. 2019. Revisions to the classification, nomenclature, and diversity of eukaryotes. *J. Eukaryot. Microbiol.*, 66:4–119.
- Annenkova, N. V., Giner, C. R. & Logares, R. 2020. Tracing the origin of planktonic protists in an ancient lake. *Microorganisms*, 8:543.
- Bachvaroff, T. R., Kim, S., Guillou, L., Delwiche, C. F. & Coats, D. W. 2012. Molecular diversity of the syndinean genus *Euduboscquella* based on single-cell PCR analysis. *Appl. Environ. Microbiol.*, 78:334–345.
- Bass, D., Czech, L., Williams, B. A. P., Berney, C., Dunthorn, M., Mahé, F., Torruella, G., Stentiford, G. D. & Williams, T. A. 2018. Clarifying the relationships between Microsporidia and Cryptomycota. *J. Eukaryot. Microbiol.*, 65:773–782.
- Bates, S. T., Clemente, J. C., Flores, G. E., Walters, W. A., Parfrey, L. W., Knight, R. & Fierer, N. 2013. Global biogeography of highly diverse protistan communities in soil. *ISME J.*, 7:652–659.
- Berger, S. A., Krompass, D. & Stamatakis, A. 2011. Performance, accuracy, and web server for evolutionary placement of short sequence reads under maximum likelihood. *Syst. Biol.*, 60:291–302.
- Berger, S. A. & Stamatakis, A. 2011. Aligning short reads to reference alignments and trees. *Bioinformatics*, 27:2068–2075.
- Berg Holtz, T., Daugbjerg, N., Moestrup, Ø. & Fernández-Tejedor, M. 2006. On the identity of *Karlodinium veneficum* and description of *Karlodinium armiger* sp. nov. (Dinophyceae), based on light and electron microscopy, nuclear-encoded LSU rDNA, and pigment composition. *J. Phycol.*, 42:170–193.
- Boere, A. C., Rijpstra, W. I. C., de Lange, G. J., Malinverno, E., Sinninghe Damsté, J. S. & Coolen, M. J. L. 2011. Exploring preserved fossil dinoflagellate and haptophyte DNA signatures to infer ecological and environmental changes during deposition of sapropel S1 in the eastern Mediterranean. *Paleoceanography*, 26:PA2204.
- Chacón, J. & Gottschling, M. 2020. Dawn of the dinophytes: a first attempt to date origin and diversification of harmful algae. *Harmful Algae*, 97:101871. <https://doi.org/10.1016/j.hal.2020.101871>
- Coats, D. W., Kim, S., Bachvaroff, T. R., Handy, S. M. & Delwiche, C. F. 2010. *Tintinnophagus acutus* n. g., n. sp. (Phylum Dinoflagellata), an ectoparasite of the ciliate *Tintinnopsis cylindrica* Daday 1887, and its relationship to *Duboscquelinium colini* Grassé 1952. *J. Eukaryot. Microbiol.*, 57:468–482.
- Czech, L., Barbera, P. & Stamatakis, A. 2018. Methods for automatic reference trees and multilevel phylogenetic placement. *Bioinformatics*, 35:1151–1158.
- Czech, L., Barbera, P. & Stamatakis, A. 2020. Genesis and Gappa: processing, analyzing and visualizing phylogenetic (placement) data. *Bioinformatics*, 36:3263–3265.
- Dale, B. 1983. Dinoflagellate resting cysts: “Benthic plankton”. In: Fryxell, G. A. (ed.), *Survival Strategies of the Algae*. Cambridge University Press, Cambridge. p. 69–136.

- de Vargas, C., Audic, S., Henry, N., Decelle, J., Mahé, F., Logares, R., Lara, E., Berney, C., Le Bescot, N., Probert, I., Carmichael, M., Poulain, J., Romac, S., Colin, S., Aury, J.-M., Bittner, L., Chaffron, S., Dunthorn, M., Engelen, S., Flegontova, O., Guidi, L., Horák, A., Jaillon, O., Lima-Mendez, G., Lukeš, J., Malviya, S., Morard, R., Mulot, M., Scalco, E., Sian, O. R., Vincent, F., Zingone, A., Dimier, C., Picheral, M., Searson, S., Kandels-Lewis, S., Tara Oceans Coordinators, Acinas, S. G., Bork, P., Bowler, C., Gorsky, G., Grimsley, N., Hingamp, P., Iudicone, D., Not, F., Ogata, H., Pesant, S., Raes, J., Sieracki, M. E., Speich, S., Stemmann, L., Sunagawa, S., Weissenbach, J., Wincker, P. & Karsenti, E. 2015. Eukaryotic plankton diversity in the sunlit ocean. *Science*, 348:1261605.
- Decelle, J., Carradec, Q., Pochon, X., Henry, N., Romac, S., Mahé, F., Dunthorn, M., Kourlaiev, A., Voolstra, C. R., Wincker, P. & de Vargas, C. 2018. Worldwide occurrence and activity of the reef-building coral symbiont *Symbiodinium* in the open ocean. *Curr. Biol.*, 28:3625–3633.
- Dunthorn, M., Otto, J., Berger, S. A., Stamatakis, A., Mahé, F., Romac, S., de Vargas, C., Audic, S., BioMarKs Consortium, Stock, A., Kauff, F. & Stoeck, T. 2014. Placing environmental next-generation sequencing amplicons from microbial eukaryotes into a phylogenetic context. *Mol. Biol. Evol.*, 31:993–1009.
- Ehrenberg, C. G. 1849. Passat-Staub und Blut-Regen: ein grosses organisches unsichtbares Wirken und Leben in der Athmosphäre; mehrere Vorträge. Königlichen Akademie der Wissenschaften, Berlin.
- Elferink, S., Neuhaus, S., Wohlrab, S., Töbe, K., Voß, D., Gottschling, M., Lundholm, N., Krock, B., Koch, B. P., Zielinski, O., Cembella, A. & John, U. 2017. Molecular diversity patterns among various phytoplankton size-fractions in West Greenland in late summer. *Deep Sea Res. Part I*, 121:54–69.
- Evitt, W. R. & Davidson, S. E. 1964. Dinoflagellate studies. I. Dinoflagellate cysts and thecae. In: Harbaugh, J. W. (ed.), Stanford University Publications in the Geological Sciences. Stanford University Press, Stanford, CA. p. 1–12.
- Fawcett, R. C. & Parrow, M. W. 2014. Mixotrophy and loss of phototrophy among geographic isolates of freshwater *Esiodinium/Bernardinum* sp. (Dinophyceae). *J. Phycol.*, 50:55–70.
- Fensome, R. A., Taylor, F. J. R., Norris, G., Sarjeant, W. A. S., Wharton, D. I. & Williams, G. L. 1993. A classification of living and fossil dinoflagellates. *Micropaleontology*. Special Publication Number, 7:1–245.
- Foissner, W. 2006. Biogeography and dispersal of micro-organisms: a review emphasizing protists. *Acta Protozool.*, 45:111–136.
- Foissner, W. 2011. Dispersal of protists: the role of cysts and human introductions. In: Fontaneto, D. (ed.), Biogeography of microscopic organisms: is everything small everywhere? Cambridge University Press, Cambridge. p. 61–87.
- Geisen, S., Mitchell, E. A. D., Adl, S., Bonkowski, M., Dunthorn, M., Ekelund, F., Fernández, L. D., Jousset, A., Krashevskaya, V., Singer, D., Spiegel, F. W., Walochnik, J. & Lara, E. 2018. Soil protists: a fertile frontier in soil biology research. *FEMS Microbiol. Rev.*, 42:293–323.
- Geisen, S., Tveit, A. T., Clark, I. M., Richter, A., Svenning, M. M., Bonkowski, M. & Urich, T. 2015. Metatranscriptomic census of active protists in soils. *ISME J.*, 9:2178–2190.
- Giner, C. R., Pernice, M. C., Balagué, V., Duarte, C. M., Gasol, J. M., Logares, R. & Massana, R. 2020. Marked changes in diversity and relative activity of picoeukaryotes with depth in the world ocean. *ISME J.*, 14:437–449.
- Gómez, F., López-García, P., Nowaczyk, A. & Moreira, D. 2009. The crustacean parasites *Ellobiopsis Caulleryi*, 1910 and *Thalassomyces Niezabitowski*, 1913 form a monophyletic divergent clade within the Alveolata. *Syst. Parasitol.*, 74:65–74.
- Gottschling, M., Chacón, J., Žerdoner Čalasan, A., Neuhaus, S., Kretschmann, J., Stibor, H. & John, U. 2020. Phylogenetic placement of environmental sequences using taxonomically reliable databases helps to rigorously assess dinophyte biodiversity in Bavarian lakes (Germany). *Freshw. Biol.*, 65:193–208.
- Gottschling, M., Söhner, S., Zinßmeister, C., John, U., Plötner, J., Schweikert, M., Aligizaki, K. & Elbrächter, M. 2012. Delimitation of the Thoracosphaeraceae (Dinophyceae), including the calcareous dinoflagellates, based on large amounts of ribosomal RNA sequence data. *Protist*, 163:15–24.
- Gottschling, M. & Söhner, S. 2013. An updated list of generic names in the Thoracosphaeraceae. *Microorganisms*, 1:122–136.
- Gu, H., Kirsch, M., Zinßmeister, C., Söhner, S., Meier, K. J. S., Liu, T. & Gottschling, M. 2013. Waking the dead: morphological and molecular characterization of extant *Posoniella tricarinelloides* (Thoracosphaeraceae, Dinophyceae). *Protist*, 164:583–597.
- Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., de Vargas, C., Decelle, J., Del Campo, J., Dolan, J. R., Dunthorn, M., Edvardsen, B., Holzmann, M., Kooistra, W. H., Lara, E., Le Bescot, N., Logares, R., Mahé, F., Massana, R., Montresor, M., Morard, R., Not, F., Pawłowski, J., Probert, I., Sauvadet, A. L., Siano, R., Stoeck, T., Vault, D., Zimmermann, P. & Christen, R. 2013. The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote small sub-unit rRNA sequences with curated taxonomy. *Nucleic Acids Res.*, 41:D597–D604.
- Henrichs, D. W., Sosik, H. M., Olson, R. J. & Campbell, L. 2011. Phylogenetic analysis of *Brachidinium capitatum* (Dinophyceae) from the Gulf of Mexico indicates membership in the Kareniaeae. *J. Phycol.*, 47:366–374.
- Hoppenrath, M., Murray, S. A. & Chomérat, N. 2014. Marine benthic dinoflagellates – Unveiling their worldwide biodiversity. Schweizerbart Science Publishers, Stuttgart.
- Jeong, H. J., Yoo, Y. D., Kang, N. S., Lim, A. S., Seong, K. A., Lee, S. Y., Lee, M. J., Lee, K. H., Kim, H. S., Shin, W., Nam, S. W., Yih, W. & Lee, K. 2012. Heterotrophic feeding as a newly identified survival strategy of the dinoflagellate *Symbiodinium*. *Proc. Natl. Acad. Sci. USA*, 109:12604–12609.
- Jones, M. D. M., Forn, I., Gadelha, C., Egan, M. J., Bass, D., Massana, R. & Richards, T. A. 2011. Discovery of novel intermediate forms redefines the fungal tree of life. *Nature*, 474:200–203.
- Katoh, K. & Standley, D. M. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.*, 30:772–780.
- Kretschmann, J., Filipowicz, N. H., Owsiany, P. M., Zinßmeister, C. & Gottschling, M. 2015. Taxonomic clarification of the unusual dinophyte *Gymnodinium limneticum* Wołosz. (Gymnodinaceae) from the Tatra Mountains. *Protist*, 166:621–637.
- Kretschmann, J., Owsiany, P. M., Žerdoner Čalasan, A. & Gottschling, M. 2018. The hot spot in a cold environment: Puzzling *Parvodinium* (Peridiniopsidaceae, Peridiniales) from the Polish Tatra Mountains. *Protist*, 230:206–230.
- Kretschmann, J., Zinßmeister, C. & Gottschling, M. 2014. Taxonomic clarification of the dinophyte *Rhabdosphaera erinaceus* Kamptner, = *Scirpsiella erinaceus* comb. nov. (Thoracosphaeraceae, Peridiniales). *Syst. Biodivers.*, 12:393–404.
- Le Bescot, N., Mahé, F., Audic, S., Dimier, C., Garet, M.-J., Poulain, J., Wincker, P., de Vargas, C. & Siano, R. 2016. Global patterns of pelagic dinoflagellate diversity across protist size classes unveiled by metabarcoding. *Environ. Microbiol.*, 18:609–626.

- Lejzerowicz, F., Pawłowski, J., Fraissinet-Tachet, L. & Marmeisse, R. 2010. Molecular evidence for widespread occurrence of Foraminifera in soils. *Environ. Microbiol.*, 12:2518–2526.
- Lentendu, G., Buosi, P. R. B., Cabral, A. F., Segovia, B. T., de Meira, B. R., Lansac-Toha, F. M., Velho, L. F. M., Ritter, C. & Dunthorn, M. 2018. Planktonic protist biodiversity and biogeography in lakes from four Brazilian river-floodplain systems. *J. Eukaryot. Microbiol.*, 66:592–599.
- Lindberg, K., Moestrup, Ø. & Daugbjerg, N. 2005. Studies on woloszynskiod dinoflagellates I: *Woloszynska coronata* re-examined using light and electron microscopy and partial LSU rDNA sequences, with description of *Tovellia* gen. nov. and *Jadwigia* gen. nov. (Tovelliaceae fam. nov.). *Phycologia*, 44:416–440.
- Luo, Z., Mertens, K. N., Bagheri, S., Aydin, H., Takano, Y., Matsuoka, K., McCarthy, F. M. G. & Gu, H. 2016. Cyst-theca relationship and phylogenetic positions of *Scrippsiella plana* sp. nov. and *S. spinifera* (Peridiniales, Dinophyceae). *Eur. J. Protistol.*, 51:188–202.
- Mahé, F., de Vargas, C., Bas, S. D., Czech, L., Stamatakis, A., Lara, E., Singer, D., Mayor, J., Bunge, J., Sernaker, S., Siemensmeyer, T., Trautmann, I., Romac, S., Berney, C., Kozlov, A., Mitchell, E. A. D., Seppey, C. V. W., Egge, E., Lentendu, G., Wirt, H. R., Trueba, G. & Dunthorn, M. 2017. Parasites dominate hyperdiverse soil protist communities in Neotropical rainforests. *Nat. Ecol. Evol.*, 1:0091.
- Mahé, F., Rognes, T., Quince, C., de Vargas, C. & Dunthorn, M. 2015. Swarm v2: highly-scalable and high-resolution amplicon clustering. *PeerJ*, 3:e1761.
- Meisterfeld, R., Holzmann, M. & Pawłowski, J. 2001. Morphological and molecular characterization of a new terrestrial allogromiid species: *Edaphoallogromia australica* gen. et spec. nov. (Foraminifera) from Northern Queensland (Australia). *Protist*, 152:185–192.
- Moestrup, Ø. & Calado, A. J. 2018. *Freshwater flora of Central Europe*, Vol. 6: Dinophyceae. Springer Spektrum, Berlin.
- Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. 2016. VSEARCH: a versatile open source tool for metagenomics. *PeerJ*, 4:e2584.
- Romeikat, C., Knechtel, J. & Gottschling, M. 2020. Clarifying the taxonomy of *Gymnodinium fuscum* var. *rubrum* from Bavaria (Germany) and placing it in a molecular phylogeny of the Gymnodiniaceae (Dinophyceae). *Syst. Biodivers.*, 18:102–115.
- Saldarriaga, J. F., McEwan, M. L., Fast, N. M., Taylor, F. J. R. & Keeling, P. J. 2003. Multiple protein phylogenies show that *Oxyrrhis marina* and *Perkinsus marinus* are early branches of the dinoflagellate lineage. *Int. J. Syst. Evol. Microbiol.*, 53:355–365.
- Saldarriaga, J. F. & Taylor, F. J. R. 2017. Dinoflagellata. In: Handbook of the Protists, 2nd edn. Springer International Publishing AG, Cham. p. 625–678.
- Santoferrara, L., Burki, F., Filker, S., Logares, R., Dunthorn, M. & McManus, G. B. 2020. Perspectives from ten years of protist studies by high-throughput metabarcoding. *J. Eukaryot. Microbiol.*, 67:612–622.
- Sayers, E. W., Cavanaugh, M., Clark, K., Ostell, J., Pruitt, K. D. & Karsch-Mizrachi, I. 2020. GenBank. *Nucleic Acids Res.*, 48:D84–D86.
- Sønstebo, J. H., Gielly, L., Brysting, A. K., Elven, R., Edwards, M., Haile, J., Willerslev, E., Coissac, E., Rioux, D., Sannier, J., Taberlet, P. & Brochmann, C. 2010. Using next-generation sequencing for molecular reconstruction of past Arctic vegetation and climate. *Mol. Ecol. Resour.*, 10:1009–1018.
- Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 30:1312–1313.
- Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M. D. M., Breiner, H. W. & Richards, T. A. 2010. Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water. *Mol. Ecol.*, 19:21–31.
- Tesson S. V. M., Weißbach A., Kremp A., Lindström Å., Rengefors K. (2018) The potential for dispersal of microalgal resting cysts by migratory birds. *Journal of Phycology*, 54 (4), 518–528. <http://dx.doi.org/10.1111/jpy.12756>
- Tillmann, U., Elbrächter, M., Gottschling, M., Gu, H., Jeong, H. J., Krock, B., Nézan, E., Potvin, E., Salas, R. & Söhner, S. 2014. The dinophycean genus *Azadinium* and related species – morphological and molecular characterization, biogeography, and toxins. In: Kim, H. G., Reguera, B., Hallegraeff, G. M., Lee, C. K., Han, H. S. & Choi, J. K. (ed.), Harmful Algae 2012 (Proceedings of the 15th International Conference on Harmful Algae). International Society for the Study of Harmful Algae. p. 149–152.
- Venter, P. C., Nitsche, F., Domonell, A., Heger, P. & Arndt, H. 2017. The protistan microbiome of grassland soil: diversity in the mesoscale. *Protist*, 168:546–564.
- Voss, C., Fiore-Donno, A. M., Guerreiro, M. A., Peršoh, D. & Bonkowski, M. 2019. Metatranscriptomics reveals unsuspected protistan diversity in leaf litter across temperate beech forests, with Amoebozoa the dominating lineage. *FEMS Microbiol. Ecol.*, 95:fiz142.
- Wall, D. 1971. Biological problems concerning fossilizable dinoflagellates. *Geosci. Mar.*, 3:1–15.
- Žerdoner Čalasan, A., Kretschmann, J. & Gottschling, M. 2019. They are young, and they are many: dating freshwater lineages in unicellular dinophytes. *Environ. Microbiol.*, 21:4125–4135.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Table S1. Voucher list.

File S1. The 269 OTUs that were assigned to the dinophytes by Mahé et al. (2017) that were extracted and used here for phylogenetic placements.

File S2. Taxon sampling for the dinophyte reference alignment.

File S3. Reference alignment for the dinophytes.

File S4. Reference tree for the dinophytes.

File S5. The 207 OTUs that were assigned to the dinophytes, which phylogenetically placed here across the reference tree with high likelihood weight scores.