



Nutrient Deficiency Tolerance in Citrus Is Dependent on Genotype or Ploidy Level

Julie Oustric¹, Raphaël Morillon², François Luro³, Stéphane Herbette⁴, Paul Martin⁵, Jean Giannettini¹, Liliane Berti¹ and Jérémie Santini^{1*}

¹ CNRS, Laboratoire Biochimie and Biologie Moléculaire du Végétal, UMR 6134 SPE, Université de Corse, Corsica, France, ² Equipe "Amélioration des Plantes à Multiplication Végétative", UMR AGAP, Département BIOS, CIRAD, Petit-Bourg, Guadeloupe, ³ UMR AGAP Corse, Station INRA/CIRAD, San-Giuliano, France, ⁴ UCA, INRA, PIAF, Clermont-Ferrand, France, ⁵ AREFLEC, San-Giuliano, France

OPEN ACCESS

Edited by:

Paula Casati,
Consejo Nacional de Investigaciones
Científicas y Técnicas (CONICET),
Argentina

Reviewed by:

Jose M. Colmenero-Flores,
Spanish National Research Council
(CSIC), Spain

Karina Eva Josefina Tripodi,
CONICET Rosario, Argentina

*Correspondence:

Jérémie Santini
jeremie.santini@univ-corse.fr

Specialty section:

This article was submitted to
Plant Abiotic Stress,
a section of the journal
Frontiers in Plant Science

Received: 13 September 2018

Accepted: 24 January 2019

Published: 11 February 2019

Citation:

Oustric J, Morillon R, Luro F,
Herbette S, Martin P, Giannettini J,
Berti L and Santini J (2019) Nutrient
Deficiency Tolerance in Citrus Is
Dependent on Genotype or Ploidy
Level. *Front. Plant Sci.* 10:127.
doi: 10.3389/fpls.2019.00127

Plants require essential minerals for their growth and development that are mainly acquired from soil by their roots. Nutrient deficiency is an environmental stress that can seriously affect fruit production and quality. In citrus crops, rootstock/scion combinations are frequently employed to enhance tolerance to various abiotic stresses. These tolerances can be improved in doubled diploid genotypes. The aim of this work was to compare the impact of nutrient deficiency on the physiological and biochemical response of diploid (2x) and doubled diploid (4x) citrus seedlings: Volkamer lemon, Trifoliolate orange × Cleopatra mandarin hybrid, Carrizo citrange, Citrumelo 4475, Fihorag1 (*Poncirus trifoliata* + and willow leaf mandarin), an allotetraploid somatic hybrid, was also included in this study. Our results showed that depending on the genotype, macronutrient and micronutrient deficiency affected certain physiological traits and oxidative metabolism differently. Tetraploid genotypes, mainly Fihorag1 and Citrumelo 4475, appeared resistant compared to the other genotypes as indicated by the lesser decrease in photosynthetic parameters (P_{net} , F_v/F_m , and G_s) and the lower accumulation of oxidative markers (MDA and H_2O_2) in roots and leaves, especially after long-term nutrient deficiency. Their higher tolerance to nutrient deficiency could be explained by better activation of their antioxidant system. For the other genotypes, tetraploidization did not induce greater tolerance to nutrient deficiency.

Keywords: antioxidant, citrus, nutrient deficiency, oxidative stress, photosynthesis, polyploid

INTRODUCTION

Polyploidization is considered as a major force in plant evolution (Wendel, 2000; Chen, 2007; Soltis and Soltis, 2009). Polyploid organisms possess more than two sets of chromosomes. There are two different types of polyploidy: autopolyploids, which result from intraspecific genome duplication occurring during incomplete mitosis, or allopolyploids, which are formed from the

Abbreviations: 2x, diploid; 4x, tetraploid; APX, ascorbate peroxidase; Asa, reduced ascorbate; Asa/DHA, redox status of ascorbate; CC2x, diploid Carrizo citrange; CC4x, tetraploid Carrizo citrange; CAT, catalase; CM2x, diploid Citrumelo 4475; CM4x, tetraploid Citrumelo 4475; DHA, oxidized ascorbate; F_0 , minimal chlorophyll *a* fluorescence; F_m , maximal fluorescence; F_v , variable fluorescence, F_v/F_m , maximum quantum efficiency of PSII; G_s , stomatal conductance; MDA, malondialdehyde; PMC2x, diploid hybrid Trifoliolate orange × Cleopatra mandarin; PMC4x, tetraploid hybrid Trifoliolate orange × Cleopatra mandarin; P_{net} , leaf net photosynthetic rate; ROS, reactive oxygen species; SOD, superoxide dismutase; tAsa, total ascorbate; VK2x, diploid Volkamer lemon; VK4x, tetraploid Volkamer lemon.

combination of at least two complete chromosome sets via interspecific hybridization (Stebbins, 1947; Doyle and Egan, 2010). Polyploidy confers several advantages: genome buffering (mutation and recombination), allele-dosage effects, increased level of allelic diversity and heterozygosity, and sub- and/or neo-functionalization of duplicated genes that can induce phenotypic variation (Comai, 2005; Udall and Wendel, 2006; Beest et al., 2012). Polyploids are usually resistant or more tolerant compared to their progenitors to adverse environmental conditions such as extreme temperatures, high radiation levels and nutrient-poor soil (Ehrendorfer, 1980; Levin, 1983; Brochmann et al., 2004; Hao et al., 2013). Citrus are generally diploid, with $2n = 2x = 18$ chromosomes (Krug, 1943); however tetraploids resulting from incomplete mitosis of somatic embryos are frequently found in seedlings of diploid apomictic genotypes (Cameron and Frost, 1968). Compared with the corresponding diploids, citrus 4x plants tend to have thicker and greener leaves, shorter thicker roots, larger fruit and depressed growth versus diploids (Ollitrault and Jacquemond, 1994; Guerra et al., 2014; Ruiz et al., 2016a,b,c). Anatomical and physiological changes between 4x and 2x counterparts (Syvertsen et al., 2000) could affect leaf mineral content.

Abiotic factors can cause oxidative stress due to an overproduction of ROS. An excess of ROS lead to protein denaturation, DNA mutation, and membrane lipid peroxidation (Mittler, 2002; Apel and Hirt, 2004; Mittler et al., 2004). It can be a tricky task to unravel the causes and the consequences in the oxidative metabolism responses. It implies to investigate both oxidative products levels like H_2O_2 as a primary marker of ROS and MDA as lipid peroxidation marker but also antioxidant responses with key antioxidant enzyme activities (mainly SOD, APX, CAT, and DHAR) and key antioxidant molecules such as ascorbate and proline. Antioxidant system has been found to be of paramount importance in the response and tolerance of trees to environmental stress (Polle et al., 1993; Mutlu et al., 2013; Santini et al., 2013; Xu et al., 2013).

In the citrus industry, one of the environmentally friendly techniques to prevent a drop in yield and fruit quality in abiotic stress conditions is grafting the scion onto a rootstock that confers better tolerance to those stresses. Rootstock selection is mainly based on the performance of the root system for coping with biotic (soil pathogens) and abiotic stress (salinity, drought, nutrient deficiency, and extreme temperature). Studies have demonstrated that using 4x seedlings or rootstocks enhances stress tolerance or allows to confer resistance when compared to their diploid counterparts (Saleh et al., 2008; Allario et al., 2013; Tan et al., 2015; Oustric et al., 2017).

When subjected to environmental stress, tolerant plants can offset the effects of oxidative damage while a resistant plant can prevent the rate of oxidative damage. These two strategies allow a plant protection from the harmful effects of stress (Mauricio et al., 1997).

Nutrient deficiency, particularly of a major nutrient (N, P, and K), is one of the most limiting factors along with drought, salinity, soil alkalinity, and extreme temperatures, on the performance of plants in their natural habitats and agricultural environments (Tewari et al., 2004). Plants require an optimal level of nutrients

for growth and normal function. An excess has a negative effect on soil biology while scarcity has a negative impact on growth and development. In addition, nutrient deficiency can disturb the plant's antioxidant system, as nutrients are needed for antioxidant biosynthesis (Kandlbinder et al., 2004). Plants possess a long distance mobile signal that allows them to respond to starvation and satiety signals throughout the plant through their vascular system. For example, the main symptoms of nutrient deficiency are stunted growth, chlorosis, necrosis and a disproportionately high allocation of biomass to the root system (Ericsson, 1995). Many genes play a central role in the acquisition and distribution of nutrients, including many protein-coding genes as well as microRNAs (miR395, miR399, miR398, miR397, and miR408) (Chiou, 2007; Sunkar et al., 2007).

In the citrus industry, nutrient deficiencies have an economic and ecologic importance. Thus, developing new citrus rootstocks that require less fertilizer is an urgent need, as well as improving our understanding of citrus responses to nutrient deficiency. The use of tetraploid rootstock could respond to this problem. However, to our knowledge, there is no information in the literature on the effect of genotype and ploidy level on resistance to nutrient deficiency. The aim of this study was to identify resistant or tolerant seedling genotypes among the most commonly used citrus rootstocks: VK, PMC, CC, CM. Moreover, in order to confirm the positive effect of polyploidization of the genotype on tolerance to nutrient deficiency, the 4x (doubled diploid) counterparts of these citrus seedlings were also studied. The allopolyploid Flhorag1 (FL)—a somatic hybrid of Trifoliate orange and Willow leaf mandarin—was also included in the experimental design (Ollitrault et al., 2000). The effects of nutrient deprivation and the plant's tolerance were evaluated at the root and leaf levels by analyzing relevant anatomical, physiological, and biochemical parameters.

MATERIALS AND METHODS

Plant Material and Growth Conditions

The experiment was performed on four citrus diploid (2x) seedling genotypes and their four doubled diploid (4x) counterparts (Table 1). The Flhorag1, an allotetraploid form, was also included. Plants were selected among seedlings made with seeds from trees maintained in the citrus germplasm collection (BCR NF 96-S-900 Citrus INRA/CIRAD) at San-Giuliano, Corsica (France). The ploidy status of 2x and 4x seedlings was checked by 10-color flow cytometry (Partec I, Germany) as described by Froelicher et al. (2007). Clonal propagation by nucellar embryogenesis was verified by genotyping using SSR markers as described by Vieira et al. (2016). The 12 seedlings of each genotype, giving a total of 108 plants were grown under identical conditions in vermiculite with ferti-irrigation and water (1 L/h) during 4 years in a tunnel greenhouse at the AREFLEC experimental station in San-Giuliano, Corsica (41°47'27" N and 09°23'40" E). The ferti-irrigation solutions were prepared and applied with a metering pump. The stock solution used for irrigation included 20-5-10 N (nitrogen) P

TABLE 1 | Citrus varieties used for physiological and biochemical analysis.

Species	Varieties	Abbreviations	Ploidy level	ICVN
<i>Citrus limonia</i> Osb.	Volkamer	VK	2x	110024
	lemon		4x	101122
<i>Poncirus trifoliata</i> L. Raf. × <i>Citrus reshni</i> Hort. ex Tan.	Trifoliata orange	PMC	2x	110155
	x Cleopatra mandarin		4x	101114
<i>Poncirus trifoliata</i> L. Raf. + <i>Citrus</i> <i>deliciosa</i> Ten.	Flhorag1	FL	4x	100951
<i>Citrus sinensis</i> L. Osb. × <i>Poncirus</i> <i>trifoliata</i> L. Raf.	Carrizo citrange	CC	2x	110181
			4x	101075
<i>Citrus paradisi</i> L. Macf. × <i>Poncirus</i> <i>trifoliata</i> L. Raf.	Citrumelo 4475	CM	2x	110410
			4x	101112

ICVN, International Citrus Variety Numbering.

(phosphorus) K (potassium) fertilizer + 2MgO (Magnesium oxide) + trace elements. This level of fertilization was based on the recommendations of the local department of agriculture. After 4 years and before starting the experiment, the vermiculite was washed with distilled water for 48 h in order to eliminate any nutritional reserves in the pot. This avoided the latency phase that could have occurred if the fertilizer remaining in the vermiculite had been completely consumed by the plants. The 4-years-old trees were then divided into two blocks: one with reference ferti-irrigation (control plants) and the other with irrigation water (without nutrient inputs). In each block, a total of three trees of each genotype was randomized, giving a total of 12 trees per block.

This experiment was carried out from May 2016 to January 2017 on homogeneous plants comprising four branches with leaves developing under stress conditions and having reached full maturity. Nutrient deficiency was applied for 210 days and then trees were returned to the baseline condition with the control fertigation solution.

The samplings and physiological measurements were carried out at different times (days) selected based on preliminary experiments: 0 (control plant), 70 (D70), 140 (D140), and 210 (D210) days after the start of nutritional deprivation and after 30 days of recovery (30DR). For physiological measurements, nine fully expanded leaves were analyzed, i.e., three per tree (nine replicates), between 7 AM and 11 AM. For biochemical analyses, three samples were collected for each genotype, i.e., one per tree, and each sample is obtained by pooling eight fully-expanded leaves.

Root samplings for the biochemical measurements were performed at four different times: 0 (control plant), D70, D210, and 30DR. The trees were removed from their pots and cleared of residual vermiculite, and then rinsed with water. For the biochemical analysis, three samples of primary and secondary roots were collected at the same time for each genotype, i.e., one per tree (three replicates).

Mineral analyses were performed at three different times: 0 (control plant), D210 and 30DR. For the mineral measurements,

three samples were collected for each genotype, i.e., one per tree, and each sample is obtained by pooling eight fully-expanded leaves (three replicates).

All leaves and roots were immediately frozen in liquid nitrogen and stored at -80°C , and each leaf sample was ground to a fine powder prior performing biochemical and mineral analysis.

Branches length, trunk diameter and leaf area were measured at four different times: 0 (control plant), D70, D210, and 30DR. Branches length was measured from the top of the trunk to the top of the highest leaf for four branches of each genotype, i.e., 12 per tree (12 replicates). Trunk diameter was measured using a caliper 5 cm from the soil surface for each genotype (three replicates). For leaf area, three fully expanded leaves for each genotype, i.e., three per tree, were selected and analyzed using ImageJ imaging software, version 1.47 (nine replicates).

Evaluation of Leaf Damage

The genotypes were visually ranked according to the degree of nutrient deficiency damage after 210 days of nutrient deficiency. Nutrient deficiency damage (chlorosis, shriveled leaves) was evaluated on fully expanded leaves and scored using the following scale: level 0, no visible signs; level 1 – light green leaves with yellow leaves; level 2 – yellow leaves; level 3 – yellow shriveled leaves. The level of nutrient deficiency damage was evaluated in 15 independent leaves for each genotype.

Foliar Mineral Analysis

All ground samples were dried at $65 \pm 10^{\circ}\text{C}$ in an oven overnight, transferred into a desiccator until cooling and sent into a CIRAD laboratory (Montpellier, France) for analysis of macro- and micro-nutrients.

Total phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg), sodium (Na), bore (B), copper (Cu), iron (Fe), zinc (Zn), and manganese (Mn) in leaves were measured using an Agilent 720 simultaneous ICP-OES after double calcination including silica removal by adding hydrofluoric acid.

Nitrogen (N) content in leaves was determined by combustion using a Leco TruMac N determinator.

Measurements of Gas Exchange

Leaf net photosynthetic rate (P_{net}) and stomatal conductance (G_s) were determined using a portable photosynthesis system (LI6400, Li-COR, Lincoln, NE, United States) with the LI6400-40 Leaf Chamber. For the P_{net} and G_s measurements, carbon dioxide concentration (CO_2) was fixed at $380 \mu\text{mol mol}^{-1}$, airflow rate at $500 \mu\text{mol s}^{-1}$ and temperature at 25°C . Photosynthetic photon flux density was provided by a red-blue light source (6400-02B no. SI-710, Li-COR, Lincoln, NE, United States) in a gas exchange chamber and was set at $1400 \mu\text{mol m}^{-2} \text{s}^{-1}$.

Measurements of Chlorophyll *a* Fluorescence

Chlorophyll *a* fluorescence was detected using a leaf fluorimeter (Handy PEA, Hansatech, Instruments, Ltd.) with a resolution

of 10 s, in a non-destructive and reproducible manner. Leaves were adapted to the dark with a leaf clip for 30 min before the measurements were started. We measured the minimum value of chlorophyll *a* fluorescence (F_0) that resulted in opening of all photosystem II (PSII) centers [oxidization of all the primary quinone acceptors (QA)] and the maximum fluorescence (F_m) that results in closing of PSII centers (reduction of all QA) after the emission of a saturating 1-s flash of light (650 nm). Values of variable fluorescence ($F_v = F_m - F_0$) and the maximum quantum efficiency of PSII primary photochemistry ($F_v/F_m [= (F_m - F_0)/F_m]$) were calculated (Maxwell and Johnson, 2000).

Evaluation of Leaf Damage

The genotypes were visually ranked according to the degree of nutrient deficiency damage after 210 days of nutrient deficiency. Nutrient deficiency damage (chlorosis, shriveled leaves) was evaluated on fully expanded leaves and scored using the following scale: level 0, no visible signs; level 1 – light green leaves with yellow veins; level 2 – yellow leaves; level 3 – yellow shriveled leaves. The level of nutrient deficiency damage was evaluated in 15 independent leaves for each genotype.

Determination of Oxidative Stress and Antioxidant Levels

Oxidative markers (malondialdehyde and hydrogen peroxide) assays in roots and leaves and ascorbate and antioxidant enzyme (superoxide dismutase, catalase, ascorbate peroxidase, and dehydroascorbate reductase) in leaves were performed as described by Santini et al. (2013). A V-630 spectrophotometer was used for all measurements (Jasco, Inc., Tokyo, Japan).

Proline content was measured in leaves using the ninhydrin reaction described by Bates et al. (1973), with slight modifications. Proline was extracted from 20 mg of leaf powder, suspended in 2 mL of sulfosalicylic acid 3% and then centrifuged at 11,500 g for 15 min at 4°C. The reaction mixture contained 600 μ L of glacial acetic acid, 600 μ L of ninhydrin acid (1.25 g of ninhydrin, 30 mL of glacial acetic acid and 20 mL of orthophosphoric acid 6 M) and 300 μ L of supernatant. After 1 h of incubation at 95°C, 2 mL of toluene was added to extracted red products and samples were vortexed for 15 s. The absorbance of the organic phase was measured at 520 nm. Proline content was calculated using a standard curve of proline. A V-630 spectrophotometer was used for all measurements (Jasco, Inc., Tokyo, Japan).

Statistical Analyses

All statistical measurements were performed with R statistical software¹. The qualitative factors studied are sampling date (days) (D0, D70, D140, and D210 after nutrient deficiency, and after 30DR of recovery for leaves and D0, D70, and D210, and 30DR after recovery for roots), genotypes subjected to nutrient stress (VK, PMC, FL, CC, and CM for leaves and roots) and ploidy level nutrient stressed genotypes (VK2x/4x, PMC2x/4x,

FL4x, CC2x/4x, CM2x/4x for leaves and roots). The influence of these three factors was analyzed using a two-way ANOVA followed by LSD test at $P < 0.05$. Heat map was made to determine the differences between genotypes, ploidy level and treatments for total content in macro- and micro-nutrients and growth parameters.

The data obtained at D0, D70, D140, and D210 of nutrient deficiency and after 30DR of recovery for the nine genotypes were analyzed by principal component analysis (PCA) of centered and reduced variables with a FactomineR package bundled with R statistical software. PCA was conducted to define a clear relationship between physiological (P_{net} , G_s , and F_v/F_m ratio) and biochemical (oxidative status, antioxidant enzymes, and non-enzymatic molecules) parameters and genotypes after different periods of total nutrient deficiency and after 30 days of recovery. PCA contributed to better understanding of similarities between many measured variables and individuals.

RESULTS

In order to minimize any effect of environmental condition changes (photoperiod, temperature in the tunnel greenhouse, etc.) during the experiments, results on stressed seedlings were expressed as ratios relative to the values obtained on control seedlings. Thus, only the effect of the nutritional deprivation was taken into account. After 30 days of recovery (30DR), PMC2x and PMC4x lost all their leaves; consequently, the physiological and biochemical parameters were not measured.

Leaf Damage and Growth Parameters


Leaf damage signs were analyzed after 210 days of nutrient stress and were ranked using a four-level system based on the color of the leaves and veins (**Figure 1**). The level 0 corresponded to the control with green leaves. The least leaf damage (level 1) was found in FL4x with light green leaves. CM4x and VK2x and VK4x had very light green leaves with yellow veins and were classified as level 2. CC2x and CC4x, CM2x and PMC2x and PMC4x had yellow leaves and were classified as level 3. Overall, branches length, trunk diameter, and leaf area continued to grow after 210 days of nutrient deprivation and at after 30 days of recovery in all genotypes (**Supplementary Table S1**). Only FL4x and PMC4x did not show an increase in their leaves between D0 and D210 and D30DR (**Supplementary Table S1**).

Change in Macro- and Micro-Nutrients

Overall after 210 days of nutrient stress all genotypes showed a significant decrease in N, Fe, Mn, Zn, and B in their leaves and a slight decrease or values close to the control in P, Mg, Ca, and Na. Only K remained substantially similar to the control for all genotype (**Supplementary Table S1**).

At 30DR, N, P, Ca, Na, Fe, Mn, Cu, Zn, and B contents were lower than the control values in all genotypes (**Table 2** and **Supplementary Table S2**). Conversely, K and Mg contents fluctuated differently according to the genotypes (**Table 2** and **Supplementary Table S2**). K content increased in VK2x and

¹<http://www.R-project.org>

TABLE 2 | Total content of nitrogen (N), phosphorus (P), potassium (K), magnesium (Mg), calcium (Ca) and sodium (Na), throughout the nutrient deficiency in leaves of 2x and 4x genotypes.

Genotype	Date	Fe ppm	Mn ppm	Cu ppm	Zn ppm	B ppm
VK2x	DO	192.050	77.900	6.200	25.150	48.250
	D210	92.600	28.500	1.450	15.750	37.050
	30DR	75.550	23.650	1.700	19.650	34.950
VK4x	DO	216.300	24.400	2.400	9.250	26.200
	D210	78.450	80.800	7.500	18.450	44.550
	30DR	83.600	17.200	3.900	16.300	25.600
PMC2x	DO	234.700	84.400	7.550	12.450	33.250
	D210	43.850	13.800	0.200	9.050	19.150
PMC4x	DO	175.900	76.950	6.900	20.900	27.600
	D210	77.800	31.950	2.200	14.700	18.150
FL4x	DO	153.800	104.900	7.200	10.400	28.350
	D210	57.200	22.867	2.233	10.900	25.167
	30DR	145.250	29.050	4.550	16.300	18.100
CC2x	DO	178.850	70.250	7.900	21.350	13.900
	D210	129.800	30.750	4.000	9.750	21.150
	30DR	103.950	36.150	2.000	11.950	31.300
CC4x	DO	213.550	70.350	7.900	28.800	27.050
	D210	166.600	25.200	5.100	19.100	18.400
	30DR	61.950	22.050	3.450	18.350	16.300
CM2x	DO	153.550	57.350	7.450	21.650	41.500
	D210	131.100	22.150	2.900	11.900	37.350
	30DR	80.650	17.150	2.400	18.950	17.650
CM4x	DO	118.500	37.050	6.500	22.900	53.100
	D210	75.500	8.800	1.200	15.600	29.800
	30DR	91.050	16.000	2.350	15.400	31.400

Contents were measured after different period of nutrient deficiency: days 0 (DO) for the control and 210 (D210) and after 30 days of recovery (30DR). The results are presented as mean of three independent measurements ($n = 3$). The heat map shows the differences between genotypes, ploidy level and treatments for each macronutrient. Values are associated with color ranging from white (low) to dark green (high).

VK4x, FL4x and CC2x while Mg content increased in FL4x, CC2x and CC4x and CM2x and CM4x. In VK2x and VK4x and PMC2x and PMC4x, K and Mg contents decreased (Table 2 and Supplementary Table S2).

Change in Photosynthetic Capacities

P_{net} began to decrease at D70 in VK2x and VK4x, PMC2x and PMC4x, FL4x and CM2x and CM4x and from D140 in CC2x and CC4x (Figure 2).

This was generally associated with a decrease in G_s (Figure 2). At D210, FL4x, and CM4x had the lowest decrease both in P_{net} and G_s . Interestingly, G_s increased at D210 in PMC2x and PMC4x while P_{net} continued to decline.

F_v/F_m decreased early on at D70 in PMC2x and CM2x and CM4x, and at D140 in VK2x and VK4x, PMC4x, FL4x and CC2x and CC4x (Figure 2). However, at D210 and 30DR,

F_v/F_m was greater in FL4x, CM4x, and VK4x than in the other genotypes.

Change in Oxidative Markers (MDA and H_2O_2) in Leaves and Roots

Overall, MDA increased lately in leaves for all genotypes, excepted in VK2x (Figure 3). At D210, PMC2x and PMC4x, CC2x and CC4x and CM2x had a peak of MDA accumulation, while there was a slight increase in FL4x and PMC4x and a decrease in VK2x. After 30 days of recovery, MDA returned to the same or a lower level than in all control genotypes. H_2O_2 increased early on FL4x, but decreased to the same values as the control at D210 (Figure 3). H_2O_2 increased from D140 in VK2x and VK4x, PMC4x, CC2x and CC4x and CM2x and CM4x and from D210 in PMC2x. Like for FL4x, H_2O_2 decreased at D210 in CC2x. At 30DR, unlike




FIGURE 1 | Leaf damages after 210 days of total nutrient deficiency on the nine genotypes (0%) compared to controls (100%). Genotypes are ranked based on the leaf symptoms from the unaffected (0) to the more affected (3).

MDA, H_2O_2 values were higher than for all control genotypes, excepted in CM4x.

Measurements of H_2O_2 and MDA contents in roots throughout the experiment revealed no large changes induced by nutrient deficiency in FL4x and CM2x and CM4x, contrary to the other genotypes (Figure 3). MDA increased early on VK2x and VK4x, PMC2x and PMC4x and CC2x and CC4x, but decreased at D210 and after 30 days of recovery (30DR). H_2O_2 content also increased early on VK4x, PMC2x and PMC4x and CC2x and CC4x but decreased in VK4x, PMC4x, and CC2x after D210.

Change in Enzymatic Antioxidants

Overall, SOD and DHAR increased or were similar to control in all genotypes (Supplementary Figure S1). SOD decreased in CM2x only throughout the deficiency period. CAT decreased in VK2x and VK4x, PMC2x and PMC4x and CC2x and CC4x, whereas it increased from D140 in FL4x (Supplementary Figure S1). APX remained stable during the deficiency period in VK2x and VK4x whereas it was higher from D140 compared to control in PMC2x and PMC4x, FL4x, CC2x and CC4x and CM2x and CM4x. However, PMC2x decreased at D210 (Supplementary Figure S1).

Change in Antioxidant Molecules

The tAsa, Asa and DHA contents increased in all genotypes during the nutrient deficiency period and after the recovery period (Supplementary Figure S2). The Asa/DHA ratio decreased from D70 in FL4x and CC2x and from D140 in VK2x and VK4x, PMC2x and PMC4x, CC4x and CM2x and CM4x (Supplementary Figure S2). However, FL4x and CM4x had an increased Asa/DHA ratio at D210. At 30DR, the Asa/DHA ratio decreased in all genotypes.

The proline content decreased as early as D70 in all genotypes, but to a lesser extent in FL4x. At 30DR, proline increased but did not reach values similar to the controls (Supplementary Figure S3).

Responses of Citrus Genotypes Evaluated Through Physiological and Chemical Parameters During the Time Course of the Stress

Based on the PCA, the first two principal axes explained 49.84, 49.55, and 52.77% for 70 (D70), 140 (D140) (Supplementary Figure S4) and 210 days (D210) after nutrient deficiency, respectively, and 55.65% for 30 days after recovery (30DR) (Figure 4). Genotypes were already split in three distinct groups based on their response to nutrient deficiency. However, genotypes showed clear differences in signs of nutrient deficiency only at D210 (Figure 1).

Interestingly, at D210, PC1 was positively correlated with oxidative markers (MDA, H_2O_2 , and DHA) and negatively with photosynthetic parameters (P_{net} and F_v/F_m ratio) and Asa/DHA ratio. PC2 was positively correlated with CAT and proline, and negatively correlated with DHAR. PC1 separated cluster 1 (FL4x and CM4x) from cluster 3 (PMC2x and PMC4x) and PC2 separated cluster 2 (CC2x and CC4x, VK2x and VK4x and CM2x) from cluster 1. Thus, FL4x and CM4x were differentiated by low oxidative marker accumulation (MDA, H_2O_2 , and DHA) and a higher photosynthetic capacity than in PMC2x and PMC4x. While CC2x and CC4x, VK2x and VK4x and CM2x had intermediate values for the variables compared to FL4x and CM4x, and PMC2x and PMC4x. These findings in FL4x and CM4x were associated with a higher Asa/DHA ratio and higher CAT in FL4x and APX in CM4x.

After 30DR, PC1 was positively correlated with photosynthetic parameters (P_{net} , G_s , and F_v/F_m ratio), proline, tAsa and DHA,




FIGURE 2 | Evolution of net photosynthesis rate (P_{net}), stomatal conductance (G_s) and chlorophyll fluorescence ratio (F_v/F_m), throughout the nutrient deficiency in leaves of 2x and 4x seedlings. The white and black bars correspond to the values obtained from 2x and 4x seedlings, respectively.

(Continued)

FIGURE 2 | Continued

Concentrations were measured after different period of nutrient deficiency: days 0 (D0) for the control, 70 (D70), 140 (D140) and 210 (D210) and after 30 days of recovery (30DR). The results obtained are expressed as ratios with respect to the values obtained on control leaves which have not been subjected to stress. The results are presented as mean (\pm standard error) of nine independent measurements ($n = 9$). Data were analyzed using ANOVA and Fisher LSD tests ($P < 0.05$). Distinct capital letter indicate significant differences between all diploid seedlings at a point of the time course. Different lower case letters indicate significant value change along the time course for in 2x seedlings. For 4x seedlings, the same procedure has been followed and the results are indicated in bold roman. An asterisk indicates significant differences between 2x and 4x seedlings of the same variety at a point of the time course.

and negatively correlated with Asa/DHA. PC2 was positively correlated with G_s , F_v/F_m ratio, Asa/DHA, DHAR and MDA, and negatively correlated with DHA, H_2O_2 , CAT, and APX. PC1 separated cluster 1 (FL4x and CM4x) from cluster 2 (CC2x and CC4x and CM2x) and cluster 3 (VK2x and VK4x) and PC2 separated cluster 2 from the cluster 3. Therefore, FL4x and CM4x were characterized by higher P_{net} , G_s , and F_v/F_m ratio and proline, and lower H_2O_2 than CC2x and CC4x and CM2x, while VK2x/VK4x had intermediate values for these variables. VK2x and VK4x were distinguished by a high Asa/DHA ratio and high DHAR activity but lower SOD, CAT, and APX activities than the other genotypes.

DISCUSSION

Comparison of the Sensitivity to Nutrient Deficiency of Citrus Seedling Genotypes

All the genotypes showed leaf damage after a long period of nutrient deficiency (Figure 1). This leaf damage suggests a decrease in the quantity of photosynthetic pigments and thus photosynthetic capacity. This is typical of leaf senescence under nutrient deprivation (Deng et al., 2001; Boussadia et al., 2010). Leaf damage was associated with an alteration of foliar mineral content. This indicates that our experimental conditions were suitable for screening genotypes for nutrient deficiency. Leaf damage is a visual sign used to rank genotypes depending on the impact of the nutrient deficiency (Srivastava, 2013). In our study, we associated them with the levels of oxidative markers and macronutrient contents at the same time in order to propose a precise classification (Figure 3 and Table 2) which will be checked later with the photosynthetic capacity and the antioxidant responses (Figures 2, 4 and Supplementary Figures S1, S4). To our knowledge, there is no information about the impact of nutrient deficiency in the citrus seedling genotypes studied in this experiment.

Level 3 of leaf damage, the significant depletion of leaf macronutrient and the large accumulation of MDA in CC2x and CC4x and PMC4x, and MDA and H_2O_2 in PMC2x and CM2x, means these genotypes are sensitive to nutrient deficiency. CM4x and VK2x and VK4x were ranked as potentially tolerant because their leaves had level 2 damage, significant macronutrient depletion and low MDA and H_2O_2 accumulation. Finally FL4x was classified as resistant because its leaves had level 1 damage and low MDA and H_2O_2 accumulation. It was the only one to avoid the depletion of macronutrients.

In our study, the effect of autotetraploidization did not systematically improve the tolerance to nutrient deficiency.

CM4x was the only 4x seedling with a significantly different response to nutrient deficiency compared to its 2x counterpart. This is in agreement with previous results showing that in salt stress conditions, CM4x was more tolerant than CM2x (Podda et al., 2013). On the other hand, results in FL4x—which was the only resistant genotype—suggest that allotetraploidization would be a mean for increasing tolerance to nutrient deficiency. This needs to be supported by additional tests on other allotetraploids (Mouhaya et al., 2010). Chromosome doubling can increase tolerance to stress, have no effect or be harmful to the plant (Comai, 2005). For example, Liu et al. (2011) showed that drought, cold and salt tolerance in 4x *Dendranthema nankingense* were improved via chromosome doubling, but heat tolerance was not. In allopolyploid species, enhanced tolerance was more frequently observed than their parental species (Ainouche et al., 2009).

What Traits Are Related to Resistance to Nutrient Deficiency in Citrus Genotypes?

The “resistant” genotype, FL4x, and also the “moderately tolerant” genotype, CM4x, belonged to the same cluster (cluster 1) (Figure 4A), despite different levels of leaf damage and macronutrients content at D210. They appeared to have less disturbance of their photosynthetic process after D210 than the other genotypes as indicated by their higher F_v/F_m ratio and P_{net} (Figures 4A,B). These results suggest a less marked photo-inhibition than in other genotypes probably due to better protection of their reaction centers of photosystem II (Huang et al., 2004). FL4x would be resistant while CM4x would be more tolerant than the genotypes of the group classified as potentially tolerant (VK2x and VK4x). Indeed, many studies have found a direct influence of nutritional stress on photosynthetic functions such as PSII photochemistry either in the electron transport chain or in a modified PSII structure (Baker and Rosenqvist, 2004; Molassiotis et al., 2006; Kalaji et al., 2014). Compared to the “moderately tolerant” VK2x and VK4x and the “sensitive” CC2x and CC4x and CM2x, which were in the same cluster at D210 (cluster 2) (Figure 4A), the smallest decrease in P_{net} was related to a lesser decrease in G_s (Figure 2), lower stomatal limitation and thus decreased CO_2 flux that governs the Rubisco carboxylation sites in the chloroplast stroma (Flexas et al., 2013). The lesser decrease in G_s could also be due to a difference in the concentration of metabolites (sugars and organic acids) and a combination of stimuli that influences opening of the stomata (Boussadia et al., 2010; Lawson and Blatt, 2014; Gago et al., 2016). However, the “sensitive” PMC2x and PMC4x (cluster 3) (Figure 4A) were the only genotypes to show similar G_s values compared to control at





FIGURE 3 | Evolution of malondialdehyde (MDA) and hydrogen peroxide (H₂O₂) concentration, throughout the nutrient deficiency in leaves and roots of 2x and 4x genotypes. The white and black bars correspond to the values obtained from 2x and 4x genotypes, respectively. Concentrations were measured after different period of nutrient deficiency in: days 0 (D0) for the control, 70 (D70), 140 (D140) and 210 (D210) and after 30 days of recovery (30DR) in leaves and days 0 (D0) for the control, 70 (D70) and 210 (D210) and after 30 days of recovery (30DR) in roots. The results obtained are expressed as ratios with respect to the values obtained, respectively, on control leaves and control roots which have not been subjected to stress. The results are presented as mean (± standard error) of three independent measurements (n = 3). Data were analyzed using ANOVA and Fisher LSD tests (P < 0.05). Distinct capital letter indicate significant differences between all 2x genotypes at a point of the time course. Different lower case letters indicate significant value change along the time course for one 2x genotype. For 4x genotypes, the same procedure has been followed and the results are indicated in bold roman. An asterisk indicates significant differences between 2x and 4x genotypes of the same variety at a point of the time course.

D210. Opening of the stomata does not seem to result in the maintenance of photosynthesis in these genotypes during nutrient deficiency.

After 30 days of recovery, the high photosynthetic parameter values (*P_{net}*, *G_s*, and *F_v/F_m*) seem to indicate better recovery

and therefore reversibility of the damage caused by nutrient deficiency in FL4x and CM4x compared to the other genotypes (Figures 4C,D). VK2x and VK4x had slightly lower recovery than FL4x and CM4x, but greater than CC2x and CC4x and CM2x (Figures 4C,D). At 30DR, no leaves resisted to the nutritional



stress in PMC2x and PMC4x, which suggests a lack of recovery on injured leaves.

On the whole, the strongest photosynthetic capacities observed in FL4x and CM4x, and to a lesser extent in VK2x and VK4x than in CM2x, CC2x and CC4x and PMC2x and PMC4x could be explained by better accumulation and/or remobilization via the phloem transport of mineral elements related to photosynthesis activity (Marschner, 1995; Reich et al., 1999; Maillard et al., 2015) during nutrient deficiency. It could also be explained by a smaller decrease in transcript level related to essential components of photosynthesis (Wang et al., 2015).

Antioxidant System Contributes to Improve Tolerance to Nutrient Deficiency in Both Roots and Leaves

During nutrient deficiency, enzymatic and non-enzymatic responses were variable and more or less effective depending on the genotype. The “resistant” FL4x and “tolerant” CM4x appeared to have a better performing non-enzymatic and enzymatic antioxidant system than the other genotypes. This helps to prevent the accumulation of toxic compounds after nutrient deficiency at D210. These results were consistent with

less decrease in photosynthetic capacity (Figures 4A,B). Unlike the other genotypes, FL4x and CM4x had lower levels of oxidative markers (H_2O_2 , MDA, and DHA) and higher Asa/DHA ratio (Figures 4A,B) and proline content (especially for FL4x; Supplementary Figure S3) at D210. At the root level, the least fluctuation of MDA and H_2O_2 in FL4x and CM4x was consistent with the tolerance found in their leaves during nutrient deficiency and after 30 days of recovery (Figure 3). All results obtained confirmed the classification as resistant and tolerant of these two genotypes. An increase in MDA and H_2O_2 at D70 was observed in other genotypes suggesting an early root response to oxidative stress in comparison to their leaves (Shin and Schachtman, 2004; Shin et al., 2005).

Regarding antioxidant enzyme systems, all genotypes had a higher or similar value in SOD and DHAR activities when compared to controls (Supplementary Figure S1). However, different variation of APX and CAT activities (Supplementary Figure S1) were observed. At D210, CM4x had the greatest APX activities whereas FL4x had the greatest CAT activities (Figures 4A,B), suggesting a difference in their H_2O_2 scavenging mechanism during nutrient deficiency (Gill and Tuteja, 2010). When subjected to salt stress, CM4x was more tolerant than its respective CM2x, which was correlated with higher constitutive

levels of antioxidant enzymes and heat shock proteins (Podda et al., 2013). High APX activity was also found at D210 in “moderately tolerant” VK2x and VK4x and in the “sensitive” CM2x and CC2x/CC4x (Figures 4A,B). No association was found with all the antioxidant enzymes in the other “sensitive” PMC2x and PMC4x, confirming the importance of an efficient antioxidant enzyme system for tolerance to nutrient deficiency.

In FL4x and CM4x, the lower H₂O₂ and MDA levels can be explained by their higher or similar CAT and APX activities at D140 and D210 responsible for removing H₂O₂ formed by SOD, respectively (Azevedo et al., 1998). When excess H₂O₂ is not eliminated by CAT and APX, high levels of hydroxyl radical production may be responsible for lipid membrane damage. This damage is highlighted by the accumulation of MDA observed at D210 (Figure 3) in CC2x and CC4x, CM2x and PMC2x and PMC4x. The high DHAR activity in all genotypes did not prevent the accumulation of DHA, especially at D210 (Supplementary Figure S2). Only FL4x and CM4x showed maintenance of Asa/DHA near the control levels at D210 (Supplementary Figure S2). This could be attributable to *de novo* synthesis of ascorbate. The large increase in DHA (Supplementary Figure S2) may indicate a key role of ascorbate during nutritional deprivation. Functional collaboration between the different enzymes and antioxidant molecules seems to be crucial for an effective antioxidant system (Blokhina et al., 2003).

During recovery, specific behaviors of antioxidant machinery were observed between “resistant,” “tolerant,” “moderately tolerant,” and “sensitive” genotypes. CC2x and CC4x and CM2x had very high H₂O₂ accumulations which was consistent with the large disturbance in photosynthetic capacity, comparatively to the other genotypes (Figures 4C,D). In FL4x and CM4x, the better recovery was probably due to the higher activity of their non-enzymatic (proline and Asa), and to a lesser extent enzymatic (SOD, CAT, and APX), systems (Figures 4C,D).

CONCLUSION

Our study used experimental design conditions that revealed differences between Citrus genotypes in terms of their resistance to nutrient deficiency. Future studies can be performed to improve citrus production in reduced input conditions. Using this experimental set up, the next steps would be to assay the mineral elements in plant organs for these different genotypes, to test the effects on growth and fruit production and quality.

This study demonstrated that total nutrient deficiency impairs the photosynthetic capacity as well as the oxidative metabolism of all citrus genotypes in different proportions. Nutrient deficiency

highly affects the photosynthesis efficiency in sensitive genotypes by strong stomatal limitation and alteration of photosystem II. During nutrient deficiency and after recovery, the resistant FL4x and tolerant CM4x were the only genotypes characterized by lower accumulation of oxidative markers (MDA and H₂O₂) in leaves and roots due to a more effective antioxidant system. However, chromosome doubling does not improve tolerance in the other genotypes. The fact that these two 4x genotypes limit the damage caused to plant growth during nutrient deficiency suggests they can be used in citrus orchards for their horticultural performance as rootstock. This would reduce the amount of fertilizer needed and have a positive impact on soil biology while preserving growth. It would be interesting to confirm these results by testing the rootstocks/scion combination in order to evaluate whether use of these 4x rootstocks can improve the tolerance to nutrient deficiency of the scion. The impact of these combinations on growth, fruit production and fruit quality of the scion during nutrient deficiency should also be studied.

AUTHOR CONTRIBUTIONS

JO collected the test data, performed the statistical analyses, interpreted the results, and drafted the manuscript. JS interpreted the results and drafted the manuscript. LB, JG, SH, FL, and RM helped to draft the manuscript. PM helped to collect the test data.

FUNDING

JO was subsidized by the “Collectivité Territoriale de Corse” (CTC). This work was funded by the “Collectivité Territoriale de Corse” as part of the research project “Sélection de porte-greffes pour le développement d’une agrumiculture durable en Corse.”

ACKNOWLEDGMENTS

We thank Jean-Claude Ribaut for hosting this study at the AREFLEC experimental station, and Gilles Paolacci and Paul Martin for installing and monitoring the ferti-irrigation system in the greenhouse.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2019.00127/full#supplementary-material>

REFERENCES

- Ainouche, M. L., Fortune, P. M., Salmon, A., Parisod, C., Grandbastien, M.-A., Fukunaga, K., et al. (2009). Hybridization, polyploidy and invasion: lessons from *Spartina* (Poaceae). *Biol. Invasion*. 11:1159. doi: 10.1007/s10530-008-9383-2
- Allario, T., Brumos, J., Colmenero-Flores, J. M., Iglesias, D. J., Pina, J. A., Navarro, L., et al. (2013). Tetraploid Rangpur lime rootstock increases drought tolerance via enhanced constitutive root abscisic acid production. *Plant. Cell. Environ.* 36, 856–868. doi: 10.1111/pce.12021
- Apel, K., and Hirt, H. (2004). Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* 55, 373–399. doi: 10.1146/annurev.arplant.55.031903.141701
- Azevedo, R. A., Alas, R. M., Smith, R. J., and Lea, P. J. (1998). Response of antioxidant enzymes to transfer from elevated carbon dioxide to air and ozone fumigation, in the leaves and roots of wild-type and a catalase-deficient

- mutant of barley. *Physiol. Plant.* 104, 280–292. doi: 10.1034/j.1399-3054.1998.1040217.x
- Baker, N. R., and Rosenqvist, E. (2004). Applications of chlorophyll fluorescence can improve crop production strategies: an examination of future possibilities. *J. Exp. Bot.* 55, 1607–1621. doi: 10.1093/jxb/erh196
- Bates, L. S., Waldren, R. P., and Teare, I. D. (1973). Rapid determination of free proline for water-stress studies. *Plant Soil* 39, 205–207. doi: 10.1016/j.dental.2010.07.006
- Beest, M., Le Roux, J. J., Richardson, D. M., Brysting, A. K., Suda, J., Kubešová, M., et al. (2012). The more the better? The role of polyploidy in facilitating plant invasions. *Ann. Bot.* 109, 19–45. doi: 10.1093/aob/mcr277
- Blokhina, O., Virolainen, E., and Fagerstedt, K. V. (2003). Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Ann. Bot.* 91, 179–194. doi: 10.1093/aob/mcf118
- Boussadia, O., Steppe, K., Zgallai, H., El Hadj, S. B., Braham, M., Lemeur, R., et al. (2010). Effects of nitrogen deficiency on leaf photosynthesis, carbohydrate status and biomass production in two olive cultivars ‘Meski’ and ‘Koroneiki. *Sci. Hortic.* 123, 336–342. doi: 10.1016/j.scienta.2009.09.023
- Brochmann, C., Brysting, A. K., Alsos, I. G., Borgen, L., Grundt, H. H., Scheen, A.-C., et al. (2004). Polyploidy in arctic plants. *Biol. J. Linnean. Soc.* 82, 521–536. doi: 10.1111/j.1095-8312.2004.00337.x
- Cameron, J., and Frost, H. B. (1968). “Genetics, breeding, and nucellar embryony,” in *The Citrus Industry*, eds W., Reuthen, H. J., Webber, L. D., Batchelor (Berkeley, CA: University of California Press), 325–370.
- Chen, Z. J. (2007). Genetic and epigenetic mechanisms for gene expression and phenotypic variation in plant polyploids. *Annu. Rev. Plant Biol.* 58, 377–406. doi: 10.1146/annurev.arplant.58.032806.103835
- Chiou, T. J. (2007). The role of microRNAs in sensing nutrient stress. *Plant. Cell. Environ.* 30, 323–332. doi: 10.1111/j.1365-3040.2007.01643.x
- Comai, L. (2005). The advantages and disadvantages of being polyploid. *Nat. Rev. Genet.* 6, 836–846. doi: 10.1038/nrg1711
- Deng, Z., Cao, S., Tang, Y., Lu, W., and Zhang, R. (2001). Comparison on some aspects of photosynthetic decline of wild rice and 9516. *J. Henan Norm. Univ.* 30, 73–77.
- Doyle, J. J., and Egan, A. N. (2010). Dating the origins of polyploidy events. *New Phytol.* 186, 73–85. doi: 10.1111/j.1469-8137.2009.03118.x
- Ehrendorfer, F. (1980). “Polyploidy and distribution,” in *Polyploidy*, (Berlin: Springer), 45–60. doi: 10.1007/978-1-4613-3069-1_3
- Ericsson, T. (1995). Growth and shoot: root ratio of seedlings in relation to nutrient availability. *Plant Soil* 16, 205–214. doi: 10.1007/BF00029330
- Flexas, J., Niinemets, Ü, Gallé, A., Barbour, M. M., Centritto, M., Diaz-Espejo, A., et al. (2013). Diffusional conductances to CO₂ as a target for increasing photosynthesis and photosynthetic water-use efficiency. *Photosynth. Res.* 117, 45–59. doi: 10.1007/s11120-013-9844-z
- Froelicher, Y., Bassene, J. B., Jedidi-Neji, E., Dambier, D., Morillon, R., Bernardini, G., et al. (2007). Induced parthenogenesis in mandarin for haploid production: induction procedures and genetic analysis of plantlets. *Plant Cell Rep.* 26, 937–944. doi: 10.1007/s00299-007-0314-y
- Gago, J., de Menezes Daloso, D., Figueroa, C. M., Flexas, J., Fernie, A. R., and Nikoloski, Z. (2016). Relationships of leaf Net photosynthesis, stomatal conductance, and mesophyll conductance to primary metabolism: a multispecies meta-analysis approach. *Plant Physiol.* 171, 265–279. doi: 10.1104/pp.15.01660
- Gill, S. S., and Tuteja, N. (2010). Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol. Biochem.* 48, 909–930. doi: 10.1016/j.plaphy.2010.08.016
- Guerra, D., Wittmann, M. T. S., Schwarz, S. F., Souza, P. V. D., de Gonzatto, M. P., and Weiler, R. L. (2014). Comparison between diploid and tetraploid citrus rootstocks: morphological characterization and growth evaluation. *Bragantia* 73, 1–7. doi: 10.1590/brag.2014.007
- Hao, G. Y., Lucero, M. E., Sanderson, S. C., Zacharias, E. H., and Holbrook, N. M. (2013). Polyploidy enhances the occupation of heterogeneous environments through hydraulic related trade-offs in *Atriplex canescens* (Chenopodiaceae). *New Phytol.* 197, 970–978. doi: 10.1111/nph.12051
- Huang, Z.-A., Jiang, D.-A., Yang, Y., Sun, J.-W., and Jin, S.-H. (2004). Effects of nitrogen deficiency on gas exchange, chlorophyll fluorescence, and antioxidant enzymes in leaves of rice plants. *Photosynthetica* 42, 357–364. doi: 10.1023/B:PHOT.0000046153.08935.4c
- Kalaji, H. M., Oukarroum, A., Alexandrov, V., Kouzmanova, M., Brestic, M., Zivcak, M., et al. (2014). Identification of nutrient deficiency in maize and tomato plants by in vivo chlorophyll a fluorescence measurements. *Plant Physiol. Biochem.* 81, 16–25. doi: 10.1016/j.plaphy.2014.03.029
- Kandlbinder, A., Finkemeier, I., Wormuth, D., Hanitzsch, M., and Dietz, K.-J. (2004). The antioxidant status of photosynthesizing leaves under nutrient deficiency: redox regulation, gene expression and antioxidant activity in *Arabidopsis thaliana*. *Physiol. Plant* 120, 63–73. doi: 10.1111/j.0031-9317.2004.0272.x
- Krug, C. A. (1943). Chromosome numbers in the subfamily Aurantioideae with special reference to the genus Citrus. *Bot. Gaz.* 104, 602–611. doi: 10.1086/335173
- Lawson, T., and Blatt, M. R. (2014). Stomatal size, speed, and responsiveness impact on photosynthesis and water use efficiency. *Plant Physiol.* 164, 1556–1570. doi: 10.1104/pp.114.237107
- Levin, D. A. (1983). Polyploidy and novelty in flowering plants. *Am. Nat.* 122, 1–25. doi: 10.1086/284115
- Liu, S., Chen, S., Chen, Y., Guan, Z., Yin, D., and Chen, F. (2011). In vitro induced tetraploid of *Dendranthema nankingense* (Nakai) Tzvel. shows an improved level of abiotic stress tolerance. *Sci. Hortic.* 127, 411–419. doi: 10.1016/j.scienta.2010.10.012
- Maillard, A., Diquélou, S., Billard, V., Lainé, P., Garnica, M., Prudent, M., et al. (2015). Leaf mineral nutrient remobilization during leaf senescence and modulation by nutrient deficiency. *Front. Plant Sci.* 6:317. doi: 10.3389/fpls.2015.00317
- Marschner, H. (1995). “Functions of mineral nutrients and macronutrients,” in *Mineral Nutrition of Higher Plants*, 2nd Edn, (Cambridge: Academic Press), 313–404. doi: 10.1016/B978-012473542-2/50011-0
- Mauricio, R., Rausher, M. D., and Burdick, D. S. (1997). Variation in the defense strategies of plants: are resistance and tolerance mutually exclusive? *Ecology* 78, 1301–1310.
- Maxwell, K., and Johnson, G. N. (2000). Chlorophyll fluorescence a practical guide. *J. Exp. Bot.* 51, 659–668. doi: 10.1093/jxb/51.345.659
- Mittler, R. (2002). Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* 7, 405–410. doi: 10.1016/S1360-1385(02)02312-9
- Mittler, R., Vanderauwera, S., Gollery, M., and Van Breusegem, F. (2004). Reactive oxygen gene network of plants. *Trends Plant Sci.* 9, 490–498. doi: 10.1016/j.tplants.2004.08.009
- Molassiotis, A., Tanou, G., Diamantidis, G., Patakas, A., and Therios, I. (2006). Effects of 4-month Fe deficiency exposure on Fe reduction mechanism, photosynthetic gas exchange, chlorophyll fluorescence and antioxidant defence in two peach rootstocks differing in Fe deficiency tolerance. *J. Plant Physiol.* 163, 176–185. doi: 10.1016/j.jplph.2004.11.016
- Mouhaya, W., Allario, T., Brumos, J., Andrés, F., Froelicher, Y., Luro, F., et al. (2010). Sensitivity to high salinity in tetraploid citrus seedlings increases with water availability and correlates with expression of candidate genes. *Funct. Plant Biol.* 37, 674–685. doi: 10.1071/FP10035
- Mutlu, S., Karadağoğlu, Ö, Atıcı, Ö, and Nalbantoğlu, B. (2013). Protective role of salicylic acid applied before cold stress on antioxidative system and protein patterns in barley apoplast. *Biol. Plant* 57, 507–513. doi: 10.1007/s10535-013-0322-4
- Ollitrault, P., Dambier, D., Seker, M., and Froelicher, Y. (2000). Rootstock breeding by somatic hybridisation for the Mediterranean citrus industry. *Acta Hortic.* 535, 157–162. doi: 10.1007/s00299-010-1000-z
- Ollitrault, P., and Jacquemond, C. (1994). Facultative apomixis, spontaneous polyploidization and inbreeding in Citrus volkameriana seedlings. *Fruits* 49, 479–480.
- Oustric, J., Morillon, R., Luro, F., Herbette, S., Lourkisti, R., Giannettini, J., et al. (2017). Tetraploid carrizo citrange rootstock (*Citrus sinensis* Osb x *Poncirus trifoliata* L. Raf.) enhances natural chilling stress tolerance of common clementine (*Citrus clementina* Hort. ex Tan). *J. Plant Physiol.* 214, 108–115. doi: 10.3389/fpls.2016.00701
- Podda, A., Checcucci, G., Mouhaya, W., Centeno, D., Rofidal, V., Del Carratore, R., et al. (2013). Salt-stress induced changes in the leaf proteome of diploid and tetraploid mandarins with contrasting Na⁺ and Cl⁻ accumulation behaviour. *J. Plant Physiol.* 170, 1101–1112. doi: 10.1016/j.jplph.2013.03.006
- Polle, A., Pfinmann, T., Chakrabarti, S., and Rennenberg, H. (1993). The effects of enhanced ozone and enhanced carbon dioxide concentrations on biomass,

- pigments and antioxidative enzymes in spruce needles (*Picea abies*, L.). *Plant Cell Environ.* 16, 311–316. doi: 10.1111/j.1365-3040.1993.tb00874.x
- Reich, P. B., Ellsworth, D. S., Walters, M. B., Vose, J. M., Gresham, C., Volin, J. C., et al. (1999). Generality of leaf trait relationships: a test across six biomes. *Ecology* 80, 1955–1969. doi: 10.1890/0012-9658(1999)080[1955:GOLTRA]2.0.CO;2
- Ruiz, M., Quiñones, A., Martínez-Alcántara, B., Aleza, P., Morillon, R., Navarro, L., et al. (2016a). Effects of salinity on diploid (2x) and doubled diploid (4x) *Citrus macrophylla* genotypes. *Sci. Hortic.* 207, 33–40. doi: 10.1016/j.plaphy.2018.12.020
- Ruiz, M., Quiñones, A., Martínez-Alcántara, B., Aleza, P., Morillon, R., Navarro, L., et al. (2016b). Tetraploidy enhances boron-excess tolerance in carrizo citrange (*Citrus sinensis* L. Osbx *Poncirus trifoliata* L. Raf.). *Front. Plant Sci.* 7:701. doi: 10.3389/fpls.2016.00701
- Ruiz, M., Quiñones, A., Martínez-Cuenca, M.-R., Aleza, P., Morillon, R., Navarro, L., et al. (2016c). Tetraploidy enhances the ability to exclude chloride from leaves in Carrizo citrange seedlings. *J. Plant. Physiol.* 205, 1–10. doi: 10.1016/j.jplph.2016.08.002
- Saleh, B., Allario, T., Dambier, D., Ollitrault, P., and Morillon, R. (2008). Tetraploid citrus rootstocks are more tolerant to salt stress than diploid. *C. R. Biol.* 331, 703–710. doi: 10.1016/j.crv.2008.06.007
- Santini, J., Giannettini, J., Pailly, O., Herbette, S., Ollitrault, P., Berti, L., et al. (2013). Comparison of photosynthesis and antioxidant performance of several Citrus and Fortunella species (Rutaceae) under natural chilling stress. *Trees* 27, 71–83. doi: 10.1007/s00468-012-0769-5
- Shin, R., Berg, R. H., and Schachtman, D. P. (2005). Reactive oxygen species and root hairs in Arabidopsis root response to nitrogen, phosphorus and potassium deficiency. *Plant Cell Physiol.* 46, 1350–1357. doi: 10.1093/pcp/pci145
- Shin, R., and Schachtman, D. P. (2004). Hydrogen peroxide mediates plant root cell response to nutrient deprivation. *Proc. Natl. Acad. Sci. U.S.A.* 101, 8827–8832. doi: 10.1073/pnas.0401707101
- Soltis, P. S., and Soltis, D. E. (2009). The role of hybridization in plant speciation. *Annu. Rev. Plant Biol.* 60, 561–588. doi: 10.1146/annurev.arplant.043008.092039
- Srivastava, A. K. (2013). Nutrient deficiency symptomatology in citrus: an effective diagnostic tool or just an aid for post-mortem analysis. *Agric. Adv.* 2, 177–194.
- Stebbins, G. L. (1947). Types of polyploids: their classification and significance. *Adv. Genet.* 1, 403–429. doi: 10.1016/S0065-2660(08)60490-3
- Sunkar, R., Chinnusamy, V., Zhu, J., and Zhu, J. K. (2007). Small RNAs as big players in plant abiotic stress responses and nutrient deprivation. *Trends Plant Sci.* 12, 301–309. doi: 10.1016/j.tplants.2007.05.001
- Syvertsen, J. P., Lee, L. S., and Grosser, J. W. (2000). Limitations on growth and net gas exchange of diploid and tetraploid Citrus rootstock cultivars grown at elevated CO₂. *J. Am. Soc. Hortic. Sci.* 125, 228–234.
- Tan, F.-Q., Tu, H., Liang, W.-J., Long, J.-M., Wu, X.-M., Zhang, H.-Y., et al. (2015). Comparative metabolic and transcriptional analysis of a doubled diploid and its diploid citrus rootstock (*Citrus junos* cv. *Ziyang xiangcheng*) suggests its potential value for stress resistance improvement. *BMC Plant Biol.* 15:89. doi: 10.1186/s12870-015-0450-4
- Tewari, R. K., Kumar, P., Tewari, N., Srivastava, S., and Sharma, P. N. (2004). Macronutrient deficiencies and differential antioxidant responses—influence on the activity and expression of superoxide dismutase in maize. *Plant Sci.* 166, 687–694. doi: 10.1016/j.plantsci.2003.11.004
- Udall, J. A., and Wendel, J. F. (2006). Polyploidy and crop improvement. *Crop Sci.* 46:S–3. doi: 10.2135/cropsci2006.07.0489tpg
- Vieira, D. D. S., Emiliani, G., Michelozzi, M., Centritto, M., Luro, F., Morillon, R., et al. (2016). Polyploidization alters constitutive content of volatile organic compounds (VOC) and improves membrane stability under water deficit in Volkamer lemon (*Citrus limonia* Osb.) leaves. *Environ. Exp. Bot.* 126, 1–9. doi: 10.1016/j.envexpbot.2016.02.010
- Wang, L., Jiang, J., Song, A., Wang, H., Li, P., Guan, Z., et al. (2015). Comparative transcriptome analysis of *Chrysanthemum nankingense* in response to nitrogen deficiency. *Sci. Hortic.* 195, 101–107. doi: 10.1016/j.scienta.2015.09.001
- Wendel, J. F. (2000). Genome evolution in polyploids. *Plant Mol. Biol.* 42, 225–249. doi: 10.1023/A:1006392424384
- Xu, J., Li, Y., Sun, J., Du, L., Zhang, Y., Yu, Q., et al. (2013). Comparative physiological and proteomic response to abrupt low temperature stress between two winter wheat cultivars differing in low temperature tolerance. *Plant Biol.* 15, 292–303. doi: 10.1111/j.1438-8677.2012.00639.x

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Oustric, Morillon, Luro, Herbette, Martin, Giannettini, Berti and Santini. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.