### Natural variation in life history strategy of *Arabidopsis thaliana* determines stress responses to drought and insects of different feeding guilds

NELSON H. DAVILA OLIVAS,\* ENRIC FRAGO,\* MANUS P. M. THOEN,\*†‡ KAREN J. KLOTH, \*†‡ FRANK F. M. BECKER,§ JOOP J. A. VAN LOON,\* GERRIT GORT,¶

JOOST J. B. KEURENTJES,§ JOOST VAN HEERWAARDEN\*\* and MARCEL DICKE\* 🝺

\*Laboratory of Entomology, Wageningen University, Wageningen, The Netherlands, †Wageningen Plant Research, Wageningen University and Research, Wageningen, The Netherlands, ‡Laboratory of Plant Physiology, Wageningen University, Wageningen, The Netherlands, §Laboratory of Genetics, Wageningen University, Wageningen, The Netherlands, ¶Biometris, Wageningen University and Research, Wageningen, The Netherlands, \*\*Plant Production Systems Group, Wageningen University, Wageningen, The Netherlands

### Abstract

Plants are sessile organisms and, consequently, are exposed to a plethora of stresses in their local habitat. As a result, different populations of a species are subject to different selection pressures leading to adaptation to local conditions and intraspecific divergence. The annual brassicaceous plant Arabidopsis thaliana is an attractive model for ecologists and evolutionary biologists due to the availability of a large collection of resequenced natural accessions. Accessions of A. thaliana display one of two different life cycle strategies: summer and winter annuals. We exposed a collection of 308 European Arabidopsis accessions, that have been genotyped for 250K SNPs, to a range of stresses: one abiotic stress (drought), four biotic stresses (Pieris rapae caterpillars, Plutella xylostella caterpillars, Frankliniella occidentalis thrips and Myzus persicae aphids) and two combined stresses (drought plus P. rapae and Botrytis cinerea fungus plus P. rapae). We identified heritable genetic variation for responses to the different stresses, estimated by narrow-sense heritability. We found that accessions displaying different life cycle strategies differ in their response to stresses. Winter annuals are more resistant to drought, aphids and thrips and summer annuals are more resistant to P. rapae and P. xylostella caterpillars. Summer annuals are also more resistant to the combined stresses of drought plus P. rapae and infection by the fungus Botryris cinerea plus herbivory by P. rapae. Adaptation to drought displayed a longitudinal gradient. Finally, trade-offs were recorded between the response to drought and responses to herbivory by caterpillars of the specialist herbivore P. rapae.

*Keywords*: biotic stress, drought, fungal pathogen, generalist, herbivory, specialist, summer annual, winter annual

Received 27 September 2015; revision received 21 February 2017; accepted 27 February 2017

#### Introduction

Biotic and abiotic conditions vary in space and time. As a result, different populations of a species are exposed to different selection pressures leading to adaptation to local conditions and intraspecific divergence (Kaltz &

Correspondence: Marcel Dicke, Fax: +31317484075; E-mail: marcel.dicke@wur.nl Shykoff 1998; Manel *et al.* 2003; Kawecki & Ebert 2004). Locally adapted populations are, therefore, expected to outperform non-native populations (Kawecki & Ebert 2004; Savolainen *et al.* 2013). Local adaptation is particularly relevant in the face of the global changes our planet is exposed to, because locally adapted populations can become maladapted if environmental conditions change. Mathematical models indicate that climate change will increase the incidence of extreme

<sup>© 2017</sup> The Authors. *Molecular Ecology* Published by John Wiley & Sons Ltd.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

temperatures, drought and flooding events (Zhao & Running 2010; Sheffield *et al.* 2012; Wheeler & von Braun 2013). With global increase in mean temperatures, it is expected that species will move towards higher elevation and latitudes (Devictor *et al.* 2012). This will result in changes in species interactions and the intensity of selection pressure that biotic and abiotic conditions impose on organisms (Hughes 2000). Thus, understanding how organisms adapt to environmental conditions and investigating how responses to environmental stresses covary with geographic gradients are central issues in exploring the ecological consequences of global change.

There is a wealth of experimental evidence for local adaptation in plants and animals (Bradshaw & Holzapfel 2001; Savolainen et al. 2007; Leimu & Fischer 2008; Hereford 2009; Leinonen et al. 2009, 2011; Zust et al. 2012; Sternberg & Thomas 2014). Local adaptation has been demonstrated along longitudinal, latitudinal and elevational gradients in several plant species (Mikola 1982; Olsson & Agren 2002; Stinchcombe et al. 2004; Montesinos-Navarro et al. 2012; Alberto et al. 2013; Debieu et al. 2013). Important variables such as temperature and precipitation covary with these gradients, and thus, the latter are useful indicators of species adaptation to local environmental conditions (Fukami & Wardle 2005; Walker et al. 2010; Halbritter et al. 2013; Sundqvist et al. 2013; Manzano-Piedras et al. 2014). Traits involved in interactions between organisms and their environment have also been observed to display variation across gradients. For example, tolerance to abiotic stresses such as drought, salinity and low temperature is known to vary with elevation and latitude (Baxter et al. 2010; Hancock et al. 2011; Alberto et al. 2013; Paccard et al. 2014). Resistance to biotic stresses such as insect herbivory is also known to vary along geographic gradients. Evidence suggests that herbivory is more intense at lower latitudes and elevations, due to increased herbivore diversity and, hence, plants will be under selection for increased defences (Sanders et al. 2003; Andrew & Hughes 2005; Moles et al. 2011; Halbritter et al. 2013; Sundqvist et al. 2013; Rasmann et al. 2014). A study on two piperaceous plant species, for example, reported higher incidence of herbivory towards the equator (Salazar & Marquis 2012). Similarly, ecotypes of Plantago lanceolata and Vicia sepium from low elevations were more resistant to herbivory than plants from high-elevation ecotypes (Pellissier et al. 2014). Although the hypothesis that 'the lower the latitude and elevation, the higher the level of plant defences' (Andrew & Hughes 2005; Moles et al. 2011) has been supported in some studies, others failed to provide support. Therefore, further experiments are needed that address plant responses throughout extensive geographic areas. As firmly rooted organisms, plants are exposed to a wide diversity of biotic and abiotic stresses which commonly occur simultaneously (Atkinson & Urwin 2012; Kissoudis *et al.* 2014; Suzuki *et al.* 2014). Much progress has been made in understanding how plants adapt to their environment and what the underlying genetic mechanisms are. Yet, we still know rather little about how plants integrate responses to multiple simultaneous stresses, and how this is shaped by local adaptation.

Several methodologies have been developed to investigate local adaptation, including reciprocal transplants and common garden experiments (Kawecki & Ebert 2004). When screening populations from a large geographic area, however, the former approach is often impractical for legal, logistic or ethical reasons, and hence, growing different genotypes in a common environment is a good alternative. Under these circumstances, environmental effects can be removed and variation in phenotypic responses can be assessed. This approach is particularly useful because it allows to investigate the role of a particular agent as driver of population differentiation (Kawecki & Ebert 2004; Alberto et al. 2013; Manzano-Piedras et al. 2014). Despite the limitations of inferring local adaptation from correlative approaches in a common environment (Savolainen et al. 2013), this method represents a powerful tool for detecting selection along environmental gradients (Gomaa et al. 2011; Montesinos-Navarro et al. 2011, 2012; Pico 2012; Debieu et al. 2013; Manzano-Piedras et al. 2014).

Model organisms like A. thaliana have become attractive to ecologists and evolutionary biologists because large collections of natural accessions with known genomic sequences are available (Shindo et al. 2007; Fournier-Level et al. 2011; Weigel 2012). Furthermore, recent studies have demonstrated the feasibility of using these natural accessions for genomewide association studies, allowing to relate variation in a trait with variation in the genes underlying that trait (Atwell et al. 2010; Weigel 2012; Bac-Molenaar et al. 2015; Thoen et al. 2017). In A. thaliana, variation in several life history traits along latitudinal and altitudinal gradients has been demonstrated, including (i) variation in flowering time, seed dormancy and growth rate (Stinchcombe et al. 2004; Montesinos-Navarro et al. 2011; Debieu et al. 2013); (ii) variation in resistance against abiotic stresses such as salt and drought stress (McKay et al. 2003; Baxter et al. 2010; Juenger 2013; Easlon et al. 2014); and (iii) variation in defence-related traits. For example, A. thaliana displays a latitudinal and longitudinal gradient for diversity in glucosinolate profiles, a main class of defensive metabolites in the Brassicaceae (Brachi et al. 2015) and, through a longitudinal gradient, this correlates with the abundance of two aphid species, that is *Brevicoryne brassicae* and *Lipaphis erysimi* (Zust *et al.* 2012).

An important characteristic of A. thaliana is that accessions exhibit one of two different life cycle strategies: summer and winter annuals (Pigliucci 1998; Koornneef et al. 2004; Shindo et al. 2007). Winter annuals are accessions that germinate in late summer and autumn, overwinter as rosette and flower in spring. Winter annuals require a vernalization period to flower. On the other hand, summer annuals or rapid cyclers germinate in spring, flower in summer and do not require vernalization for flowering (Koornneef et al. 2004; Stinchcombe et al. 2004; Shindo et al. 2007). Furthermore, the winter annual life cycle strategy is mostly expressed in temperate areas, whereas the summer annuals occur in warmer regions (Johanson et al. 2000; Michaels et al. 2003). This plasticity in life cycle strategies has been hypothesized to be the basis for the highly adaptive capacity of A. thaliana (Shindo et al. 2007). For instance, there is a strong positive genetic correlation between flowering time and water-use efficiency such that late-flowering accessions display higher water-use efficiency than early-flowering ones (McKay et al. 2003; Juenger 2013; Easlon et al. 2014). Despite the different phenological, morphological and stress-response traits observed in A. thaliana accessions displaying different life cycle strategies, these differences have rarely been considered when studying local adaptation and responses to abiotic and biotic stresses.

The overall goal of this study was to address this knowledge gap. We have investigated whether accessions of the two *Arabidopsis* life history strategies differ in resistance to a range of biotic and abiotic stresses. We used a collection of 308 European *A. thaliana* accessions and exposed them in a controlled environment to a diverse range of abiotic and biotic stresses.

Phytohormones are major players in the signalling network underlying responses to both biotic and abiotic stresses (Pieterse et al. 2009). For instance, chewing insect herbivores elicit especially the jasmonic acid (JA), abscisic acid (ABA) and ethylene (ET) signalling pathways; phloem-sucking insects and biotrophic microbial pathogens elicit especially the salicylic acid (SA) pathway; and drought elicits the abscisic acid (ABA) pathway (Pieterse et al. 2009). These phytohormonal responses exhibit extensive crosstalk, resulting in specific changes in plant phenotype in response to individual stresses (De Vos et al. 2005; Pieterse et al. 2009). As biotic stresses, we selected four insect herbivores that induce different phytohormonal pathways so as to cover a range of stress types. The herbivores used in this study are two leaf-chewing caterpillars, Pieris rapae and Plutella xylostella, and two piercing-sucking insects, the phloem-feeding aphid Myzus persicae and the cellcontent-feeding thrips *Frankliniella occidentalis*. Moreover, stress combinations were imposed by combining drought plus *P. rapae* and the necrotrophic fungus *Botrytis cinerea* plus *P. rapae*. *Arabidopsis thaliana*'s response to *P. rapae*, *P. xylostella*, *F. occidentalis*, *B. cinerea* and drought as single stresses is highly diverse, yet at the same time regulated by the plant hormones JA and/or ABA (De Vos *et al.* 2005; Broekgaarden *et al.* 2007; Verhage *et al.* 2011). In contrast, the response to the aphid *M. persicae* is regulated by the plant hormone SA (De Vos *et al.* 2005).

The main goal of this study was to investigate whether winter and summer annuals differ in their responses to a set of different stresses, including biotic and abiotic stresses and some stress combinations. To further assess the basis for the expected differential responses, we also addressed the following questions for the population of Arabidopsis accessions that we used here: (a) Is there heritable genetic variation in A. thaliana's responses to the different stresses studied? (b) Does the proportion of winter and summer annuals vary along latitude, longitude and elevation? (c) Do A. thaliana plants from different latitudes, longitudes and elevations differ in their responses to these stresses? Life history theory predicts that for a specific genotype, the expression of a particular defensive or resistance trait will constrain the expression of others, so we finally asked the following: (d) Are there tradeoffs among A. thaliana responses to different stresses? To identify candidate genes underlying the responses of winter annuals vs. summer annuals, we carried out a multitrait genomewide association analysis.

### Materials and methods

### Arabidopsis thaliana populations and molecular markers

In this study, we used a population of 350 Arabidopsis thaliana (L.) Heynh. accessions from the Hapmap popu-(http://bergelson.uchicago.edu/wp-content/ lation uploads/2015/04/Justins-360-lines.xls) (Fig. S1, Supporting information). To avoid geographic outliers, however, our analyses were limited to the data on the European accessions found at a latitude  $\geq$ 30, longitude between -50 and 50 and elevation ≤2000 m (Fig. S2, Supporting information). This resulted in a subset of 308 accessions on which all further analyses were performed. The Hapmap population has been genotyped for 250K bi-allelic SNPs (Baxter et al. 2010; Platt et al. 2010; Chao et al. 2012), and after quality control and imputation, this SNP set was reduced to a set of 214 051 SNPs. To get a deeper understanding of these correlates, we obtained a set of environmental variables

from the locations where the *A. thaliana* accessions had been collected (Data S1, Supporting information). Sources of these environmental data are summarized in Table S1 (Supporting information). The environmental variables were then correlated with latitude, longitude and elevation through Spearman correlation tests (Fig. S3, Supporting information).

### Genetic diversity in European accessions from the Hapmap population

Determination of geographically informative genetic groups was done as follows. Principal component analysis was performed on the scaled 0,1 molecular marker matrix of dimensions  $n \times m$ , where *n* is the number of individuals and m the number of markers (van Heerwaarden et al. 2011, 2013; Odong et al. 2013). Spatial autocorrelation of individual PCs was determined by Moran's I statistic. Geographic coordinates were converted to spatial weight classes by the functions graph2nb and nb2listw in the R package spdep. The first k principal components with the highest spatial autocorrelation and associated P values below 0.001 were retained as spatially informative and were used to calculate a Euclidean distance matrix. A dendrogram was produced by Ward clustering, and genetic groups were assigned by splitting the dendrogram into groups using the R function cutree. Phenotypic differences between genetic groups were tested by ANOVA after correction for life cycle strategy.

### *Classification of accessions as winter and summer annuals*

Flowering time and flowering time after vernalization under greenhouse conditions have been recorded for this plant population as reported by Bac-Molenaar *et al.* (2016). Accessions were classified as winter annuals if vernalization was required for flowering (flowering time  $\geq$ 75 days) or as summer annuals otherwise (Data S1, Supporting information).

### Plants, insects and fungi

Arabidopsis thaliana plants were grown in a controlled environmental chamber at  $24 \pm 1$  °C,  $70 \pm 10\%$  relative humidity, 200 µmol m<sup>-2</sup> s<sup>-1</sup> photosynthetically active radiation from fluorescent lights (TL-D 58W/840; www. philips.com) and a diurnal cycle of 8:16 L:D. In all experiments, seeds were vernalized at 4 °C for 5 days to induce even germination. Plants were individually grown in 0.08 L pots (http://uk.poeppelmann.com/te ku/home/) in a pasteurized (4 h at 80 °C) commercial *Arabidopsis* potting soil, which was mixed 1:1 (v/v) with autoclaved sand in experiment (1). In experiments (2), (3) and (4) (see later), plants were grown in pasteurized (4 h at 80 °C) commercial potting soil. Individual pots were accommodated in trays, randomly distributed within the growth chamber. Plants were watered three times per week by adding water to the tray. Once per week, they received *Steinernema feltiae* entomopathogenic nematodes (Entonem; http://www.kop pert.nl/) to prevent infestation by fungus gnats.

*Pieris rapae* L. (Small Cabbage White butterfly; Lepidoptera; Pieridae) were reared on Brussels sprouts plants (*B. oleracea* var. *gemmifera* cv Cyrus) in a growth chamber at  $21 \pm 1$  °C, 50–70 % relative humidity and a diurnal cycle of 16:8 L:D.

*Plutella xylostella* L. (Diamondback Moth; Lepidoptera; Yponomeutidae) were reared on Brussels sprouts plants (*B. oleracea* var. *gemmifera* cv Cyrus) in a growth chamber at  $22 \pm 1$  °C, 40–50% relative humidity and a diurnal cycle of 16:8 L:D.

The Western flower thrips (*Frankliniella occidentalis* (Pergande)) used in this study were originally collected from *Chrysanthemum* flowers and reared in glass bottles on green common bean pods (*Phaseolus vulgaris*) in a climate cabinet at  $25 \pm 1$  °C, 50–70% relative humidity and a diurnal cycle of 16:8 L:D.

*Myzus persicae* (Sulzer) (Green Peach Aphid; Hemiptera: Aphididae) were reared on radish plants, *Raphanus sativus* L., at  $19 \pm 2$  °C, 50–70% relative humidity and a diurnal cycle of 16:8 L:D.

The necrotrophic fungus *Botrytis cinerea*, strain B0510 (Van der Ent *et al.* 2008), was grown on half-strength PDA plates containing penicillin (100  $\mu$ g mL<sup>-1</sup>) and streptomycin (200  $\mu$ g mL<sup>-1</sup>) for 2 weeks at room temperature. Spores were collected and resuspended in half-strength potato dextrose broth (Difco Laboratories) to a final density of  $1.0 \times 10^5$  spores mL<sup>-1</sup>. After a 3-h incubation period, the spores were used for inoculation (Thomma *et al.* 1998; Pre *et al.* 2008; Van der Ent *et al.* 2008).

### Experimental design and treatments

We performed four different experiments in which plants were exposed to the following stresses (Fig. S4, Supporting information): experiment (1): drought, damage by *P. rapae* caterpillars alone or preceded by drought, or infestation by *B. cinerea*; experiment (2): damage by *P. xylostella* caterpillars; experiment (3): damage by the cell-content-feeding thrips *F. occidentalis* and experiment (4): infestation by the phloem-feeding aphid *M. persicae*.

In experiment (1), plant responses were quantified for all 308 accessions with six replicates per accession. Bioassays were performed in 10 temporal blocks. Each block included 28–33 random accessions. To correct for variation among temporal blocks, a fixed set of three accessions (CS28780; Tsu-0, CS76113; Col-0 and CS76129; Fei-0) was included in each temporal block. The spatial location of each plant was recorded. Within temporal blocks, plants were allocated in trays and the position of the tray in the rearing chamber recorded as its position in either of the six racks, each with four shelves. The position of each plant in the trays was also recorded in terms of X and Y coordinates. In each temporal block, accessions were exposed to the following five treatments with a total of six replicates per accession and treatment: (a) no stress, (b) drought stress, (c) *P. rapae* herbivory, (d) drought plus P. rapae herbivory or (e) B. cinerea infection plus P. rapae herbivory (Fig. S4A, Supporting information). Plants were grown under similar conditions during the first 3 weeks. Drought stress was imposed by withholding water for 7 days during the third week, while the rest of the plants was watered every 2 days with 1 L of water per tray. Withholding water for 7 days resulted in plants that displayed retarded growth and were smaller than well-watered plants. Botrytis cinerea inoculation was carried out 24 h prior to P. rapae inoculation. Plants were inoculated with B. cinerea by pipetting 5 µL of spores suspended in half-strength PDB (Difco Laboratories) at a concentration of  $1 \times 10^5$  spores mL<sup>-1</sup> onto two of the leaves in the rosette. To ensure successful infection by B. cinerea, plants were kept at 100% RH for 24 h. Four-week-old plants were exposed to herbivory by P. rapae as single or combined stress. Infestation with this species was carried out by placing two newly hatched first instar (L1) caterpillars on one of the leaves; they were allowed to feed for 5 days. To prevent caterpillars from moving between plants, individual plants were placed on the inverted lid of a Petri dish and trays were filled with water at the moment of inoculation. While plants were on the Petri dishes, they received the same watering regime as described above; however, watering was carried out by adding 20 mL of water to each Petri dish. Subsequently, plant rosette fresh weight was quantified for all treatments.

In experiment (2), all 308 accessions were used. However, some of the accessions displayed some germination problems so we did not have enough replicates for the experiment; this reduced the data set from 308 to 265 accessions. Four replicates were used for each of these accessions. The bioassays were performed in four temporal blocks. In each block, all accessions were phenotyped; one replicate per accession per temporal block, so block is the level of replication. Thus, the four temporal blocks resulted in four replicates per accession. Within blocks, accessions were randomly distributed over 40 trays. In this experiment, accession Col-0 was included to control for possible positional effects within the chamber. Each tray contained both control and treatment plants for Col-0 and the other accessions. Plant positions were randomized within the trays. Within each block, accessions were exposed simultaneously to either (a) no stress or (b) herbivory by *P. xy-lostella*. Plants were 4 weeks old when they were inoculated with two L2 larvae. Larvae were allowed to feed for 5 days. To prevent caterpillars from moving between plants, the same procedure was used as described above for experiment (1). Subsequently, plant rosette fresh weight was quantified for all treatments (Fig. S4B, Supporting information).

In experiment (3), bioassays were carried out for all 308 accessions with five replicates per accession. Bioassays were performed in five temporal blocks. Within each block, one replicate per accession was screened and so block is the level of replication. Leaves from 5-week-old plants were cut (one leaf per plant) and transferred to Petri dishes (diameter 5 cm; BD falcon, Product Number: 351006) containing a film of 1% agar. The petiole was inserted into the agar film (Fig. S4C, Supporting information). Leaves were exposed to three juvenile (second-larval instar, L2) F. occidentalis for 6 days (Fig. S4C, Supporting information). Feeding damage was estimated in mm<sup>2</sup> after 6 days by counting the number of small 'silver damage' feeding spots, where one small feeding spot accounts for 3 mm<sup>2</sup> damage (bigger spots were counted as 2–5 small spots).

In experiment (4), bioassays were carried out for all 308 accessions with two to three replicates per accession. The experiment was performed in three temporal blocks, each divided into four sub-blocks, representing four consecutive days. Three-week-old plants were inoculated with one 0–24 h-old *M. persicae* nymph per plant. Individual plants were placed on the inverted lid of a Petri dish in trays with a soap-diluted water barrier to prevent aphids from moving between plants. Each tray contained 16 plants. Fourteen days after infestation, the number of aphids per plant was counted (Fig. S4D, Supporting information).

The bioassay data are part of plant response data collected for a global population of 350 accessions. The data for this global population have been used previously to investigate the genetic architectures underlying single and combined stress responses (Davila Olivas *et al.* 2017; Thoen *et al.* 2017). Here, we used the stressresponse data of only the 308 European accessions to specifically investigate differences in stress responses between summer annuals and winter annuals.

#### Statistical analysis

*Genotypic means.* For each genotype, we estimated a genotypic mean response to each stress (Jimenez-Gomez *et al.* 2010; Filiault & Maloof 2012; Riedelsheimer *et al.* 

2012). These estimated means (or predicted values) were extracted from linear mixed effect models that were fitted with the ASReml package in R (Harrell 2009).

Experiment 1:  $Y = \mu + GEN + TRT + GEN \times TRT + random$ terms + e, Experiment 2:  $Y = \mu + GEN + TRT + GEN \times TRT + random$ terms + e, Experiment 3:  $Y = \mu + GEN + random$  terms + e, Experiment 4:  $Y = \mu + GEN + random$  terms + e.

In all these models, Y represents the measured variable in each experiment (i.e. A. thaliana rosette fresh weight, thrips silver damage or aphid reproduction), and e is the residual error. TRT is the treatment factor and GEN is genotype (accession) which were included as fixed factors. Random terms were included to correct for temporal and positional effects. Below, B refers to blocks, SB to sub-blocks, R to racks, S to shelves, T to trays, X to x-coordinate within tray and Y to y-coordinate within tray. The random part of experiment 1 consisted of random terms for B, R, S,  $B \times R \times S$ ,  $B \times R \times S \times T$ ,  $B \times R \times S \times T \times X$ , and  $B \times R \times S \times T \times Y$ . For experiment 2, it consisted of random terms for B, B  $\times$  T, B  $\times$  T  $\times$  X and B  $\times$  T  $\times$  Y, and for experiments 3 and 4, random terms for B and B/SB. In the first two experiments, predicted means from the fitted models were used to estimate variables that capture the response to each stress. The response to each stress consisted of A. thaliana fresh weight in the form of percentage reduction relative to control plants not exposed to stress. In the treatment where plants were exposed to both drought and herbivory by P. rapae, percentage weight reduction was relative to plants that experienced drought. From here onward, we will refer to the response to each stress as P. rapae, Drought, Drought&Pieris, Botrytis&Pieris and P. xylostella, respectively. Feeding damage by thrips will be referred to as F. occidentalis and number of aphids as M. persicae. These variables are summarized in Data S1 (Supporting information). Furthermore, these variables were used in downstream analyses as response variables in phylogenetic mixed models and to estimate heritability as explained in the section below.

*Phylogenetic mixed models, heritability estimations.* When comparing traits among species or among different populations of the same species, one needs to control for the nonindependence of data points (correlated residuals) due to shared ancestry. This may be achieved by

statistical methods such as phylogenetic mixed models or animal models (Hadfield 2010; Hadfield & Nakagawa 2010). A phylogenetic mixed model is a type of mixed effects model where, in addition to any other fixed or random effects, a pedigree representing the genealogy of the individuals is included in the model. This genealogy is then transformed into a variance-covariance matrix of relatedness between individuals and included in the model as a random additive genetic effect (Kruuk, 2004; Postma & Charmantier, 2007). In our study, such genealogy was based on the kinship matrix obtained from the 214 051 A. thaliana SNPs (Atwell et al. 2010) as is common in genetic association studies (Kim et al. 2007; Li et al. 2010; Bac-Molenaar et al. 2015). By doing so, we accounted for the nonindependence of plant phenotypic responses among genetically similar plant accessions. This is particularly relevant, for example when specific genotypes aggregate in a particular geographic location not because of the action of natural selection but because of genetic drift. To test hypotheses about the effect of plant life cycle strategy and geographic location on plant responses to stress, we built a different phylogenetic mixed model for each stress. Phenotypic responses of the plants were included as response variable with a Gaussian distribution. As predictors we included (i) plant life cycle strategy as a categorical variable (winter vs summer annual) and (ii) latitude, (iii) longitude and (iv) elevation as continuous variables (phylogenetic model type 1). The effect of environmental gradients (latitude, longitude and elevation) on A. thaliana plant life cycle strategy was also modelled with similar phylogenetic mixed models but with plant life cycle strategy included as a categorical response variable (phylogenetic model type 2). To explore potential trade-offs among plant responses to different stresses, a different phylogenetic mixed model was also built for each of the single stresses studied here. In each phylogenetic mixed model, we included plant responses to a particular stress as a response variable with a Gaussian distribution, and as predictors, we included responses to the other stresses and plant life cycle strategy (phylogenetic model type 3). All phylogenetic mixed models were fitted with Bayesian Markov chain Monte Carlo (MCMC) techniques implemented in the MCMCglmm package in R (Hadfield 2010; Hadfield & Nakagawa 2010). In the phylogenetic mixed models types 1 and 3, the MCMC chain ran for 150 000 iterations. To prevent autocorrelation among subsequent iterations, the chain was sampled every 50 iterations with the first 50 000 removed as burn-in. In the phylogenetic mixed model type 2, all MCMC parameters where ten times larger so that the chain ran for 1.5 million iterations. Autocorrelation between consecutive values was always lower than 0.1, and convergence of the chains

was confirmed by visual inspection so that there were no trends in the chain and posterior distributions were not skewed. Fixed effects are presented as the posterior mean (PM) with the credible intervals (CI) of the estimate, with significance reported as the pMCMC statistic (Hadfield 2010; Hadfield & Nakagawa 2010). In a Bayesian model, probability distributions need to be specified via specific priors for the fixed effects and the covariance matrices. Because we did not have any a priori knowledge on the distribution of our data, we used flat priors. For phylogenetic mixed models type 1 and 3, we used priors with scale = 0.002 and degree of belief = 0.002 (i.e. V = 1, nu = 0.002). For the phylogenetic mixed models type 2, we used an inverse Wishart prior with scale = I\*0.002 and degree of belief = 2.002 (i.e. V = diag(1)\*(0.002/2.002), nu = 2.002). Narrowsense heritability for each response was also estimated with the heritability package in R (Kruijer et al. 2015).

*Multi-Trait Mixed Model (MTMM) Genomewide Association analysis.* Two trait vectors were generated based on their response in summer vs. winter annuals (Fig. 2):

*p1*: Traits where summer annuals were more resistant than winter annuals (*P. rapae, P. xylostella,* Drought&*Pieris, Botrytis*&*Pieris*).

*p*2: Traits where winter annuals were more resistant than summer annuals (*F. occidentalis, M. persicae,* Drought).

The first principal component of each vector was used as a phenotype for the MTMM GWA analysis. MTMM was carried out as described by Korte et al. (2012), using 214k SNPs with a minor allele frequency >0.05. Briefly, a multiple trait mixed model was fitted for phenotypes of *n* individuals as  $Y = X\beta + G + E$ , with Y being the genotypes by traits  $(n \times p)$  matrix of phenotypic observations. The terms  $X\beta$ , G and E stand for, respectively, the fixed effects (including trait specific intercepts and SNP effects) and the random genetic and environmental effects. G follows a zero mean matrix-variate normal distribution with row-covariance (marker-based kinship) matrix K and column (trait) covariance matrix  $V_{g}$ .  $V_{g}$  is a  $p \times p$  matrix modelling the genetic correlations between traits. Similarly, vec(E) follows a zero mean normal distribution with covariance  $V_{\rm e} \otimes I_{\rm n\prime}$  where  $V_{\rm e}$  accounts for the nongenetic correlations between traits.

Using MTMM with multiple traits, we can find causal loci with (a) a common effect across traits, (b) trait-specific loci or (c) loci with opposite effects. For this purpose, MTMM uses a generalized least square (GLS) F test. For any two traits, it can be constructed as follows:

$$y = \begin{bmatrix} p1\\ p2 \end{bmatrix} = s_1\mu_1 + s_2\mu_2 + \mathbf{x}\beta + (\mathbf{x} \times \mathbf{s}_1) + \mathbf{\alpha} + \mathbf{v},$$

where *x* is the marker,  $s_1$  is a vector of 1 for all values belonging to the first trait and 0 otherwise, and  $s_2$  is a vector of 1 for all values belonging to the second trait and 0 otherwise. v (0,cov(y)) is a random variable that captures the error and the genetic random effect. MTMM applies the following tests:

(full); The full model tested against a null model where  $\beta = 0$  and  $\alpha = 0$ . This test identifies both loci with common and differing effects in one model but suffers in power from the extra degree of freedom.

(common); A model  $\alpha = 0$  tested against a null model where  $\beta = 0$  and  $\alpha = 0$ . This test identifies loci with common effect.

(specific); The full model tested against a model where  $\alpha = 0$ . This test identifies loci specific or with opposite effect.

*Candidate gene selection.* For selection of candidate genes, an arbitrary threshold was considered. Regions containing SNPs with  $-\log_{10}(P) \ge 4$  were considered for further analysis as described by El-Soda *et al.* (2015). A search window was defined by a 20-Kb region around significant SNPs. All genes within a search window were considered to be potential candidate genes. Genes were annotated according to TAIR 10. Relative gene expression data for *A. thaliana* (Col-0) plants exposed to jasmonic acid or salicylic acid were obtained from http://bar.utoronto.ca/.

### Results

### *Population structure of the European accessions of the Hapmap population*

We first examined the patterns of population structure of the European accessions in the Hapmap population. The first nine genetic principal components, explaining 12% of total variance, showed strong geographic autocorrelation and were used to subdivide the European accessions into ten geographically distinct genetic groups (Fig. 1).

### *Heritability of* A. thaliana *responses to abiotic and biotic stresses*

An important condition for natural selection to act upon a trait (and therefore to allow local adaptation) is that this trait has phenotypic variation which is genetically determined. This condition was met for *A. thaliana* responses to different stresses as the different traits that



Fig. 1 Population structure in the European accessions from the Hapmap population. Clusters of accessions were generated based on kinship matrix. (a) Geographic distribution of genetic clusters is shown in colours. (b) Dendrogram shows the relationship among accessions. Branch colours correspond to the colours in the map. Life cycle strategy is indicated by red circles (summer annuals) and white circles (winter annuals). Dendrogram was generated with itol (http://itol.embl.de/index.shtml). [Colour figure can be viewed at wileyonlinelibrary.com]

we quantified showed substantial heritable variation (Table 1). The largest trait variation was observed for the response to *P. xylostella* (CV = 138%), while the response to *M. persicae* displayed the lowest variation (CV = 20%). No relationship was observed between degree of trait variation and heritability. The largest heritability was observed for feeding damage by thrips (*F. occidentalis*) ( $h^2 = 0.90$ ), while plant biomass reduction after *P. xylostella* feeding had the lowest heritability ( $h^2 = 0.25$ ). The response to Drought&*Pieris* exhibited less variation and a lower heritability (CV = 33%,  $h^2 = 0.39$ ) than the response to *P. rapae* alone (CV = 43%,  $h^2 = 0.60$ ). The response to *Botrytis*&*Pieris* (CV = 117%,  $h^2 = 0.67$ ) had more variation and a higher heritability than the response to *P. rapae* alone.

# *Effect of life history strategy and genetic structure on stress responses in* A. thaliana

We classified *A. thaliana* accessions as winter and summer annuals based on their flowering time. Of the 308 accessions analysed, 89 did not produce flowers after 75 days and were therefore classified as winter annuals, while 219 behaved as summer annuals (Data S1, Supporting information). Within this collection of

genotypes, winter annuals were more often found in lower elevations and higher latitudes, whereas longitude was not a significant explanatory variable (Fig. S5, Supporting information; Table 2). Furthermore, the two life history strategies were not equally distributed among genetic groups. For example, cluster 8 and cluster 3 had mostly winter annuals while cluster 10 comprised summer annuals exclusively (Figs 1B, S5B, Supporting information). Phenotypic differences between genetic groups were tested by ANOVA after correction for life cycle strategy. Overall, stress responses did not differ significantly between genetic clusters, with the exception of the response to F. occi*dentalis* ( $P < 10^{-6}$ ) (Fig. S6, Supporting information).

A different phylogenetic mixed model was built for each stress factor, to explore whether plant responses varied between winter and summer annuals and among accessions located along geographic gradients. We found that life cycle strategy significantly affected all responses measured (Table 2; Fig. 2). Summer annuals were more resistant than winter annuals with regard to stresses that included caterpillars. The percentage of *A. thaliana* biomass reduction as a result of feeding by *P. rapae* and *P. xylostella* caterpillars was 6% and 10% larger in winter than in summer annuals, respectively.

Response trait	Experiment	Min.	Mean	Max.	CV (%)	Ν	$h^2$	va	ve
Drought*	1	-25.94	19.22	50.07	75	308	0.41	83.32	119.96
P. rapae*	1	-5.03	32.40	87.80	43	308	0.60	114.79	76.72
Drought&Pieris*	1	-3.39	48.25	143.80	33	308	0.39	96.15	153.52
Botrytis&Pieris*	1	-31.44	14.42	90.13	117	308	0.67	190.83	95.41
P. xylostella*	2	-24.86	14.15	78.16	138	265	0.25	94.08	287.25
F. occidentalis <sup>†</sup>	3	0.00	21.98	56.51	43	308	0.90	77.46	8.28
M. persicae <sup>‡</sup>	4	13.12	27.89	44.12	20	299	0.34	10.57	20.58

Table 1 Summary of trait values for A. thaliana resistance to abiotic and biotic stresses.

 $h^2$  = Narrow-sense heritability, va = additive genetic variance, ve = residual variance, CV = coefficient of variation, N = number of observations after subset for European accessions.

\*Trait values are expressed in %.

<sup>†</sup>Trait values are expressed in mm<sup>2</sup>.

<sup>‡</sup>Trait values are counts.

The percentage of biomass reduction by the combined stresses Drought&Pieris and Botrytis&Pieris was 8% and 12% larger in winter than summer annuals, respectively. Winter annuals were more resistant to drought stress and piercing-sucking insects than summer annuals. The percentage of biomass reduction in A. thaliana caused by drought was 10% lower in winter annuals than in summer annuals. Feeding damage by thrips (F. occidentalis) was 8% lower in winter annuals than in summer annuals, and the number of offspring produced by aphids (M. persicae) was 6% lower on winter annuals than on summer annuals. In all phylogenetic mixed models for plant responses to stress, and in accordance with our heritability estimates, the importance of A. thaliana genealogy was moderate, but relatively higher for the models on plant responses to thrips (F. occidentalis; Table 2).

# *Effect of environmental gradients on stress responses in* A. thaliana

We hypothesized that latitude, longitude and elevation of the locations from which the accessions had been collected would affect plant responses and, hence, these variables were also included in the phylogenetic mixed models as predictors. Although the three gradients were highly correlated with several environmental variables measured at each sampling location (Fig. S3, Supporting information), we found only a single significant association with A. thaliana responses to stress (Table 2). In particular, we found a significant, positive association between longitude and percentage of biomass reduction caused by drought (Table 2; Fig. S7, Supporting information). Furthermore, a longitudinal gradient was observed for flowering time and a latitudinal gradient was observed for flowering time after vernalization (Table S2, Fig. S7, Supporting information).

### Trade-offs in A. thaliana responses to stress

We built a separate phylogenetic mixed model for each of the single stresses and assessed whether they correlated with the other stresses. Because of its importance, plant life cycle strategy was also included as a cofactor in these models. Among the different stresses, we found a clear negative relationship between response to drought and to *P. rapae* herbivory (posterior mean = -0.12, *P* = 0.049), which indicates that these two responses trade-off (Fig. 3; Table 3). Responses to *P. rapae* herbivory were also negatively correlated with reproduction by the aphid *M. persicae* (posterior mean = -0.42, *P* = 0.005).

#### Multi-trait GWA analysis

As our models showed that life history strategy is a predictor for all of the tested stress responses, we explored which genes were associated with flowering-time-related resistance by genomewide association mapping. Two trait vectors were generated by taking the first principal component (PC) of stresses that displayed a resistance response in summer annuals (*P. rapae, P. xylostella,* Drought&*Pieris, Botrytis*&*Pieris,* PC1 = 55%) and the first PC of stresses with a resistance response in winter annuals (*F. occidentalis, M. persicae,* Drought, PC1 = 40%).

MTMM analysis identified a total of 23 trait-specific SNPs at  $-\log_{10}(P)$ -value  $\geq 4$ , which were distributed across 12 QTL regions, containing a total of 168 genes, in the genome (Fig. 4, Data S2, Supporting information). These 12 regions (QTLs) contained putative biotic stress-related genes, such as a pathogenesis-related thaumatin gene (*At5g40020*), and several life history strategy-related candidates. Two genes in QTL 6 (*At3g57390: AGL18* and *At3g57300:INO80*) are members of the flowering-time

	Life cycle strategy		Elevation		Latitude		Longitude		
Response trait	PM	Ρ	PM	Р	PM	Ρ	PM	Р	r1ant genealogy PM
Drought	-9.21 (-12.59 to -5.62)	<0.001	0.003 (-0.004 to 0.010)	0.389	-0.12 (-0.56 to 0.27)	0.566	0.229 (-0.004 to 0.426)	0.043	1.51 (<0.001-7.133)
P. rapae	4.61 (1.38–8.18)	0.007	-0.005(-0.012  to  0.001)	0.167	0.12 (-0.29 to 0.53)	0.575	-0.03 ( $-0.24$ to $0.19$ )	0.768	4.22 (<0.001-21.26)
Drought&Pieris	7.37 (3.10–11.13)	<0.001	-0.002 (-0.011  to  0.005)	0.534	0.23 (-0.24 to 0.68)	0.352	-0.09 ( $-0.32$ to $0.14$ )	0.432	2.34 (<0.001-11.81)
Botrytis&Pieris	10.68 (6.76 - 14.81)	<0.001	-0.004 ( $-0.013$ to $0.003$ )	0.329	0.25 (-0.23 to 0.73)	0.326	-0.11 (-0.38 to 0.12)	0.383	3.05 (< 0.001 - 14.65)
P. xylostella	10.40 (4.86–15.75)	<0.001	0.004 (-0.007  to  0.015)	0.444	-0.05 (-0.74  to  0.58)	0.867	-0.07 (-0.45  to  0.25)	0.683	5.49 (<0.001-28.58)
F. occidentalis	-3.50 (-6.00  to  -1.31)	0.004	-0.003 ( $-0.008$ to $0.001$ )	0.149	0.02 (-0.29 to 0.32)	0.893	0.16 (-0.03 to 0.35)	0.102	38.99 (0.029–99.53)
M. persicae	-1.75 ( $-3.20$ to $-0.19$ )	0.018	-0.001 ( $-0.003$ to $0.002$ )	0.629	-0.08 (-0.27 to 0.09)	0.357	0.008 (-0.118 to 0.123)	0.851	3.78 (<0.001-17.17)
Flowering type	NA	NA	-0.26 (-0.55  to  -0.01)	0.031	15.58 (4.21 to 29.39)	0.003	-0.971 ( $-6.61$ to $4.31$ )	0.732	0.01 (< 0.001 - 0.02)

Table 2 Bayesian phylogenetic mixed model analysis to assess differences in life cycle strategy and geographic gradients for each stress. For each variable, the posterior mean

gene network (Bouché *et al.* 2016). Interestingly, *ino80* mutants display a late-flowering phenotype, while the *agl18agl15* double mutant displays an early-flowering phenotype (Fernandez *et al.* 2014; Zhang *et al.* 2015). Furthermore, ELF4, a homologue of *At1g72630*: *ELF4-L2* in QTL 4, has also been implicated in flowering time, where the *elf4* single mutant displays a late-flowering phenotype (Doyle *et al.* 2002). Another interesting candidate is *At1g30060*:*COP1-interacting protein* in QTL 1 that has not been directly implicated in flowering time; however, *cop1* displays a late-flowering phenotype (McNellis *et al.* 1994). Even though their direct role in plant resistance remains to be investigated, these flowering-related candidate genes suggest that life-history strategy may affect plant responses to stress.

### Discussion

Plants are exposed to a wide range of biotic and abiotic stresses and need to deal with this ensemble of stresses in order to survive and reproduce. Yet, responses of plants to stress are usually investigated for individual stresses and for individual plant accessions. Plant accessions may differ in life history, which may decisively influence their exposure to a subset of the total set of environmental stresses and thus the selection pressures expressed. Here, we executed an extensive study to investigate whether life history affects the response of 308 European A. thaliana accessions to seven different stresses, including both single stresses and combinations of stresses. The most salient result of our study is that life history of Arabidopsis plants has a major effect on resistance to these stresses, rather than latitude, longitude or altitude. This is in line with a recent metaanalysis that shows that genetic variation in life history characteristics has a strong correlation with resistance to stress (Carmona et al. 2011). In particular, life history traits such as flowering time exhibited the strongest genetic correlations with herbivore susceptibility (Carmona et al. 2011). Our multitrait GWA analysis suggests the involvement of life history strategy-related genes in plant responses to different types of stresses. These genes are related to the timing of flowering and thus further indicate that genes influencing life history strategy may be involved in plant responses to stress. Because life history strategies such as being summer annual or winter annual have consequences for the plant's exposure to environmental stresses such as drought or herbivory, a link between the two traits may likely have been under positive selection. Which genes underlie the contrasting responses of summer annuals and winter annuals to stresses remains to be experimentally investigated. The fact that there are floweringrelated genes in the 12 QTLs identified should incite the

### LIFE HISTORY STRATEGY DETERMINES STRESS RESPONSES 2969



Fig. 2 Stress responses of A. thaliana accessions belonging to either winter annual or summer annual life cycle strategies. Accessions that required vernalization for flowering were classified winter annuals (blue, n = 89); the rest were classified summer annuals (orange, n = 219). Bars show mean value  $\pm$  SE. Stress responses to P. rapae caterpillars, P. xylostella caterpillars, Drought, Drought&Pieris and Botrytis&Pieris are represented by the percentage of biomass reduction in A. thaliana in response to each stress. Response to M. persicae is represented by the number of aphids produced. Response to F. occidentalis thrips is represented by the amount of feeding damage in mm<sup>2</sup>. Bayesian Pvalues are indicated as  $*P \le 0.05$ , \*\**P* < 0.01, \*\*\**P* < 0.001. [Colour figure can be viewed at wileyonlinelibrary.com]

search to their possible involvement. However, it should be realized that a QTL allele effect in an association analysis represents the conditional difference between two groups of genotypes with alternative versions of a SNP and epistatic interactions may play a role (Caicedo *et al.* 2004). Furthermore, our data show that summer annuals of *A. thaliana* are more resistant to stress involving caterpillars, either as single stress or as component of a combination of stresses, whereas winter annuals are more resistant to drought and herbivores such as aphids and thrips. The summer annual is considered the derived state (Michaels *et al.* 2003; Amasino 2004). The present study implies that the evolutionary shift from winter annual to summer annual imposes new selection pressures.

# *Geographic patterns in genetic diversity of European accessions*

The 308 European *A. thaliana* accessions displayed a strong isolation by distance (IBD) which implies that,

relative to distantly located accessions, accessions from the same location are genetically similar. Our findings are in line with several studies that have reported strong similar IBD patterns in A. thaliana accessions from Europe (Sharbel et al. 2000; Koornneef et al. 2004; Ostrowski et al. 2006; Schmid et al. 2006; Platt et al. 2010), although such isolation is weaker among the accessions introduced in North America (Platt et al. 2010). Many factors can contribute to isolation-by-distance patterns, but A. thaliana populations are suggested to become genetically isolated at a fast rate due to the self-pollinating nature of the species (Alonso-Blanco & Koornneef 2000). This trait is thought to be also important in explaining why A. thaliana populations are highly inbred with large genetic variation within and among populations (Nordborg et al. 2005; Horton et al. 2012). This kind of population structure has implications for our study, especially when inferring local adaptation. If similar genotypes are clustered in space, correlations between genotype and environment may arise due to spatial clustering and lead to misleading



Fig. 3 Trade-off between response to abiotic and biotic stresses among A. thaliana accessions. Accessions that required vernalization for flowering were classified winter annuals (blue); the rest were classified summer annuals (orange). Stress responses to P. rapae and Drought are represented by the percentage of biomass reduction in A. thaliana in response to each stress. Response to M. persicae is represented by the number of aphids produced. Lines represent a linear regression fit and shades the 95% confidence interval. These lines were just used for better appreciation of the data because correlations tests were carried out using a linear mixed model as described in Materials and Methods. Posterior means (PM) and (P) Bayesian P-values are indicated. [Colour figure can be viewed at wileyonlinelibrary.com]

interpretations (Platt et al. 2010). For example, Allard et al. (1972) provided a classical example of local adaptation in Avena barbata in California, but a deeper analysis revealed that these populations were not locally adapted but spatially structured (Latta & Gardner 2009). In our study, we corrected for these potentially misleading associations by implementing a nd Barrocian int adible O E O/ 4 niablo C ç - lobo Por incord ÷ nhvlo 5 D22 Table 3

	Life cycle strategy		Drought		P. rapae		P. xylostella		F. occidentalis		M. persicae	
Response trait	PM	Р	PM	Р	PM	Р	PM	Р	Md	Р	PM	Р
Drought	-6.70 (-10.77 to -2.86)	<0.001			-0.13 (-0.26	0.032	-0.08 (-0.16	0.075	0.14 (-0.03	0.133	0.09 (-0.21	0.551
P. rapae	2.26 (-2.09 to 6.13)	0.276	-0.12(-0.24)	0.049	(10.0- 0)		0.05 (-0.03	0.227	0.12 (-0.06	0.187		0.005
P. xylostella	7.7 (2.42–13.35)	0.003	-0.16(-0.32)	0.084	0.12 (-0.06	0.196	10 0.14/		-0.15(-0.40)	0.262	<pre>(c1.0 - 0) </pre>	0.988
F. occidentalis	-3.72 (-6.39 to -1.11)	0.011	0.05(-0.03)	0.198	0.06 (-0.02 0.06 (-0.02	0.153	-0.03(-0.08)	0.314	(01.0 Q)		-0.05(-0.25)	0.649
M. persicae	-1.09 (-2.93 to 0.43)	0.191	0.01 (-0.03 to 0.07)	0.524	-0.07 (-0.12 to -0.02)	0.002	<pre>&lt;0.001 (-0.04 </pre> <pre>to 0.04)</pre>	0.973	0.01 (-0.08 to 0.08)	0.809	(c1.0 M	

© 2017 The Authors. Molecular Ecology Published by John Wiley & Sons Ltd.



**Fig. 4** Multi-trait GWA analysis. Red line indicates an arbitrary threshold set at  $-\log 10(P) \ge 4$  as described in the Methods section for detecting significant associations. QTL regions are indicated with numbers. Detailed information about each QTL is presented in Data S2 (Supporting information). [Colour figure can be viewed at wileyonlinelibrary.com]

phylogenetic mixed model that accounted for the genetic resemblance among individuals and, therefore, their spatial colocation.

# Arabidopsis responses to abiotic and biotic stresses differ between life history strategies

Two life history strategies have been described in A. thaliana: winter and summer annuals (Pigliucci 1998; Koornneef et al. 2004; Shindo et al. 2007). The winter annual is considered the ancestral state because loss and reduced function alleles at two genes (FRI and FLC) confer the summer annual phenotype (Michaels et al. 2003; Amasino 2004). It is generally accepted that summer annuals occur closer to the equator while winter annuals occur in temperate regions (Johanson et al. 2000; Michaels et al. 2003). Despite this general assumption, no clear geographic patterns were observed for the occurrences of these two life strategies in an earlier study (Shindo et al. 2007). Here, we observed a higher proportion of summer annuals at high elevations and southern latitudes. In line with this observation, an increase of summer annuals with increasing altitude in Spain (400-1700 m) and the Swiss alps (600-2700 m) has been reported (Pico 2012; Luo et al. 2015). It is interesting that a decrease in grasshopper abundance is correlated with a decrease in resistance of plants to chewing insect herbivores (Descombes et al. 2017). This, together with data on the correlation of life history with resistance to herbivory (Carmona et al. 2011) further supports the connection of life history strategy and selection pressure by herbivores. The predominance of summer annuals at high elevations has been attributed to the extended and cold winters that A. thaliana may be able to withstand as seed but not as rosette. On the

other hand, several studies have reported a predominance of summer annuals at low altitudes along elevational gradients in the Iberian peninsula (100–1600 m; Montesinos-Navarro *et al.* 2011; Mendez-Vigo *et al.* 2013) that was explained by the extreme drought conditions in summer that may select for early flowering (Luo *et al.* 2015). Furthermore, we also observed genetic clusters consisting mostly of winter annuals at northern latitudes. This is also consistent with literature reports where nonrandom distribution of strong winter annuals has been observed in Scandinavia (Shindo *et al.* 2005).

Independent of the biogeographic distribution of life cycle strategies, our study revealed that life cycle strategy is the most important factor explaining plant responses to most of the stresses studied here, except for resistance to thrips. We found that the degree of damage to thrips (*F. occidentalis*) was strongly influenced by geographic-genetic structure. This was also reflected by the high heritability values estimated for thrips resistance. Interestingly, this geographic-genetic structure resembles the geographic-genetic distribution of glucosinolates, a well-established defence mechanism against generalist herbivores, including thrips (Zust *et al.* 2012; Brachi *et al.* 2015).

Several authors have suggested that at higher latitudes, herbivore pressure is lower and hence plants are less defended (Pennings & Silliman 2005; Salazar & Marquis 2012; Halbritter et al. 2013). Assuming that winter annuals are more common in these habitats, we expected that relative to summer annuals, winter annuals would be less defended against herbivory (Pennings & Silliman 2005; Salazar & Marquis 2012; Halbritter et al. 2013). Indeed, we found that winter annuals were more susceptible to damage by the two chewing specialist caterpillars *P. rapae* and *P. xylostella* than summer annuals. However, the opposite was observed for the two piercing-sucking generalist herbivores: the phloemfeeding aphid M. persicae and the cell-content-feeding thrips F. occidentalis. Whether this difference between the effects of life history on chewing and phloem-sucking herbivores is correlated with the differences in feeding style or phenology remains to be investigated. Although we only tested four different insect herbivore species, these results add to the growing body of literature that suggests that the degree of specialization (Mathur et al. 2011; Ali & Agrawal 2012; Barrett & Heil 2012) and insect feeding guild (De Vos et al. 2005; Bodenhausen & Reymond 2007; Bidart-Bouzat & Kliebenstein 2011) may exert different selective pressures on plants.

*Arabidopsis thaliana* accessions displaying different life cycle strategies cope with drought stress in a different manner. For instance, winter annuals employ drought

avoidance (i.e. mechanisms that maintain the internal water status under limited water conditions such as stomatal closure and increased root growth) and summer annuals employ drought escape (i.e. shift in phenology that allows plants to grow and reproduce by avoiding activity during periods of water scarcity; McKay et al. 2003; Des Marais et al. 2012; Juenger 2013; Easlon et al. 2014). Furthermore, a link has been observed between life cycle strategies and heat tolerance, such that late-flowering plants are more sensitive to heat stress (Bac-Molenaar et al. 2015). We, therefore, predicted that under our controlled experimental conditions, summer annuals would be less adapted to drought than winter annuals. Our hypothesis was confirmed. When exposed to drought stress, summer annuals gained less weight than winter annuals. A strong correlation between flowering life history and drought resistance has already been observed in the laboratory in A. thaliana (McKay et al. 2003; Juenger 2013; Easlon et al. 2014). In fact, winter annuals have higher wateruse efficiency than summer annuals (Juenger 2013; Lovell et al. 2013). The association between these two traits has been suggested to be partially caused by alleles of the FLC and FRI genes (McKay et al. 2003; Scarcelli et al. 2007). A recent study that exposed a large collection of A. thaliana accessions to controlled moderate drought found the opposite, that is that summer annuals were more resistant than winter annuals (Bac-Molenaar et al. 2016). The discrepancy between these studies may arise from different levels or time patterns of drought stress applied. For instance, it has been determined that severe drought and moderate drought elicit different physiological and molecular responses in A. thaliana (Skirycz et al. 2011). In addition, studies on other taxa also found evidence that flowering-time genes can have pleiotropic effects on other traits such as water-use efficiency in Brassica rapa (Franks 2011), vegetative biomass in A. barbata (Latta & Gardner 2009) and size at reproduction in B. rapa (Haselhorst et al. 2011).

Here, we tested the hypothesis that *A. thaliana* responses to stress would vary with elevation, latitude and longitude. The only significant correlation found was between longitude and *A. thaliana* responses to drought: drought resistance (i.e. reduced weight loss when experiencing the stress) increased eastwards. Interestingly, flowering time in European *A. thaliana* accessions has also been found to correlate with longitude, where the proportion of early-flowering accessions increases eastwards (Samis *et al.* 2008). In the present study, we also observed a longitudinal gradient for flowering time decreasing eastwards. Interestingly, experimental evolution research on several plant species demonstrated that wet soil and late-season

drought conditions selected for early-flowering accessions which displayed low water-use efficiency. On the other hand, early-season drought selected for higher water-use efficiency (Heschel & Riginos 2005; Sherrard & Maherali 2006). Large differences in physiological and transcriptional changes upon drought stress have been reported between winter and summer annuals in *A. thaliana* (Des Marais *et al.* 2012), further underlining the importance of life history in responses to stress. Future efforts should be devoted to understanding at the mechanistic level how accessions displaying distinct life cycle strategies cope with abiotic and biotic stresses.

### Plant responses against different stresses exhibit tradeoffs

Upon stress, plants are able to elicit defence or resistance mechanisms that are specific for the attacker or adverse abiotic condition (Reymond et al. 2000; De Vos et al. 2005; Verhage et al. 2011; Appel et al. 2014; Stam et al. 2014). However, plant responses to a specific stress are usually modulated at the individual or population level by cooccurrence with other stresses, because plants have limited resources and are locally exposed to different selection forces. Thus, the differential responses of summer and winter annuals to different stress types are likely to reflect differential selection pressures experienced by plants exhibiting these two life history strategies. In addition, life history theory predicts that the defensive repertoire is also genetically constrained so that adaptation to different stresses can trade-off. For example, abiotic stresses tend to have a negative impact on how plants deal with pathogens or herbivores (Suzuki et al. 2014; Ramegowda & Senthil-Kumar 2015). We hypothesized that the different A. thaliana accessions would be limited in their responses to stresses so that responses would be negatively correlated. A negative correlation was observed between responses to P. rapae herbivory and drought stress or aphid reproduction on the plants. Trade-offs are expected when the defences are specifically tailored and are costly to the plant (Erb et al. 2011). This suggests that responses to drought and *P. rapae* are specific and costly. An example of specific defences that constrain each other is the interaction between jasmonic acid (JA) and salicylic acid (SA) signalling pathways (Sendon et al. 2011; Van der Does et al. 2013; Caarls et al. 2015). Caterpillar feeding and drought induce an overlapping response at the transcriptional level (Reymond et al. 2000; Bodenhausen & Reymond 2007; Verhage et al. 2011; Vos et al. 2013a), vet with clear specific aspects as well (Coolen et al. 2016). At the metabolic level, clear differences can be recorded (Weldegergis et al. 2015). Recently, it has been shown that JA signalling is required to increase ABA levels under water stress conditions (De Ollas *et al.* 2015) and that ABA is involved in plant defence against caterpillar feeding (Vos *et al.* 2013b). A plausible explanation for the observed trade-off between resistance to drought and feeding by the specialist herbivore *P. rapae* may be resource allocation. Furthermore, extensive down-regulation under drought stress has been observed for genes that are upregulated by *P. rapae* herbivory, suggesting a mechanistic explanation for this trade-off (Coolen *et al.* 2016).

#### Conclusion

We exposed 308 well-genotyped European *A. thaliana* accessions to diverse biotic and abiotic and combined stresses. We found that plant life cycle strategy is a major determinant of responses to the different stresses that we applied in this study. Moreover, an extensive trade-off between *A. thaliana*'s response to *P. rapae* herbivory and drought and between the responses to *P. rapae* and aphids was observed. Several putative genes, including several life history strategy-related candidates, resulted from an MTMM GWA analysis. Future studies should aim to understand the mechanisms of how accessions with different life history strategies deal with different stresses.

### Acknowledgments

We are grateful to Mark G. M. Aarts and Kim M. C. A. Vermeer for constructive comments on a previous version of the manuscript. This work was supported by The Netherlands Organization for Scientific Research (NWO) through the Technology Foundation, Perspective Programme 'Learning from Nature' [STW10988]. E.F. was funded by Marie Curie Intra-European Fellowship FP7-PEOPLE-2012-IEF#329648.

#### References

- Alberto FJ, Aitken SN, Alia R *et al.* (2013) Potential for evolutionary responses to climate change evidence from tree populations. *Global Change Biology*, **19**, 1645–1661.
- Ali JG, Agrawal AA (2012) Specialist versus generalist insect herbivores and plant defense. *Trends in Plant Science*, **17**, 293–302.
- Allard RW, Babbel GR, Kahler AL, Clegg MT (1972) Evidence for coadaptation in Avena barbata. Proceedings of the National Academy of Sciences of the United States of America, 69, 3043– 3048.
- Alonso-Blanco C, Koornneef M (2000) Naturally occurring variation in *Arabidopsis*: an underexploited resource for plant genetics. *Trends in Plant Science*, **5**, 22–29.
- Amasino R (2004) Vernalization, competence, and the epigenetic memory of winter. *Plant Cell*, **16**, 2553–2559.
- Andrew NR, Hughes L (2005) Herbivore damage along a latitudinal gradient: relative impacts of different feeding guilds. *Oikos*, **108**, 176–182.

- Appel HM, Fescemyer H, Ehlting J *et al.* (2014) Transcriptional responses of *Arabidopsis thaliana* to chewing and sucking insect herbivores. *Frontiers in Plant Science*, **5**, 565.
- Atkinson NJ, Urwin PE (2012) The interaction of plant biotic and abiotic stresses: from genes to the field. *Journal of Experimental Botany*, 63, 3523–3543.
- Atwell S, Huang YS, Vilhjalmsson BJ et al. (2010) Genome-wide association study of 107 phenotypes in *Arabidopsis thaliana* inbred lines. *Nature*, **465**, 627–631.
- Bac-Molenaar JA, Fradin EF, Becker FFM *et al.* (2015) Genomewide association mapping of fertility reduction upon heat stress reveals developmental stage-specific QTLs in *Arabidop*sis thaliana. The Plant Cell, 27, 1857–1874.
- Bac-Molenaar JA, Granier C, Keurentjes JJB, Vreugdenhil D (2016) Genome wide association mapping of time-dependent growth responses to moderate drought stress in *Arabidopsis*. *Plant, Cell & Environment*, **39**, 88–102.
- Barrett LG, Heil M (2012) Unifying concepts and mechanisms in the specificity of plant-enemy interactions. *Trends in Plant Science*, **17**, 282–292.
- Baxter I, Brazelton JN, Yu DN *et al.* (2010) A coastal cline in sodium accumulation in *Arabidopsis thaliana* is driven by natural variation of the sodium transporter AtHKT1;1. *PLoS Genetics*, **6**, e1001193.
- Bidart-Bouzat MG, Kliebenstein D (2011) An ecological genomic approach challenging the paradigm of differential plant responses to specialist versus generalist insect herbivores. *Oecologia*, **167**, 677–689.
- Bodenhausen N, Reymond P (2007) Signaling pathways controlling induced resistance to insect herbivores in *Arabidopsis*. *Molecular Plant-Microbe Interactions*, **20**, 1406–1420.
- Bouché F, Lobet G, Tocquin P, Périlleux C (2016) FLOR-ID: an interactive database of flowering-time gene networks in Arabidopsis thaliana. Nucleic Acids Research, 44(Database), D1167– D1171.
- Brachi B, Meyer CG, Villoutreix R et al. (2015) Co-selected genes determine adaptive variation in herbivore resistance throughout the native range of Arabidopsis thaliana. Proceedings of the National Academy of Sciences of the United States of America, 112, 4032–4037.
- Bradshaw WE, Holzapfel CM (2001) Genetic shift in photoperiodic response correlated with global warming. *Proceedings of the National Academy of Sciences of the United States of America*, 98, 14509–14511.
- Broekgaarden C, Poelman EH, Steenhuis G et al. (2007) Genotypic variation in genome-wide transcription profiles induced by insect feeding: *Brassica oleracea-Pieris rapae* interactions. *BMC Genomics*, 8, 239.
- Caarls L, Pieterse CMJ, Van Wees SCM (2015) How salicylic acid takes transcriptional control over jasmonic acid signaling. *Frontiers in Plant Science*, 6, 170.
- Caicedo AL, Stinchcombe JR, Olsen KM, Schmitt J, Purugganan MD (2004) Epistatic interaction between Arabidopsis FRI and FLC flowering time genes generates a latitudinal cline in a life history trait. *Proceedings of the National Academy of Sciences of the United States of America*, **101**, 15670–15675.
- Carmona D, Lajeunesse MJ, Johnson MTJ (2011) Plant traits that predict resistance to herbivores. *Functional Ecology*, **25**, 358–367.
- Chao DY, Silva A, Baxter I et al. (2012) Genome-wide association studies identify heavy metal ATPase3 as the primary

© 2017 The Authors. Molecular Ecology Published by John Wiley & Sons Ltd.

determinant of natural variation in leaf cadmium in Arabidopsis thaliana. PLoS Genetics, 8, e1002923.

- Coolen S, Davila Olivas NH, Huang P *et al.* (2016) Whole transcriptome analysis of *Arabidopsis thaliana* under sequential biotic and abiotic stresses. *The Plant Journal*, **86**, 249–267.
- Davila Olivas NH, Kruijer W, Gort G et al. (2017) Genomewide association analysis reveals distinct genetic architectures for single and combined stress responses in Arabidopsis thaliana. New Phytologist, 213, 838–851.
- De Ollas C, Arbona V, Gómez-Cadenas A (2015) Jasmonoyl isoleucine accumulation is needed for abscisic acid build-up in roots of *Arabidopsis* under water stress conditions. *Plant*, *Cell and Environment*, **38**, 2157–2170.
- De Vos M, Van Oosten VR, Van Poecke RMP et al. (2005) Signal signature and transcriptome changes of Arabidopsis during pathogen and insect attack. Molecular Plant-Microbe Interactions, 18, 923–937.
- Debieu M, Tang C, Stich B *et al.* (2013) Co-variation between seed dormancy, growth rate and flowering time changes with latitude in *Arabidopsis thaliana*. *PLoS ONE*, **8**, e61075.
- Des Marais DL, McKay JK, Richards JH et al. (2012) Physiological genomics of response to soil drying in diverse Arabidopsis accessions. The Plant Cell, 24, 893–914.
- Descombes P, Marchon J, Pradervand JN et al. (2017) Community-level plant palatability increases with elevation as insect herbivore abundance declines. *Journal of Ecology*, 105, 142–151.
- Devictor V, van Swaay C, Brereton T *et al.* (2012) Differences in the climatic debts of birds and butterflies at a continental scale. *Nature Climate Change*, **2**, 121–124.
- Doyle MR, Davis SJ, Bastow RM et al. (2002) The ELF4 gene controls circadian rhythms and flowering time in Arabidopsis thaliana. Nature, 419, 74–77.
- Easlon HM, Nemali KS, Richards JH *et al.* (2014) The physiological basis for genetic variation in water use efficiency and carbon isotope composition in *Arabidopsis thaliana*. *Photosynthesis Research*, **119**, 119–129.
- El-Soda M, Kruijer W, Malosetti M, Koornneef M, Aarts MGM (2015) Quantitative trait loci and candidate genes underlying genotype by environment interaction in the response of *Arabidopsis thaliana* to drought. *Plant Cell and Environment*, **38**, 585–599.
- Erb M, Balmer D, De Lange ES *et al.* (2011) Synergies and trade-offs between insect and pathogen resistance in maize leaves and roots. *Plant Cell and Environment*, **34**, 1088–1103.
- Fernandez DE, Wang CT, Zheng Y et al. (2014) The MADSdomain factors AGAMOUS-LIKE15 and AGAMOUS-LIKE18, along with SHORT VEGETATIVE PHASE and AGAMOUS-LIKE24, are necessary to block floral gene expression during the vegetative phase. *Plant Physiology*, **165**, 1591–1603.
- Filiault DL, Maloof JN (2012) A genome-wide association study identifies variants underlying the *Arabidopsis thaliana* shade avoidance response. *PLoS Genetics*, 8, e1002589.
- Fournier-Level A, Korte A, Cooper MD et al. (2011) A map of local adaptation in Arabidopsis thaliana. Science, 334, 86–89.
- Franks SJ (2011) Plasticity and evolution in drought avoidance and escape in the annual plant *Brassica rapa*. New Phytologist, 190, 249–257.
- Fukami T, Wardle DA (2005) Long-term ecological dynamics: reciprocal insights from natural and anthropogenic gradients.

Proceedings of the Royal Society B-Biological Sciences, 272, 2105–2115.

- Gomaa NH, Montesinos-Navarro A, Alonso-Blanco C, Pico FX (2011) Temporal variation in genetic diversity and effective population size of Mediterranean and subalpine *Arabidopsis thaliana* populations. *Molecular Ecology*, **20**, 3540–3554.
- Hadfield JD (2010) MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *Journal of Statistical Software*, **33**, 1–22.
- Hadfield JD, Nakagawa S (2010) General quantitative genetic methods for comparative biology: phylogenies, taxonomies and multi-trait models for continuous and categorical characters. *Journal of Evolutionary Biology*, 23, 494–508.
- Halbritter AH, Alexander JM, Edwards PJ, Billeter R (2013) How comparable are species distributions along elevational and latitudinal climate gradients? *Global Ecology and Biogeography*, **22**, 1228–1237.
- Hancock AM, Brachi B, Faure N *et al.* (2011) Adaptation to climate across the *Arabidopsis thaliana* genome. *Science*, **334**, 83– 86.
- Harrell FE Jr (2009) Hmisc: Harrell Miscellaneous. R Package Version 3.0. R Foundation for Statistical Computing, Vienna, Austria. http://cran.r-project.org/web/packages/Hmisc/ index.html
- Haselhorst MSH, Edwards CE, Rubin MJ, Weinig C (2011) Genetic architecture of life history traits and environmentspecific trade-offs. *Molecular Ecology*, 20, 4042–4058.
- van Heerwaarden J, Doebley J, Briggs WH *et al.* (2011) Genetic signals of origin, spread, and introgression in a large sample of maize landraces. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, 1088–1092.
- van Heerwaarden J, Odong TL, van Eeuwijk FA (2013) Maximizing genetic differentiation in core collections by PCAbased clustering of molecular marker data. *Theoretical and Applied Genetics*, **126**, 763–772.
- Hereford J (2009) A quantitative survey of local adaptation and fitness trade-offs. *American Naturalist*, **173**, 579–588.
- Heschel MS, Riginos C (2005) Mechanisms of selection for drought stress tolerance and avoidance in *Impatiens capensis* (Balsaminacea). *American Journal of Botany*, **92**, 37–44.
- Horton MW, Hancock AM, Huang YS *et al.* (2012) Genomewide patterns of genetic variation in worldwide *Arabidopsis thaliana* accessions from the RegMap panel. *Nature Genetics*, 44, 212–216.
- Hughes L (2000) Biological consequences of global warming: is the signal already apparent? *Trends in Ecology & Evolution*, 15, 56–61.
- Jimenez-Gomez JM, Wallace AD, Maloof JN (2010) Network analysis identifies *ELF3* as a QTL for the shade avoidance response in *Arabidopsis*. *PLoS Genetics*, 6, e1001100.
- Johanson U, West J, Lister C et al. (2000) Molecular analysis of FRIGIDA, a major determinant of natural variation in Arabidopsis flowering time. Science, 290, 344–347.
- Juenger TE (2013) Natural variation and genetic constraints on drought tolerance. *Current Opinion in Plant Biology*, 16, 274– 281.
- Kaltz O, Shykoff JA (1998) Local adaptation in host-parasite systems. *Heredity*, 81, 361–370.
- Kawecki TJ, Ebert D (2004) Conceptual issues in local adaptation. *Ecology Letters*, 7, 1225–1241.

- Kim S, Plagnol V, Hu TT et al. (2007) Recombination and linkage disequilibrium in Arabidopsis thaliana. Nature Genetics, 39, 1151–1155.
- Kissoudis C, van de Wiel C, Visser RGF, van der Linden G (2014) Enhancing crop resilience to combined abiotic and biotic stress through the dissection of physiological and molecular crosstalk. *Frontiers in Plant Science*, **5**, 207.
- Koornneef M, Alonso-Blanco C, Vreugdenhil D (2004) Naturally occurring genetic variation in *Arabidopsis thaliana*. *Annual Review of Plant Biology*, **55**, 141–172.
- Korte A, Vilhjalmsson BJ, Segura V, Platt A, Long Q, Nordborg M (2012) A mixed-model approach for genome-wide association studies of correlated traits in structured populations. *Nature Genetics*, 44, 1066–1071.
- Kruijer W, Boer MP, Malosetti M *et al.* (2015) Marker-based estimation of heritability in immortal populations. *Genetics*, **199**, 379–393.
- Kruuk LEB (2004) Estimating genetic parameters in natural populations using the 'animal model'. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 359, 873–890.
- Latta RG, Gardner KM (2009) Natural selection on pleiotropic quantitative trait loci affecting a life-history trade-off in *Avena barbata*. *Evolution*, **63**, 2153–2163.
- Leimu R, Fischer M (2008) A meta-analysis of local adaptation in plants. *PLoS ONE*, **3**, e4010.
- Leinonen PH, Sandring S, Quilot B et al. (2009) Local adaptation in European populations of Arabidopsis lyrata (Brassicaceae). American Journal of Botany, **96**, 1129–1137.
- Leinonen PH, Remington DL, Savolainen O (2011) Local adaptation, phenotypic differentiation, and hybrid fitness in diverged natural populations of *Arabidopsis lyrata*. *Evolution*, 65, 90–107.
- Li Y, Huang Y, Bergelson J, Nordborg M, Borevitz JO (2010) Association mapping of local climate-sensitive quantitative trait loci in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 21199– 21204.
- Lovell JT, Juenger TE, Michaels SD *et al.* (2013) Pleiotropy of FRIGIDA enhances the potential for multivariate adaptation. *Proceedings of the Royal Society B-Biological Sciences*, **280**, 20131043.
- Luo Y, Widmer A, Karrenberg S (2015) The roles of genetic drift and natural selection in quantitative trait divergence along an altitudinal gradient in *Arabidopsis thaliana*. *Heredity*, 114, 220–228.
- Manel S, Schwartz MK, Luikart G, Taberlet P (2003) Landscape genetics: combining landscape ecology and population genetics. *Trends in Ecology & Evolution*, **18**, 189–197.
- Manzano-Piedras E, Marcer A, Alonso-Blanco C, Pico FX (2014) Deciphering the adjustment between environment and life history in annuals: lessons from a geographically-explicit approach in *Arabidopsis thaliana*. *PLoS ONE*, **9**, e87836.
- Mathur V, Ganta S, Raaijmakers CE *et al.* (2011) Temporal dynamics of herbivore-induced responses in *Brassica juncea* and their effect on generalist and specialist herbivores. *Entomologia Experimentalis et Applicata*, **139**, 215–225.
- McKay JK, Richards JH, Mitchell-Olds T (2003) Genetics of drought adaptation in *Arabidopsis thaliana*: I. Pleiotropy contributes to genetic correlations among ecological traits. *Molecular Ecology*, **12**, 1137–1151.

- McNellis TW, von Arnim AG, Deng XW (1994) Overexpression of Arabidopsis COP1 results in partial suppression of lightmediated development: evidence for a light-inactivable repressor of photomorphogenesis. The Plant Cell, 6, 1391–1400.
- Mendez-Vigo B, Gomaa NH, Alonso-Blanco C, Xavier Pico F (2013) Among- and within-population variation in flowering time of Iberian *Arabidopsis thaliana* estimated in field and glasshouse conditions. *New Phytologist*, **197**, 1332–1343.
- Michaels SD, He YH, Scortecci KC, Amasino RM (2003) Attenuation of FLOWERING LOCUS C activity as a mechanism for the evolution of summer-annual flowering behavior in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America*, **100**, 10102–10107.
- Mikola J (1982) Bud-set phenology as an indicator of climatic adaptation of Scots pine in Finland (*Pinus sylvestris*). Silva Fennica, **16**, 178–184.
- Moles AT, Bonser SP, Poore AGB, Wallis IR, Foley WJ (2011) Assessing the evidence for latitudinal gradients in plant defence and herbivory. *Functional Ecology*, **25**, 380–388.
- Montesinos-Navarro A, Wig J, Pico FX, Tonsor SJ (2011) Arabidopsis thaliana populations show clinal variation in a climatic gradient associated with altitude. New Phytologist, 189, 282–294.
- Montesinos-Navarro A, Pico FX, Tonsor SJ (2012) Clinal variation in seed traits influencing life cycle timing in *Arabidopsis thaliana*. *Evolution*, **66**, 3417–3431.
- Nordborg M, Hu TT, Ishino Y et al. (2005) The pattern of polymorphism in Arabidopsis thaliana. Plos Biology, 3, 1289–1299.
- Odong TL, van Heerwaarden J, van Hintum TJL, van Eeuwijk FA, Jansen J (2013) Improving Hierarchical clustering of genotypic data via principal component analysis. *Crop Science*, **53**, 1546–1554.
- Olsson K, Agren J (2002) Latitudinal population differentiation in phenology, life history and flower morphology in the perennial herb *Lythrum salicaria*. *Journal of Evolutionary Biology*, **15**, 983–996.
- Ostrowski MF, David J, Santoni S *et al.* (2006) Evidence for a large-scale population structure among accessions of *Arabidopsis thaliana*: possible causes and consequences for the distribution of linkage disequilibrium. *Molecular Ecology*, **15**, 1507–1517.
- Paccard A, Fruleux A, Willi Y (2014) Latitudinal trait variation and responses to drought in *Arabidopsis lyrata*. Oecologia, 175, 577–587.
- Pellissier L, Roger A, Bilat J, Rasmann S (2014) High elevation *Plantago lanceolata* plants are less resistant to herbivory than their low elevation conspecifics: is it just temperature? *Ecography*, **37**, 950–959.
- Pennings SC, Silliman BR (2005) Linking biogeography and community ecology: latitudinal variation in plant-herbivore interaction strength. *Ecology*, 86, 2310–2319.
- Pico FX (2012) Demographic fate of *Arabidopsis thaliana* cohorts of autumn- and spring-germinated plants along an altitudinal gradient. *Journal of Ecology*, **100**, 1009–1018.
- Pieterse CMJ, Leon-Reyes A, Van der Ent S, Van Wees SCM (2009) Networking by small-molecule hormones in plant immunity. *Nature Chemical Biology*, **5**, 308–316.
- Pigliucci M (1998) Ecological and evolutionary genetics of *Arabidopsis*. *Trends in Plant Science*, **3**, 485–489.
- Platt A, Horton M, Huang YS *et al.* (2010) The scale of population structure in *Arabidopsis thaliana*. *PLoS Genetics*, **6**, e1000843.

- Postma E, Charmantier A (2007) What 'animal models' can and cannot tell ornithologists about the genetics of wild populations. *Journal of Ornithology*, **148**, S633–S642.
- Pre M, Atallah M, Champion A *et al.* (2008) The AP2/ERF domain transcription factor ORA59 integrates jasmonic acid and ethylene signals in plant defense. *Plant Physiology*, 147, 1347–1357.
- Ramegowda V, Senthil-Kumar M (2015) The interactive effects of simultaneous biotic and abiotic stresses on plants: mechanistic understanding from drought and pathogen combination. *Journal of Plant Physiology*, **176**, 47–54.
- Rasmann S, Pellissier L, Defossez E, Jactel H, Kunstler G (2014) Climate-driven change in plant-insect interactions along elevation gradients. *Functional Ecology*, 28, 46–54.
- Reymond P, Weber H, Damond M, Farmer EE (2000) Differential gene expression in response to mechanical wounding and insect feeding in *Arabidopsis. Plant Cell*, **12**, 707–719.
- Riedelsheimer C, Lisec J, Czedik-Eysenberg A et al. (2012) Genome-wide association mapping of leaf metabolic profiles for dissecting complex traits in maize. Proceedings of the National Academy of Sciences of the United States of America, 109, 8872– 8877.
- Salazar D, Marquis RJ (2012) Herbivore pressure increases toward the equator. Proceedings of the National Academy of Sciences of the United States of America, 109, 12616–12620.
- Samis KE, Heath KD, Stinchcombe JR (2008) Discordant longitudinal clines in flowering time and PHYTOCHROME C in Arabidopsis thaliana. Evolution, 62, 2971–2983.
- Sanders NJ, Moss J, Wagner D (2003) Patterns of ant species richness along elevational gradients in an arid ecosystem. *Global Ecology and Biogeography*, **12**, 93–102.
- Savolainen O, Pyhajarvi T, Knurr T (2007) Gene flow and local adaptation in trees. Annual Review of Ecology Evolution and Systematics, 38, 595–619.
- Savolainen O, Lascoux M, Merila J (2013) Ecological genomics of local adaptation. *Nature Reviews Genetics*, **14**, 807–820.
- Scarcelli N, Cheverud JM, Schaal BA, Kover PX (2007) Antagonistic pleiotropic effects reduce the potential adaptive value of the FRIGIDA locus. Proceedings of the National Academy of Sciences of the United States of America, 104, 16986–16991.
- Schmid K, Torjek O, Meyer R et al. (2006) Evidence for a largescale population structure of Arabidopsis thaliana from genome-wide single nucleotide polymorphism markers. Theoretical and Applied Genetics, **112**, 1104–1114.
- Sendon PM, Seo HS, Song JT (2011) Salicylic acid signaling: biosynthesis, metabolism, and crosstalk with jasmonic acid. *Journal of the Korean Society for Applied Biological Chemistry*, 54, 501–506.
- Sharbel TF, Haubold B, Mitchell-Olds T (2000) Genetic isolation by distance in *Arabidopsis thaliana*: biogeography and postglacial colonization of Europe. *Molecular Ecology*, 9, 2109– 2118.
- Sheffield J, Wood EF, Roderick ML (2012) Little change in global drought over the past 60 years. *Nature*, **491**, 435–+.
- Sherrard ME, Maherali H (2006) The adaptive significance of drought escape in Avena barbata, an annual grass. Evolution, 60, 2478–2489.
- Shindo C, Aranzana MJ, Lister C *et al.* (2005) Role of FRIGIDA and FLOWERING LOCUS C in determining variation in flowering time of *Arabidopsis. Plant Physiology*, **138**, 1163– 1173.

- Shindo C, Bernasconi G, Hardtke CS (2007) Natural genetic variation in *Arabidopsis*: tools, traits and prospects for evolutionary ecology. *Annals of Botany*, 99, 1043–1054.
- Skirycz A, Vandenbroucke K, Clauw P *et al.* (2011) Survival and growth of *Arabidopsis* plants given limited water are not equal. *Nature Biotechnology*, **29**, 212–214.
- Stam JM, Kroes A, Li YH et al. (2014) Plant interactions with multiple insect herbivores: from community to genes. Annual Review of Plant Biology, 65, 689–713.
- Sternberg ED, Thomas MB (2014) Local adaptation to temperature and the implications for vector-borne diseases. *Trends in Parasitology*, **30**, 115–122.
- Stinchcombe JR, Weinig C, Ungerer M et al. (2004) A latitudinal cline in flowering time in Arabidopsis thaliana modulated by the flowering time gene FRIGIDA. Proceedings of the National Academy of Sciences of the United States of America, 101, 4712–4717.
- Sundqvist MK, Sanders NJ, Wardle DA (2013) Community and ecosystem responses to elevational gradients: processes, mechanisms, and insights for global change. *Annual Review* of Ecology, Evolution, and Systematics, 44, 261–280.
- Suzuki N, Rivero RM, Shulaev V, Blumwald E, Mittler R (2014) Abiotic and biotic stress combinations. *New Phytologist*, 203, 32–43.
- Thoen MPM, Davila Olivas NH, Kloth KJ et al. (2017) Genetic architecture of plant stress resistance: multi-trait genome-wide association mapping. *New Phytologist*, 213, 1346–1362.
- Thomma B, Eggermont K, Penninckx I *et al.* (1998) Separate jasmonate-dependent and salicylate-dependent defenseresponse pathways in *Arabidopsis* are essential for resistance to distinct microbial pathogens. *Proceedings of the National Academy of Sciences of the United States of America*, **95**, 15107– 15111.
- Van der Does D, Leon-Reyes A, Koornneef A et al. (2013) Salicylic acid suppresses jasmonic acid signaling downstream of SCFCOI1-JAZ by targeting GCC promoter motifs via transcription factor ORA59. Plant Cell, 25, 744–761.
- Van der Ent S, Verhagen BWM, Van Doorn R et al. (2008) MYB72 is required in early signaling steps of rhizobacteriainduced systemic resistance in *Arabidopsis*. *Plant Physiology*, 146, 1293–1304.
- Verhage A, Vlaardingerbroek I, Raaymakers C et al. (2011) Rewiring of the jasmonate signaling pathway in Arabidopsis during insect herbivory. Frontiers in Plant Science, 2, 47.
- Vos IA, Pieterse CMJ, van Wees SCM (2013a) Costs and benefits of hormone-regulated plant defences. *Plant Pathology*, 62, 43–55.
- Vos IA, Verhage A, Schuurink RC *et al.* (2013b) Onset of herbivore-induced resistance in systemic tissue primed for jasmonate-dependent defenses is activated by abscisic acid. *Frontiers in Plant Science*, **4**, 539.
- Walker LR, Wardle DA, Bardgett RD, Clarkson BD (2010) The use of chronosequences in studies of ecological succession and soil development. *Journal of Ecology*, 98, 725–736.
- Weigel D (2012) Natural variation in *Arabidopsis*: from molecular genetics to ecological genomics. *Plant Physiology*, **158**, 2–22.
- Weldegergis BT, Zhu F, Poelman EH, Dicke M (2015) Drought stress affects plant metabolites and herbivore preference but not host location by its parasitoids. *Oecologia*, **177**, 701–713.

- Wheeler T, von Braun J (2013) Climate change impacts on global food security. *Science*, 341, 508–513.
- Zhang C, Cao L, Rong L *et al.* (2015) The chromatin-remodeling factor AtINO80 plays crucial roles in genome stability maintenance and in plant development. *The Plant Journal*, **82**, 655–668.
- Zhao MS, Running SW (2010) Drought-induced reduction in global terrestrial net primary production from 2000 through 2009. *Science*, **329**, 940–943.
- Zust T, Heichinger C, Grossniklaus U *et al.* (2012) Natural enemies drive geographic variation in plant defenses. *Science*, **338**, 116–119.

N.H.D.O., M.P.M.T., K.J.K. performed experiments. N.H.D.O., E.F., K.J.K., G.G., J.V.H. performed data analysis. N.H.D.O., J.J.A.V.L, M.D., G.G., K.J.K., M.P.M.T conceived the experimental design. F.F.M.B. and J.J.B.K. performed experiments related to flowering time. N.H.D.O., E.F., M.D., J.J.A.V.L., K.J.K., J.J.B.K., J.V.H. contributed to the interpretation of the analysis. All authors contributed to writing the manuscript.

### Data accessibility

All data are available on the Molecular Ecology website.

### Supporting information

Additional supporting information may be found in the online version of this article.

**Table S1** Summary of the climate variables mined for this study that vary along geographical gradients.

**Table S2** Bayesian phylogenetic mixed model analysis to assess differences in flowering time without and after vernalization and geographical gradients.

Fig. S1 Geographic distribution of *Arabidopsis thaliana* accessions from the Hapmap population.

Fig. S2 Subset of accessions for Europe.

Fig. S3 Correlation matrix between geography and climate variables for the 308 *A. thaliana* accessions investigated.

Fig. S4 Experimental Design and Treatments Scheme.

Fig. S5 Spatial and genetic distribution of summer annual (219 accessions) and winter annual (89 accessions) life histories in *A. thaliana*.

Fig. S6 Feeding damage (mm<sup>2</sup>) by thrips per genetic clusters.

Fig. S7 Variables that display a geographical gradient.

**Data S1** Data used for downstream analysis. Columns A&B present the unique identifiers for each accession. Column C represents the genetic groups (See M&M, section: Genetic diversity in European accessions from the Hapmap population). Columns D-J present the variables that capture the response of each accession to the corresponding stress (See M&M, section: Statistical analysis).

**Data S2** Results from the MTMM GWAs analysis. MAF (Minor allele frequency). PCp1 = First principal component for traits that were resistant in summer annuals. PCp2 = First principal component for traits that were resistant in winter annuals. MTMM *P*-values for a full model, trait specific and common are presented. Details of the different models are explained in the M&M section. Symbols and descriptions were obtained from TAIR 10. Minimum (Min) and Maximum (Max) relative expression values (log2) for plants exposed to jasmonic acid (JA) or salicylic acid (SA) were obtained from http://bar.utor onto.ca/. In red are indicated the associations identified through MTMM while in black are SNPs in LD ( $r2 \ge 0.5$ ) in a half window of 20 Kb.