



▲ Swarm of migratory locusts in Madagascar.

Crop damage associated with the last major outbreak (1996-2000) led to pesticide spraying of an area of over 4 million ha, thus highlighting the need for an operational warning network.

Main teams

EA GESTER - Gestion des Sociétés, des Territoires et des Risques (UM3)

31 scientists involved in the tobic Director: lean-Marie Miossec. jean-marie.miossec@univ-montp3.fr http://recherche.univ-montp3.fr/gester

LGEI - Laboratoire Génie de l'Environnement Industriel et des Risques Industriels et Naturels (EMA)

20 scientists, with 4 involved in the topic Director: Miguel Lopez-Ferber, miguel.lopez-ferber@ema.fr www.ema.fr/LGEI

BRGM - Service Géologique Régional, Operational services « Eau » et «Aménagement et risques naturels »

(BRGM Languedoc-Roussillon) 29 scientists, with 4 involved in the topic Director: Marc Audibert, m.audibert@brgm.fr www.brgm.fr

UMR Géosciences Montpellier (see page 28)

UMR LISAH - Laboratoire d'étude des Interactions Sol - Agrosystème -Hydrosystème (see page 18)

UMRTETIS - Geoinformation and Earth Observation for Environment and Land Management (see page 8)

...continued on page 50

Spatial analysis to prevent locust outbreaks in Madagascar

The Malagasy migratory locust, Locusta migratoria capito Sauss., is able to change phase according to its population density. The solitary and gregarious phases are characterized by specific behaviours with markedly different morphological, physiological and ecological traits. In the solitary phase, these locusts are dispersed and do not threaten crops. They migrate by flying in search of suitable conditions for their development. If such suitable areas are spatially reduced, the locusts form high density groups. They can then, through a few generations, switch to the gregarious phase, where they form hopper bands and swarms that are capable of destroying crops and pastures.

An operational research project carried out by EMPA (CIRAD), in collaboration with the Malagasy National Locust Centre, is aimed at improving locust forecasting and monitoring systems in order to control this plague. This involves combining monthly information on rainfall, habitats and the biology of the locust (density, phase and stage). Spatial data can also be used to delimit potential grouping areas where the high population densities could trigger phase transformation and subsequent outbreaks.

A spatial delimitation of habitats was carried out through an analysis of around 12 high resolution Landsat images. These spatial data on habitats structures are updated monthly with field data: the rainfall levels and biological features of the migratory locust are integrated in a GIS for interpolation over the entire outbreak source area, i.e. 100 000 km². It is thus possible, on a monthly basis, to monitor the evolution of areas suitable for locust grouping and prevent outbreak risks.

Spatial monitoring of pest locusts in Madagascar, and elsewhere, represents extremely targeted research. It requires in-depth field knowledge, effective management of spatial information and long-term partners in concerned countries.

Contact: Alex Franc, alex.franc@cirad.fr