Representation and functions of shoot apical meristems in FSPMs

KEYNOTE

Romain Barillot¹, Frédéric Boudon², Didier Combes¹, Jean-Louis Durand¹ and Gaëtan Louarn¹ ¹INRAE, URP3F, Lusignan, France ²CIRAD, UMR AGAP Institut, F-34398 Montpellier, France ³UMR AGAP Institut, Univ. Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France

Functional Structural Plant Models (FSPM) are individual-based models that explicitly account for the interactions between plant architecture and its abiotic and biotic environment. Plant morphogenesis, growth and grain or fruit production are among the most represented processes in FSPMs. These biological processes are largely determined at the Shoot Apical Meristem (SAM) level as it drives the rate of leaf initiation, phyllotaxy, leaf geometry, floral induction and the production of reproductive organs. Nevertheless, not all FSPMs are based on an explicit representation of SAMs, their functioning is then ignored in the case of non-dynamic models or embedded in parametric or statistical functions of plant morphogenesis in dynamic models. For FSPMs that include an explicit representation of SAMs, they are mainly constructed on the L-systems formalism which is well suited to describe plant development (Boudon et al., 2012). First, one of the main functions of SAMs in these FSPMs is the production of new leaves or growth units, usually at a constant rate expressed in thermal unit. In some cases, the rate of leaf production by SAMs is coordinated with the rate of leaf emergence, leading to the concept of self-regulated architecture. However, the effect of substrate or water availability on the functioning of the SAM in terms of the size or properties of the emitted primordia is not accounting for. Secondly, the representation of apices in models is also used to simulate axillary bud break and the production of tillers, branches or new growth units. In these cases, the transition from a latent to an active SAM is controlled by light intensity and its spectrum (Verdenal *et al.*, 2008; Faverjon et al., 2019), or temperature or hormones (Prusinkiewicz et al., 2009) or is based on stochastic approaches. Finally, some FSPMs also account for the production of reproductive organs like grains and fruits by SAMs (Boudon et al., 2020; Rouet et al., 2022). Nevertheless, this aspect of SAM functioning remains poorly described in FSPMs despite its importance in plant production, yield and ecology. We believe that a better integration of SAMs functioning and control in FSPMs is a promising way to assess functional hypotheses and predict plant plasticity to the environment.

References

Boudon F, Persello S, Jestin A, Briand A-S, Grechi I, Fernique P, Guédon Y, Léchaudel M, Lauri P-É, Normand F. 2020. V-Mango: a functional-structural model of mango tree growth, development and fruit production. Annals of Botany 126: 745-763.

Boudon F, Pradal C, Cokelaer T, Prusinkiewicz P, Godin C. 2012. L-Py: An L-System Simulation Framework for Modeling Plant Architecture Development Based on a Dynamic Language. Frontiers in Plant Science 3.

Faverjon L, Escobar-Gutiérrez A, Litrico I, Julier B, Louarn G. 2019. A generic individual-based model can predict yield, nitrogen content, and species abundance in experimental grassland communities. Journal of Experimental Botany 70: 2491–2504.

Prusinkiewicz P, Crawford S, Smith RS, Ljung K, Bennett T, Ongaro V, Leyser O. 2009. Control of bud activation by an auxin transport switch. Proceedings of the National Academy of Sciences 106: 17431-17436.

Rouet S, Durand J-L, Leclercq D, Bernicot M-H, Combes D, Escobar-Gutiérrez A, Barillot R. 2022. L-GrassF: a functional-structural and phenological model of Lolium perenne integrating plant morphogenesis and reproductive development. in silico Plants 4: diac012. Verdenal A, Combes D, Escobar-Gutiérrez AJ. 2008. A study of ryegrass architecture as a self-regulated system, using functional-structural plant modelling. Functional Plant Biology 35: 911–924.

Contact: romain.barillot@inrae.fr