Diseases Caused by Bacteria and Phytoplasmas

First Report of Bacterial Leaf Blight Disease of Rice Caused by *Xanthomonas oryzae* pv. *oryzae* in Madagascar

H. Raveloson,^{1,2} R. Rabekijana,^{3,4} N. M. Rakotonanahary,¹ B. Szurek,⁴[©] B. Muller,⁵ K. vom Brocke,⁵ and M. Hutin^{4,†}

- ¹ Centre Régional de Recherche du FOFIFA/DP SPAD Antsirabe, Antsirabe, Madagascar
- ² Institut d'Enseignement Supérieur d'Antsirabe Vakinankaratra (IESAV), Antsirabe, Madagascar
- ³ Université d'Antananarivo, École Supérieure des Sciences
- Agronomiques (ESSA), Antananarivo, Madagascar
- ⁴ IRD, Institut de Recherche Pour le Développement UMR PHIM, Plant Health Institute of Montpellier, University of Montpellier, CIRAD, INRAE, Institut Agro, IRD, Montpellier, France
- ⁵ CIRAD, UMR AGAP Institut, University of Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France

Funding: We acknowledge funding from the European Union through DeSIRA (Development Smart Innovation through Research in Agriculture), as part of the DINAAMICC project (FOOD/2021/422-791). Plant Dis. 107: 2510, 2023; published online as https://doi.org/10.1094/PDIS-03-23-0411-PDN. Accepted for publication 21 April 2023.

In Madagascar, rice is the staple food, and rice production represents a major challenge for food security. Bacterial leaf blight (BLB) caused by Xanthomonas oryzae pv. oryzae (Xoo) is a devastating disease of rice crops in many countries. BLB was first reported in Japan in 1884 and is now prevalent in most of the rice-producing countries of Asia. In Africa, Xoo was reported for the first time in Mali and Cameroon (Buddenhagen et al. 1979) and later in many West African countries. In contrast, in East Africa, BLB seems less prominent and has only been reported in Tanzania and Uganda (Oliva et al. 2019). To our knowledge, in Madagascar, despite continuous monitoring for rice diseases since the 1980s, BLB had never been reported until this study. In December 2019, BLB-like symptoms were observed in two fields in Vakinankaratra, the Central Highlands of Madagascar. In December, most rice fields in Vakinankaratra were at the seedling stage where typical symptoms are often not visible, making it impossible to assess disease severity and incidence. Typical symptoms were characterized by yellow to grayish, water-soaked lesions that start from the leaf tip and progress along the central vein or leaf margin. At an advanced stage of the

disease, the leaves were completely desiccated sometimes with droplets of yellow exudate at the leaf margin. Four symptomatic leaf fragments from two fields were successively disinfected, rinsed in sterile water, and ground. Leaf powder was resuspended in 1.5 ml of sterile water, of which 10 µl was plated on semiselective PSA medium and incubated at 28°C for 3 to 7 days. Typical Xoo-like single colonies were purified from four individual leaf samples from two distinct sites. The multiplex PCR developed for the identification of X. oryzae pathovars (Lang et al. 2010) produced, for eight isolates (two per leaf), the diagnostic two-band pattern characteristic of Xoo. Fiveweek-old plants of Oryza sativa cultivar IR24 were subjected to a pathogenicity test performed with four strains. Leaves were clipped with scissors and dipped in a bacterial suspension at an optical density of 0.2 as previously described (Kauffman et al. 1973). Fourteen days after incubation of rice plants in a greenhouse (27 \pm 1°C with a 12-h photoperiod), the inoculated leaves (n = 8) developed typical BLB lesions identical to those challenged with the Xoo reference strain PXO99^A. In contrast, water-inoculated leaves remained asymptomatic. The lesions were subjected to strain isolation, and purified strains were validated as Xoo by multiplex PCR as described above, thus fulfilling Koch's postulates. Finally, we amplified by PCR and sequenced the housekeeping gene gyrB for two of the isolates using the primers XgyrB1F and XgyrB1R (Young et al. 2008). Analysis of 825 bp of the gyrB sequence of the strains Md5 and Md11 revealed 100% identity with that of the Indian Xoo strain BXO1 (accession no. CP033201) and highlighted 10 and 12 polymorphic nucleotides compared with the gyrB sequence of the Xoo reference strains PXO99^A from the Philippines (accession no. CP000967) and BAI3 from West Africa (accession no. CP025610). To our knowledge, this is the first report of BLB in Madagascar. Surveys conducted in the Vakinankaratra and Menabe regions between 2020 and 2022 confirmed the presence of BLB in Madagascar and a sharp increase in disease incidence. Further surveys will be necessary to evaluate if the disease has already spread over the island, and further analysis will be needed to understand the origin of the outbreak and identify strategies to control BLB in Madagascar.

References:

Buddenhagen, I. W., et al. 1979. Int. Res. Newsl. 4:11. Kauffman, H. E., et al. 1973. Plant Dis. Rep. 57:537. Lang, J. M., et al. 2010. Plant Dis. 94:311. Oliva, R., et al. 2019. Nat. Biotechnol. 37:1344. Young, J. M., et al. 2008. Syst. Appl. Microbiol. 31:366.

The author(s) declare no conflict of interest.

e-Xtra

Keywords: bacterial leaf blight, Madagascar, Xanthomonas oryzae

[†]Indicates the corresponding author. M. Hutin; mathilde.hutin@ird.fr

This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of the source. The American Phytopathological Society, 2023.