

Genetic bases of fruity and flora aroma of the Nacional cocoa variety

K. Colonges^{1,2}; J.C. Jimenez⁴; A. Saltos⁴; M.C. Lahon¹; M. Lebrun¹; I. Maraval¹; D. Calderon⁴; C. Subia⁴; I. Sotomayor⁴; F. Fernández⁴; P. Costet⁴; E.d Seguine⁵; O. Fouet²; B. Rhoné²; X. Argout²; S. Assemat¹; F. Davrieux¹; E. Cros¹; G. Loor Solorzano⁴; C. Lanaud²; R. Boulanger¹

1 Qualisud, Univ Montpellier, Avignon Université, Cirad, Institut Agro, IRD, Université de La Réunion, Montpellier, France.

2 AGAP Institut, Univ Montpellier, Cirad, INRAE, Institut Agro, Montpellier, France.

3 Instituto Nacional de Investigacion Agropecurias, INIAP, Ecuador.

4 Valrhona, Tain l'Hermitage, France

5 Guittard, United-States

Montpellier, France December 5-7



AmaCacao

Project Leader:
C. Lanaud/R. Boulanger

Total Budget total: 427K€

- MUSE
- Valrhona



Museo
Nacional
de Colombia



MUSEO ANTROPOLOGICO Y
DE ARTE CONTEMPORANEO


Guittard
CHOCOLATIER



VALRHONA


International
Symposium on
Cocoa
Research


AmaZcacao



- History of past domestication of aromatic fine cocoa trees using paleogenomics
- Rescue and exploitation of new genetic resources related to fine cocoa
- Study of genetic and biochemical determinants of fine cocoa aromas
- Genetic basis of cadmium accumulation
- Study of dynamics of fine cocoa markets



Location of cultivation sites in Ecuador




The plant material were composed of a collection from Ecuador :

- Pichilingue experimental station of INIAP
- Coleccion de Cacao de Aroma Tenguel (CCAT) of Tenguel

This population (152) represents the Nacional variety currently grown in Ecuador

A second population of native cocoa trees (202) in the south of the Ecuadorian Amazon in the area of origin of Nacional

Phénotypes for Genome Wide Association Study (GWAS)



Methods

Genome Wide Association Study (GWAS)

Genotypes

Genotyping 152 trees

SSR + SNP (GBS)

O. Fouet X. Argout


Nacional

Floral

Lanaud *et al*, PAG 2012
Collaboration CIRAD/INIAP

Amelonado

Cacao



Phenotypes

Sensorial characterization

36 sensorial traits

(Fruity, Floral, Bitterness,
Astringency ...)

E. Seguine



Biochemical characterization

65 volatile compounds


unroasted beans

GC/MS R. Boulanger E. Cros J-C. Jimenez

87 volatile compounds

roasted beans

GC/MS R. Boulanger E. Cros A. Saltos




8 non volatile compounds


unroasted beans

NIRS F. Davrieux A. Saltos


Association genetics (GWAS): example of results for aroma compounds



Association genetics (GWAS): example of results for floral notes



Candidates genes research in association areas



Determining the floral and fruity aromas of modern Nacional

Floral notes - 16 sensory descriptors, 27 VOCs

48 Association areas (9 sens.; 39 VOCs)

2 Main biosynthetic pathways:

- Monoterpene biosynthetic pathway (**9 Candidate Genes**)
- L-phenylalanine degradation pathway (**19 Candidate Genes**)



Fruity notes - 13 sensory notes and 35 VOCs

233 Association areas (39 sens.; 194 VOCs)


5 Biosynthetic pathways:

- Monoterpene biosynthetic pathway
- Pathway of L-phenylalanine
- Degradation of proteins, sugars and fatty acids

(514 Candidate Genes)



Example for candidate genes, floral and fruity notes L-phenylalanine pathway



58 gènes
candidats

Determining the floral and fruity aromas of National ancestral

A second population of native cocoa trees resulting from surveys carried out in the south of the Ecuadorian Amazon in the area of origin of Nacional

Floral notes – 16 sensory descriptors, 27 COVs

393 Zones d'associations (38 sens.; 355 COVs)

- monoterpene biosynthetic pathway (114 Candidate Genes)
- L-phenylalanine degradation pathway (41 Candidate Genes)



Fruity notes - 13 sensory descriptors and 35 COVs

502 Zones d'associations (22 sens.; 480 COVs)

- monoterpene biosynthetic pathway
- pathways of L-phenylalanine (1108 Candidate Genes)
- degradation of proteins, sugars and fatty acids



54 Common genes between the two populations

Determining the floral and fruity aromas of modern Nacional

- 5 biosynthetic pathways highlighted:
 - 2 for the floral notes 
 - 5 for the fruity notes 
- Characterization of genome zones in relation to aroma types
- Selection of areas of the genome favourable to the desired notes

Nest step

- Expression analyses at different stages



*Thanks for your
attention*



This work, part of the MUSE Amazcacao project, was publicly funded through ANR (the French National Research Agency) under the 'Investissement d'avenir' programme with the reference ANR-16-IDEX-0006.

We thank the I-Site MUSE, Valrhona and the U.S. Department of Agri-culture (USDA-ARS), for their financial support of this project.