





# Afforestation of savannah using cocoa agroforestry: impacts on ecosystem services and effects of associated tree species on soil fertility

Jean-Michel Harmand, Annemarijn Nijmeijer, Marie Sauvadet, Seguy Enock, Jean-Daniel Essobo, Eltson Fonkeng, Pierre-Eric Lauri, Patrick Jagoret, Stephane Saj







#### Preliminary: Natural encroachment of forest into savannah

In the North of the Congo Basin, different past and recent studies showed **natural forest expansion over savannah** 

Gillet et al, 2001. J Trop Ecol: 809–832.
<u>http://journals.cambridge.org/abstract\_S0266467401001614</u>
Mitchard ETA, 2011. Remote Sens Environ 115: 2861–2873.
<u>https://www.sciencedirect.com/science/article/abs/pii/S0034425711001337</u>
Mitchard ETA, 2013. Phil Trans R Soc B 368: 20120406.
<u>https://royalsocietypublishing.org/doi/10.1098/rstb.2012.0406</u>

© JM Harmand, Cirad

But this natural expansion of forest over savannah is currently **disrupted and often stopped by human activities**, such as cropping, **leading to degraded gramineous savannah ecosystems which are burned every year and periodically cultivated**.



Forest and Cocoa agroforest - Savannah boundary in the Bokito area (Cameroun)

#### Afforestation of savannah with cocoa

- Farmers have proven their ability to create cocoa agroforests on savannah
- What is observed: Full-grown cocoa AFS created on savannah and in forest show comparable multistrata structures.
- Question : Do cocoa agroforests created on savannah have the same potential of cocoa plantations created in neighbouring forests ?



Young cocoa plants under plantain and young trees



Mature cocoa plantation set up after savannah

Jagoret P. et al., 2012. Agrofor Syst, 493-504. <u>https://doi.org/10.1007/s10457-012-9513-9</u>

2022 International Symposium on Cocoa Research (ISCR), Montpellier, France

3

### **Objective of the first study**

- Assess the impacts of previous land uses (forest or savannah) of cocoa agroforesty plantations on ecosystem services over time (PhD Annemarijn Nijmeijer).



Rainfall: 1400 mm yr<sup>-1</sup>, 3 months of dry season

Nijmeijer A, et al 2019. Agr Ecos Env 275: 100-111. <u>https://doi.org/10.1007/s10457-017-0182-6</u>

2022 International Symposium on Cocoa Research (ISCR), Montpellier, France

#### **Ecosystem attributes (EA) of cocoa agroforests**



Similar values of EA in cocoa plantations after savannah and forest some decades after establishment

- EA of cocoa plantations set up after savannah increased with time and reached similar levels of EA in cocoa plantations after forest.
- 20 to 30 years to reach the same values for species diversity, cocoa yield, leaf litterfall
- 50 to 70 years to reach the same values for C storage and some components of soil quality.

Adapted from: Nijmeijer A, et al 2019. Agr Ecos Env 275: 100-111. https://doi.org/10.1007/s10457-017-0182-6

2022 International Symposium on Cocoa Research (ISCR), Montpellier, France

#### **Objective of the second study**

 Assess the impacts of agroforestry and different shade trees on soil fertility (top 10 cm soil layer) and cocoa yield in relation to tree functional traits



Sauvadet M et al, 2020. J Applied Ecol 57 (3):476-487. <u>https://doi.org/10.1111/1365-2664.13560</u>

6

#### Dynamic of soil C of the top 10 cm for different land uses



AGRICULTURE, FOREST MANAGEMENT, RESEARCH SUMMARY

#### Improving soil fertility in cocoa agroforests using the most suitable shade tree species

Journal of Applied Ecology

#### THE APPLIED ECOLOGIST'S BLOG

Bridging the gap between researchers, practitioners and policymakers

http://appliedecologistsblog.com/2020/02/18/improving-soil-fertility-in-cocoa-agroforests-using-the-most-suitable-shade-tree-species



Sauvadet M et al, 2020. J Applied Ecol 57 (3):476-487. https://doi.org/10.1111/1365-2664.13560

2022 International Symposium on Cocoa Research (ISCR), Montpellier, France

### Cocoa yield



- No visible impact of shade on yield (high variability of data);
- Young cocoa monoculture of 10 years old not more productive than old cocoa agroforestry system

#### **Cocoa nutritional status**



In comparison to cocoa monoculture:

- soil inorganic P,
- and cocoa leaf P concentration
   were increased in cAFS with
   differences between tree species.

#### Conclusions

- Planting cocoa trees together with shade trees in savannah can increase soil fertility, species diversity and carbon storage in biomass and soil over time;
- Cocoa-based AFS set up after forest or after savannah display with time convergent profiles of ecosystem attributes (including cocoa production);
- Tree species effects on **soil properties** can be explained by **leaf litter traits**;
- Our results suggest that deep uptake of nutrients by associated tree species and their recycling through litterfall (data not shown) increase top soil nutrient contents and contribute to nutrient export by cocoa harvest;
- Using specific shade trees for the provision of goods, and other trees for soil quality, both compatible with cocoa, can ensure multifunctionality and sustainability of cocoa systems;
- Further studies are necessary to understand how plant diversity can help to adapt to climate change in the context of the forest-savannah transition zone.

## Acknowledgements





D P Agroforesterie Cameroun







Institut de Recherche pour le Développement F R A N C E





## Thank you for your attention



agropolis fondation

