

Detection of Crimean–Congo haemorrhagic fever virus in *Hyalomma marginatum* ticks, southern France, May 2022 and April 2023

Célia Bernard^{1,2}, Charlotte Joly Kukla^{1,2,7}, Ignace Rakotoarivony^{1,2}, Maxime Duhayon^{1,2}, Frédéric Stachurski^{1,2}, Karine Huber^{1,2}, Carla Giupponi^{1,2}, Lyonna Zortmann^{1,2}, Philippe Holzmüller^{1,2}, Thomas Pollet^{1,2}, Mélanie Janneau^{1,2}, Alice Mercey^{1,2}, Nathalie Vachery^{1,2}, Thierry Lefrançois⁸, Claire Garros^{1,2}, Vincent Michaud^{1,2}, Loïc Comtet⁶, Léa Despois⁶, Philippe Pourquier⁶, Caroline Picard^{3,4}, Alexandra Journeaux^{3,4}, Damien Thomas^{3,5}, Sabine Godard^{3,5}, Elodie Moissonnier^{3,5}, Stéphane Mely^{3,5}, Manon Segal^{3,5}, Delphine Pannetier^{3,5}, Sylvain Baize^{3,4}, Laurence Vial^{1,2}

1. Centre de coopération internationale en recherche agronomique pour le développement (CIRAD), University of Montpellier (UMR) Animal Santé Territoires Risques Écosystèmes (ASTRE), Montpellier, France
2. ASTRE UMR, CIRAD, Institut national de la recherche agronomique (INRAE), Montpellier, France
3. National Reference Center for Viral Hemorrhagic Fevers, Lyon, France
4. Unité de Biologie des Infections Virales Emergentes, Institut Pasteur - Centre International de Recherche en Infectiologie (CIRI), Université de Lyon, Institut national de la santé et de la recherche médicale (INSERM) U1111, Ecole Normale Supérieure de Lyon, Université Lyon 1, CNRS UMR5308, Lyon, France
5. Laboratoire P4 INSERM Jean Mérieux, INSERM Lyon, France
6. Innovative Diagnostics, Grabels, France
7. Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail (ANSES), INRAE, Ecole Nationale Vétérinaire d'Alfort, UMR BIPAR, Laboratoire de Santé Animale, Maisons-Alfort, France
8. CIRAD, DG, Paris, France

Correspondence: Célia Bernard (celia.bernard@cirad.fr)

Citation style for this article:

Bernard Célia, Joly Kukla Charlotte, Rakotoarivony Ignace, Duhayon Maxime, Stachurski Frédéric, Huber Karine, Giupponi Carla, Zortmann Lyonna, Holzmüller Philippe, Pollet Thomas, Janneau Mélanie, Mercey Alice, Vachery Nathalie, Lefrançois Thierry, Garros Claire, Michaud Vincent, Comtet Loïc, Despois Léa, Pourquier Philippe, Picard Caroline, Journeaux Alexandra, Thomas Damien, Godard Sabine, Moissonnier Elodie, Mely Stéphane, Segal Manon, Pannetier Delphine, Baize Sylvain, Vial Laurence. Detection of Crimean–Congo haemorrhagic fever virus in *Hyalomma marginatum* ticks, southern France, May 2022 and April 2023. Euro Surveill. 2024;29(6):pii=2400023. <https://doi.org/10.2807/1560-7917.ES.2024.29.6.2400023>

Article submitted on 10 Jan 2024 / accepted on 01 Feb 2024 / published on 08 Feb 2024

Crimean–Congo haemorrhagic fever (CCHF), a potentially severe zoonotic viral disease causing fever and haemorrhagic manifestations in humans. As the Crimean–Congo haemorrhagic fever virus (CCHFV) has been detected in ticks in Spain and antibodies against the virus in ruminant sera in Corsica, it was necessary to know more about the situation in France. In 2022–2023, CCHFV was detected in 155 ticks collected from horses and cattle in southern France.

The emergence and spread of the tick-borne viral disease Crimean–Congo haemorrhagic fever (CCHF) pose significant challenges to public health. Transmission of the virus to humans occurs predominantly via bites of *Hyalomma* ticks, in Europe especially the species *H. marginatum* and *H. lusitanicum*, or via exposure to infected blood or tissues of viraemic animals or humans [1–3]. Here we describe the detection of Crimean–Congo haemorrhagic fever virus (CCHFV) from ticks in southern France.

Risk-based sampling


As ticks are the only known natural reservoirs of CCHFV [4], we focused on field collection of ticks. In May 2022, during the peak activity of ticks, we collected ticks mainly from horses in four Mediterranean departments (NUTS-3) on the French mainland bordering Spain

(Figure 1). Horses are the likely hosts of adult stages of *H. marginatum* [5,6].

Cattle are considered good amplifiers of CCHFV and thus enhance local virus circulation [7]. In 2023, we optimised the sampling by collecting ticks from cattle in the Pyrénées-Orientales department (Figure 1) where antibodies against CCHFV were identified in 2021–2022 from cattle (maps showed in <https://www.anses.fr/fr/system/files/SABA2020SA0039Ra.pdf>). Cattle farms with the highest within-herd seroprevalences were selected, as well as a few seronegative farms in the same zones. In addition, farms with horses (not previously tested with serology), located in the neighbourhood of the seropositive cattle farms were also visited, especially when the cattle farmers did not give their consent to sampling. As *Hyalomma* ticks are not located within cattle barns but likely in the natural farm environments, specific pastures (e.g. shrublands) were selected. In spring, cattle from different farms are gathered and grazed in such shrublands, and these spring pastures were thus considered as suitable sites for collection of *H. marginatum* [8]. Ticks were collected in April when adult *H. marginatum* search for hosts and potentially infect naïve cattle.

FIGURE 1

Map showing areas where ticks were collected from cattle and horse farms for analysis of Crimean–Congo haemorrhagic fever virus, France, May 2022 and April 2023 (n =57)



Boxes indicate the name of each department, the animal species (horse or cattle) from which ticks were collected and the number of ticks tested positive for Crimean–Congo haemorrhagic fever virus. The departments coloured green were sampled in 2022 and the one coloured purple was sampled in 2022 and 2023.

Laboratory investigations

Ticks were species identified morphologically by experienced acarologists using relevant identification keys [9,10], and only *H. marginatum* ticks were included in the further analysis and stored at -80°C.

After being washed in bleach for 30 s and rinsed three times in water, the ticks were crushed individually in 400 µL of Dulbecco's Modified Eagle Medium (DMEM) (Eurobio Scientific, Les Ulis, France) supplemented with 10% fetal calf serum. Total RNA was extracted using a kit, NucleoMag VET extraction kit (Macherey-Nagel, Düren, Germany), according to the instructions of the manufacturer, using IDEALTM 96 extraction robot (Innovative Diagnostics, Montpellier, France). Then, CCHFV simplex reverse transcription (RT) quantitative PCRs were run, adapted from previously published studies [11,12]. Optimisations, including modifications to the PCR thermoprofile and mix, were carried out by Innovative Diagnostics, and allowed the quickest detection and the best amplification curve for a synthetic RNA used as positive control. The PCR detection limit was estimated to five viral genomic copies.

Positive tick homogenates as well as a set of negative ones were sent for confirmation to the Biosafety Level 4 laboratory of the National Reference Center for Viral Hemorrhagic Fevers (NRC), Lyon, France. After removing tick tissues from suspensions, RNA was extracted using the QIAamp viral RNA mini kit (Qiagen, Hilden, Germany) following the manufacturer's instructions. Thereafter, detection of the virus was performed using the QIAGEN OneStep RT-PCR kit (Qiagen), according to a protocol adapted from Wölfel et al. [12].

The NRC selected samples with quantification cycle (Cq) values <25 for whole genome sequencing. The RNA was subjected to a DNase digest (Turbo DNase, Thermo Fisher Scientific, Waltham, the United States (US)). Then, library preparation was performed using the NEBNext Ultra II RNA Library Prep Kit for Illumina (New England Biolabs, Ipswich, US). Sequencing was performed using a MiniSeq platform (Illumina Inc, San Diego, US). The paired-end reads generated were trimmed and de-novo assembled using naviralSPAdes (version 3.15.4) on Galaxy platform (version 23.2.rc1) (<https://usegalaxy.org/>). The CCHFV genomes (segment S, M and L) were searched from the contigs using Basic Local Alignment Search Tool (BLAST)

(version 2.14.1). We used the BioEdit software (version 7.1.3.0) (<https://bioedit.software.informer.com/>) to align the nucleotide sequences obtained with published sequences belonging to the different lineages of CCHFV.

Findings of ticks and Crimean–Congo haemorrhagic fever virus

In 2022, an average of 30 *H. marginatum* ticks per location, and in 2023 all ticks, were analysed for CCHFV.

In 2022, ticks were collected on 33 horse and three cattle farms. In total, 997 ticks of *H. marginatum* were identified and CCHFV was detected in 13 (1.3%) ticks (all from the same cattle farm in Pyrénées-Orientales) (Figure 1). In 2023, in Pyrénées-Orientales, ticks were collected on 15 cattle farms, three farms with cattle and horses and three farms with horses. A total of 1,001 *H. marginatum* ticks were collected and analysed, and 142 (14.2%) were positive for CCHFV. Most of these positive ticks were collected from cattle farms ($n = 11$), except for 12 ticks from two horse farms (Figure 1).

Considering 2022 and 2023 data, the proportion of infected ticks in positive farms varied from 3.1% to 55.8% (median: 7.4%) or 1 to 66 positive ticks per farm, with four farms with infection rates up to 20%. The Cq values ranged from 18.25 to 40.82 (median: 36.39), which suggested a high viral load in some ticks. At the NRC, CCHFV RNA was confirmed in 132 (85.2%) of 155 samples, and six additional ones among the 114 initially negative samples. All Cq values from the 23 non-confirmed positive samples were high (around 40). All obtained sequences clustered with CCHFV strains, such as the Caceres strain previously isolated from ticks in Spain, within the genotype III (Figure 2).

Discussion

For the first time, CCHFV was detected in resident tick populations in the mainland of France, confirming its circulation. Crimean–Congo haemorrhagic fever is one of the most widespread viral tick-borne zoonoses worldwide with severe consequences for healthcare, such as intensive case management, 5–30% fatality rate in haemorrhagic patients and the need to prevent nosocomial infections [13]. Endemic in Africa, the Middle East and other Asian countries and in the Balkans, CCHF was recently detected in Spain, western Europe [14]. Transmission of the virus to humans occurs predominantly via tick bites or via exposure to infected blood or tissues of viraemic animals or patients [1]. Given the recent findings of *H. marginatum*, one main competent vector for CCHFV, in the south of France [8,15] near the Spanish border, it was crucial to evaluate the epidemiological situation of CCHFV in France.

We considered that CCHFV circulation in France would be at a low level. Indeed, immature stages of *H. marginatum* are mainly found on birds and lagomorphs. Most bird species are refractory to CCHFV and thus unable to reinfect naïve ticks, whereas lagomorphs are good


amplifiers of the virus. In France, bird populations are sizeable in the south of the country. The populations of hares and rabbits have decreased in France since the outbreak of myxomatosis in the 1950s and the current occurrence of rabbit viral haemorrhagic disease. Consequently, the probability for *H. marginatum* to become infected at the immature tick stage in France is considered low [7]. Furthermore, adult stages of *H. marginatum* have a marked trophic preference for horses in France [5], and horses are poor amplifiers of CCHFV (unable to reinfect new ticks) although they are very abundant in the south of France. Thus, the probability for *H. marginatum* to become infected at the adult stage is considered low [7].

Our findings of a high proportion of infected ticks in some farms further reinforced that this cannot be due to sporadic introduction of infected *H. marginatum* ticks from endemic countries but local virus transmission. However, such introduction events remain a risk to be monitored, especially to detect the emergence of new CCHFV genotypes [16–18]. All CCHFV isolates sequenced in this study were highly identical and belonged to the same genotype. Ticks infected with CCHFV were predominantly from cattle although many horses were examined, with high proportions of infected ticks in four facilities where very few animals were examined (4–6 animals per farm). This strengthens the importance of cattle in the CCHFV transmission and the possibility of having sampled viraemic animals. As horses are not able to replicate sufficiently CCHFV to infect naïve ticks [19], finding a few infected ticks on horses indicated that these *H. marginatum* were infected at immature stages or through transovarial transmission, and then maintained the virus along their development.

Because of the risk-based sampling method used, the proportion of infected ticks does not reflect the prevalence of *H. marginatum* ticks infected with CCHFV. The prevalence of infected ticks is presumably lower which may explain that no human cases have, so far, been notified in France. Additionally, genotype III, which was identified from the first Spanish cases [20], is considered of low virulence. At present, in France, humans may become infected via bites of adult *H. marginatum* ticks. This risk exists during the activity period of the ticks, namely from April to July. Only ticks seeking hosts in the environment should be considered as possible vehicles for virus transmission; those already attached to animals cannot detach and reattach. Risk areas for CCHFV transmission present conditions conducive to the presence of *H. marginatum* (open dry natural environments, typical of the Mediterranean area), but infected ticks were not found in all locations infested by *H. marginatum* and further parameters suitable for local circulation of CCHFV between ticks and animals, remain to be determined. Another CCHFV transmission pathway could be contact with infected animal material, as animals can be viraemic, albeit asymptomatic, during the seasonal activity of *H. marginatum* e.g. from April to July when adult stages

FIGURE 2

Phylogenetic analysis of Crimean–Congo haemorrhagic fever virus S, M and L segment sequences, France obtained in May 2022 and April 2023 and other countries



^a This designation is without prejudice to positions on status and is in line with United Nations Security Council Resolution 1244/99 and the International Court of Justice Opinion on the Kosovo Declaration of Independence.

The analysis was performed by using IQ-TREE software (version 2.1.2) (<http://www.iqtree.org/>).

Bootstrap values (>50%) are shown at nodes. Scale bar represents the estimated number of substitutions per site. Individual sequences are named with strain name, country of origin and GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) accession number, if available. Each colour represents a genotype already described. All phylogenetic analyses were conducted on the complete coding DNA sequence (nucleoprotein for the S segment and RNA-dependent RNA polymerase for the L segment) except for the M segment for which partial sequences were obtained (from Nairovirus M polyprotein-like region to codon STOP).

parasite wild and domestic ungulates and from August to October when immature stage parasite lagomorphs. Considering such risks, preventive recommendations towards at-risk populations are needed.

Ethical statement

We informed farmers and horse owners of the objectives of our study and asked them to sign individual agreements for the collection and analysis of ticks on their animals. A legal procedure has been also set up to protect breeders' personal data. All data have been anonymised.

Funding statement

The authors thank the funders who made this work possible: French Ministry of Agriculture—General Directorate for Food (DGAL, grant agreement: SPA17 number 0079-E), European Funds for Regional Development (FEDER, Grand-Est), French Establishment for Fighting Zoonoses (ELIZ) and the Association Nationale Recherche Technologie (ANRT, grant agreement number: 2019-1145), Défi clé RIVOC (Occitanie Region): Holis-tiques project. In addition, the National Reference Center for Viral Hemorrhagic Fevers is funded by Santé Publique France (Saint Maurice, France).

Data availability

Sequences were deposited on GenBank (Accession Numbers: PPO67047 to PPO67088). The location and identification of the farms included in the study have been protected.

Acknowledgements

The authors would like to thank Harold Noël, Alexandra Maille and Julie Figoni of Santé Publique France for their help. We would also like to specifically thank Christophe Leculier and Béatrice Labrosse, head of biological safety at the BSL-4 INSERM Jean Mérieux Laboratory for their involvement in the logistical and operational management of the samples in the containment laboratories and Vincent Lotteau as BSL-4 director. We would like to thank all cattle and horse owners for collaborating with us on this study.

Conflict of interest

None declared.

Authors' contributions

Célia Bernard and Laurence Vial designed the study and contacted the stakeholders in the field. Sampling was carried out by Célia Bernard, Laurence Vial, Charlotte Joly Kukla, Ignace Rakotoarivony, Maxime Duhayon, Frederic Stachurski, Karine Huber, Carla Giupponi, Lyonna Zortmann, Philippe Holzmuller, Thomas Pollet and Alice Mercey. Laurence Vial and Frédéric Stachurski did the morphological tick identifications. Sample treatments were carried out by Célia Bernard, Charlotte Joly Kukla, Alice Mercey, Melanie Jeanneau, Loïc Comtet, Lea Despois and Philippe Pourquier. Confirmation and sequencing were done by Caroline Picard, Alexandra Journeaux, Damien Thomas, Sabine Godard, Elodie Moissonnier, Stéphane Mely, Manon Segal, Delphine Pannetier and Sylvain Baize. Célia Bernard, Laurence Vial, Loïc Comtet, Lea Despois, Sylvain Baize, Delphine Pannetier and Caroline Picard wrote the article. Laurence Vial, Nathalie

Vachieri, Thierry Lefrançois, Claire Garros, Sylvain Baize and Vincent Michaud communicated about the detection and had contacts with the governmental institutions.

References

1. Fillâtre P, Revest M, Tattevin P. Crimean-Congo hemorrhagic fever: An update. *Med Mal Infect.* 2019;49(8):574-85. <https://doi.org/10.1016/j.medmal.2019.09.005> PMID: 31607406
2. Gargili A, Estrada-Peña A, Spengler JR, Lukashev A, Nuttall PA, Bente DA. The role of ticks in the maintenance and transmission of Crimean-Congo hemorrhagic fever virus: A review of published field and laboratory studies. *Antiviral Res.* 2017;144:93-119. <https://doi.org/10.1016/j.antiviral.2017.05.010> PMID: 28579441
3. Bonnet SI, Bertagnoli S, Falchi A, Figoni J, Fite J, Hoch T, et al. An update of evidence for pathogen transmission by ticks of the genus *Hyalomma*. *Pathogens.* 2023;12(4):513. <https://doi.org/10.3390/pathogens12040513> PMID: 37111399
4. Hawman DW, Feldmann H. Crimean-Congo haemorrhagic fever virus. *Nat Rev Microbiol.* 2023;21(7):463-77. <https://doi.org/10.1038/s41579-023-00871-9> PMID: 36918725
5. Grech-Angelini S, Stachurski F, Lancelot R, Boissier J, Allienne JF, Marco S, et al. Ticks (Acari: Ixodidae) infesting cattle and some other domestic and wild hosts on the French Mediterranean island of Corsica. *Parasit Vectors.* 2016;9(1):582. <https://doi.org/10.1186/s13071-016-1876-8> PMID: 27842608
6. Stachurski F, Vial L. Installation de la tique *Hyalomma marginatum*, vectrice du virus de la fièvre hémorragique de Crimée-Congo, en France continentale. [Emergence of *Hyalomma marginatum* tick, vector of the Crimean–Congo haemorrhagic fever virus, in continental France]. *Bulletin épidémiologique Santé animale – alimentation* 2018;84(8):1-5. French. Available from: https://be.anses.fr/sites/default/files/O-005_2018-07-23_Hyalomma-Stachurski_VF.pdf
7. Bernard C, Holzmuller P, Bah MT, Bastien M, Combes B, Jori F, et al. Systematic review on Crimean–Congo Hemorrhagic Fever enzootic cycle and factors favoring virus transmission: special focus on France, an apparently free-disease area in Europe. *Front Vet Sci.* 2022;9:932304. <https://doi.org/10.3389/fvets.2022.932304> PMID: 35928117
8. Bah MT, Grosbois V, Stachurski F, Muñoz F, Duhayon M, Rakotoarivony I, et al. The Crimean-Congo haemorrhagic fever tick vector *Hyalomma marginatum* in the south of France: Modelling its distribution and determination of factors influencing its establishment in a newly invaded area. *Transbound Emerg Dis.* 2022;69(5):e2351-65. <https://doi.org/10.1111/tbed.14578> PMID: 35511405
9. Estrada-Peña A. Ticks as vectors: taxonomy, biology and ecology. *Rev Sci Tech.* 2015;34(1):53-65. <https://doi.org/10.20506/rst.34.1.2345> PMID: 26470449
10. Pérez-Eid C. Les tiques : identification, biologie, importance médicale et vétérinaire. [Ticks: identification, biology, importance in medicine and veterinary medicine]. Cachan: Lavoisier; 2007. p. 339. French.
11. Sas MA, Vina-Rodriguez A, Mertens M, Eiden M, Emmerich P, Chaintoutis SC, et al. A one-step multiplex real-time RT-PCR for the universal detection of all currently known CCHFV genotypes. *J Virol Methods.* 2018;255:38-43. <https://doi.org/10.1016/j.jviromet.2018.01.013> PMID: 29408661
12. Wölfel R, Paweska JT, Petersen N, Grobbelaar AA, Leman PA, Hewson R, et al. Virus detection and monitoring of viral load in Crimean-Congo hemorrhagic fever virus patients. *Emerg Infect Dis.* 2007;13(7):1097-100. <https://doi.org/10.3201/eid1307.070068> PMID: 18214191
13. Leblebicioglu H, Bodur H, Dokuzoguz B, Elaldi N, Guner R, Koksali I, et al. Case management and supportive treatment for patients with Crimean-Congo hemorrhagic fever. *Vector Borne Zoonotic Dis.* 2012;12(9):805-11. <https://doi.org/10.1089/vbz.2011.0896> PMID: 22607078
14. Al-Abri SS, Abaidani IA, Fazlalipour M, Mostafavi E, Leblebicioglu H, Pshenichnaya N, et al. Current status of Crimean-Congo haemorrhagic fever in the World Health Organization Eastern Mediterranean Region: issues, challenges, and future directions. *Int J Infect Dis.* 2017;58:82-9. <https://doi.org/10.1016/j.ijid.2017.02.018> PMID: 28259724
15. Vial L, Stachurski F, Leblond A, Huber K, Vourc'h G, René-Martellet M, et al. Strong evidence for the presence of the tick *Hyalomma marginatum* Koch, 1844 in southern continental France. *Ticks Tick Borne Dis.* 2016;7(6):1162-7. <https://doi.org/10.1016/j.ttbdis.2016.08.002> PMID: 27568169
16. Monsalve Arteaga L, Muñoz Bellido JL, Negrodo AI, García Criado J, Vieira Lista MC, Sánchez Serrano JA, et al. New

- circulation of genotype V of Crimean-Congo haemorrhagic fever virus in humans from Spain. *PLoS Negl Trop Dis*. 2021;15(2):e0009197. <https://doi.org/10.1371/journal.pntd.0009197> PMID: 33617538
17. Moraga-Fernández A, Ruiz-Fons F, Habela MA, Royo-Hernández L, Calero-Bernal R, Gortazar C, et al. Detection of new Crimean-Congo haemorrhagic fever virus genotypes in ticks feeding on deer and wild boar, Spain. *Transbound Emerg Dis*. 2021;68(3):993-1000. <https://doi.org/10.1111/tbed.13756> PMID: 32738065
 18. Palomar AM, Portillo A, Santibáñez P, Mazuelas D, Arizaga J, Crespo A, et al. Crimean-Congo hemorrhagic fever virus in ticks from migratory birds, Morocco. *Emerg Infect Dis*. 2013;19(2):260-3. <https://doi.org/10.3201/eid1902.121193> PMID: 23347801
 19. Spengler JR, Estrada-Peña A, Garrison AR, Schmaljohn C, Spiropoulou CF, Bergeron É, et al. A chronological review of experimental infection studies of the role of wild animals and livestock in the maintenance and transmission of Crimean-Congo hemorrhagic fever virus. *Antiviral Res*. 2016;135:31-47. <https://doi.org/10.1016/j.antiviral.2016.09.013> PMID: 27713073
 20. Negrodo A, de la Calle-Prieto F, Palencia-Herrejón E, Mora-Rillo M, Astray-Mochales J, Sánchez-Seco MP, et al. Autochthonous Crimean-Congo Hemorrhagic Fever in Spain. *N Engl J Med*. 2017;377(2):154-61. <https://doi.org/10.1056/NEJMoa1615162> PMID: 28700843

License, supplementary material and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence and indicate if changes were made.

Any supplementary material referenced in the article can be found in the online version.

This article is copyright of the authors or their affiliated institutions, 2024.